An Analysis on the Determination of Climatic Drought Indices and Zoning in Khorasan Razavi Province

Mehdi boroghani

Watershed engineering PhD student in Faculty of Nor Natural Resource, Tarbeat Modares University, Mazandaran Province, Nor city, Faculty of Natural Resources.

hamidreza Moradi*

Associate professor of Watershed Engineering Department in Natural Resource Faculty of Nor, Tarbeat Modares University, Mazandaran province, Nor city, Faculty of Natural Resource, Department of Watershed Engineering.

mohamadali Zangane asadi

Associate professor of Geomorphology, Geography Faculty of Hakim Sabzevari University, Sabzevar, Tovhid Shar, Hakim Sabzevari University, Geographic Faculty.

Extended Abstract

Introduction

Drought is a creeping environmental phenomenon that has many damaging effects on the economy, agriculture and society. Drought is a situation of shortage of rainfall and incoming temperature increases that may occur in any climate. This phenomenon is in parts of eastern central and southern of Iran, due to dry and semidry weather will be more. The geographic distribution of droughts occurred in the percentage of the country shows that the southern and eastern areas of the country's is more extensive, and whatever from the southern and eastern parts of the country took distance the intensity and frequency of drought is also reduced. The reason for this is the impact of subtropical high-pressure systems that significant amount of the rainforest of South and Eastern parts in the country towards the northern and Western parts of the significantly reduced and prevents the effect of Northern and Western systems in these area.

Methodology

Khorasan Razavi with 143864.118 Km² is the fourth largest province in Iran. The highest point of the province is in the North of Nishabor with a height of Binalud heights 2116 m above sea level and the lowest point of the province is Sarakhs plain with a height of 300 meters above sea level which is located in Iran and Turkmenistan border. The average annual precipitation of Khorasn Razavi is between 75 to 390 mm. At the first data related to precipitation in the Razavi Khorasan Province was obtained from the country's water resources management organization. Precipitation data from 1980 to 2010 were averaged on an annual basis. This data is them prepared in note pad file, and by using DIP software, the index of SPI, PNI and ZSI was calculated for all situations. After the calculation of this indicator for each year, the data were entered into ArcGIS software and then used ordinary Kriging method for drought zoning for the end of the 11 years (2000-2010).

Discussion and result

The results indicated that changes in wet and drought year during the studied period didn't follow a regular process but changes in indicator are similar. Results also indicate that the year of 1992 and 2008 respectively are the rainiest and driest year during the studied period in all three indicators.

The results of drought indices zoning of PNT, ZSI and SPI in the period of case study suggest that changes in different years and parts of Province does not follow the regular process. The result of drought index zoning in Khorasan Razavi Province suggested that PNI drought index made the condition wetter and ZSI drought index drier. Also the results show that drought index SPI and ZSI

^{*}Email: hrmoradi@modares.ac.ir

are more consistent than PNI drought index. The result of drought index PNI, ZSI and SPI indicated that in 2008, 2001 and 2000 years respectively are the driest zoning year. SPI and ZSI drought index in 2003 were the wettest year drought zoning and drought index in 2003 in addition to 2004 and 2009 were the wettest year. API drought index calculated the minimum amounts of rainfall in the studied station in the years of happening dryness, which regarding two other indexes are more accuracy. The result also shows that ZSI and PNI index have the same accuracy. In total SPI drought index for dry zoning in Khorasan Razavi province is more accurate and this index is suggested for drought studies.

Conclusion

The results indicated that none of the index drought zoning followed the regular processes. The results suggested that SPI index is the best performance of other indices. Also PNI and ZSI indices have the same accuracy in determination of drought in Khorasan Razavi Province. This result had compliance with the results of Naserzade and Ahmadi (2012), Ansafi Moghadam (2007), Karimi et al (2011), Piry et al (2013) and Khoram bakht et al (2014)which in their studies determined SPI index as the best index. Besides the results didn't match with the result of Shokohi (2012) which SPI index was not the best index.

References

- 1- Ansafi Moghadam, T., "Evaluation of several climatic droughts index and determine the most appropriate indicator in the salt lake basin fitted", Iranian Journal of research in rangeland and desert, 14(2): 271-288. (In Persian)
- 2- Beheshti Rad, M., beheshti Rad, M., 2013, "Reviews of the performance of methods of zoning of the severity of the drought in the province of Kerman", Quarterly regional planning, 9(1): 89-101. (In Persian)
- 3- Boroghani, M., Taie, M., Mirnia, S.K., 2013, "Analysis of relationship between hydrogeological and climatological droughts using SWI and SPI indices in Sabzevar Plain", quarterly of Scientific-research Iranian desert and rangeland research, 20(4): 733-744. (In Persian)
- 4- Edwards, D.C., McKee, T.B., 1997. "Characteristics of 20th century drought in the United States at multiple timescales", Colorado State University: Fort Collins. Climatology Report No. 97-102.
- 5- Gholam Ali, M., Younes, K.H., Esmaeil, A., Fatemeh, T., 2011. Assessment of Geostatistical Methods for Spatial Analysis of SPIand EDI Drought Indices, World Applied Sciences Journal, 15 (4):474-482.
- 6- Goovaerts, p., 2000. Geostatistical approaches for incorporating elevation into the spatial interpolation of rainfall, Journal of Hydrology, 228: 113-129.
- 7- Herweijer, C., Seager, R., Cook, K., Geay, E., 2013. North American Droughts of the Last Millennium from a Gridded Network of Tree-Ring Dat, Lamont-Doherty Earth Observatory, Columbia University, Palisades, New York, Drying Technology: An International Journal, Volume 31, Issue 15.
- 8- Herweijer, C., Seager, R., 2008. The global footprint of persistent extra-tropical drought in the instrumental era, International Journal of Climatology, 28(13):1761-1774
- 9- Hong, W and Hayes, J., 2001. An Evaluation of the standardized precipitation Index, The China-Z Index and the statistical Z-Score, Journal of Climatol, 21:745-758.
- Hong, W., Hayes, M.J., Welss, A., Hu, Q., 2001. An evaluation the standardized precipitation index, the china-z index and the statistical z-score, International Journal of Climatology, 21:745-758.
- 11- Hoseini Safa, H., Marid, S., 2008, "Probabilistic analysis of spatial characteristics of the severity of the drought in the province of Tehran", Journal of agricultural sciences and resources, 11(5): 203-212. (In Persian)

- 12- Karimi, V., Habib nejad, M., Abkar, A., 2011, "Reviews of meteorological drought index in Mazandaran synoptic stations", Journal of irrigation and water engineering, 5(3): 15-25. (In Persian)
- 13- Keyantash, J.A and Dracup, J.A., 2004. An aggregate drought index: assessing drought severity based on fluctuation in the hydrologic cycle and surface water storage, water resources research, 40: 1-13.
- 14- Khalili, A., bazr Afshan, J., 2003, "the evaluation of the performance of meteorological drought index in a few examples of the different climate of Iran", Journal of Nivar, 48(1): 79-93. (In Persian)
- 15- Khoram Bakht, A., Moshiri, S.R., Mahdavi, M., 2014, "Evaluation of the climatic drought in the larestan region attributes", Quarterly regional planning, 13(1): 103-120. (In Persian)
- 16- Lloyd-Hughes, B., Saunders. M.A., 2002. "A drought climatology for Europe", International Journal of Climatology 22: 1571–1592.
- 17- Loukas, A., Vasiliades, L.N.R., Dalezios, N.R., 2003. Intercom parison of meteorological drought indices for drought assessment and monitoring in Greece. Proceeding of the 8 international conferences on environmental science and Technology. Lemons Island and Greece 8-10 September 2003.
- 18- Mirzaie, A.A., Sbah, Gh., 2011, "Specialized software for water engineering", Publishing Kian green computer, First printing. (In Persian)
- 19- Moradi, H.R., Rajabi, M., Faragzadeh, M., 2011. Investigation of meteorological drought characteristics in Fars province, Iran, Catena, 84(2): 35-46.
- 20- Mosavi Baygi, M., Ashraf, B., 2011, "the study of the patterns of synoptic leads to winter and autumn droughts in Khorasan Razavi Province". Journal of soil and water conservation research, 18(4): 167-184. (In Persian)
- 21- NaserZade, M.H., Ahmadi, A., 2012, "A study on the performance of meteorological drought indices in evaluation of drought and its regionalization in Qazvin province", Journal of geographical sciences, applied research, 27(2): 141-162. (In Persian)
- 22- Nikbakht, J., Tabari, H., Talaee, P., 2013. Stream flow drought severity analysis by percent of normal index (PNI) in northwest Iran, Theoretical and Applied Climatology. 112: 565-573.
- 23- Palmer, W.C., 1965. "Meteorological Drought", U.S. Department of Commerce Weather Bureau, Washington, D.C. Research Paper No. 45.
- 24- Palmer, W.C., 1968. Keeping track of crop moisture conditions, nationwide: the new Crop Moisture Index, Weatherwise, 21: 156-161.
- 25- Peri, H., Abas Zade, M., Rahdari, V., Malki, S., 2013, "Comparative evaluation of four meteorological drought indexes using the method of cluster analysis (case study of Sistan and Baluchestan province)", Journal of water resources engineering, 6(2): 25-36. (In Persian)
- 26- Razeei, T., DaneshKar Araste, P., Akhtari, R.A., Saghafean, B., 2007, "Reviews of meteorological droughts (climate) in the province of Sistan and Baluchestan with use SPI and Markov chain model", Iranian Journal of research in water resources, 3(1): 25-35. (In Persian)
- 27- Salehvand, A., Montazeri, M., Momeni, M., 2013, "drought zoning with (ZSI, CZI, DI, PNI and SPI) and the sequence in the cities in Khuzestan province in GIS software", Journal of geography and urban planning the Zagros landscape, 17(3): 35-52. (In Persian)
- 28- Shafer, B.A.,Dezman, L.E., 1982. Development of a Surface Water Supply Index (SWST) to assess the severity of drought conditions in snow pack runs off areas, In Proceedings of the (50th) Annual Western Snow Conference, 164-175.
- 29- Shokohi, A., 2012, "comparing SPI and RDI indexes for analyses of drought in station scale with use agricultural drought (case study: Ghazvin and takestan)", Journal of irrigation and water engineering, 9(3): 111- 122. (In Persian)

- 30- Tsakiris, G., Vangelis, H., 2004. Towards a drought watch system based on spatial SPI, Water Resources Management, 18: 1-12.
- 31- Yazdani, V., Davari, K., Ghahraman, B., Zare Abyane, H., 2013, "the meteorological drought index four validated based on growth rings of trees wood (case study Mashhad)", Water engineering magazine, 23-37.