# Assessing Changes in Patterns of Spatial Autocorrelation of Maximum Temperature of Iran

Kazem Aliabadi<sup>\*</sup> Geography Department, Hakim Sabzevari University, Sabzevar, Iran AbbasAli Dadashi Roudbari Ph.D. Student Urban Climatology, Shahid Beheshti University, Tehran, Iran

**Extended Abstract** 

## Introduction

Temperature is one of the important elements of weather, resulting in part from solar energy absorbed by the effects of the Earth's surface and is converted to thermal energy. Maximum temperatures in the hot period of the year, especially in the hours after the city, should be given more attention. Analyzing the maximum temperature can also be normal evaluation mechanism to evaluate changes and track climate change. This study aimed to identify the spatio-temporal patterns and determine the maximum monthly patches of country, so the main focus of the maximum temperature will be identified.

### **Research Methodology**

The common statistics and homogeneous 125 synoptic stations with a common statistic period of 30 years (2010-1980) is used, then the station data using kriging interpolation software in the area of data with ArcGIS 10.2.2 of cells to dimensions  $15 \times 15$  km was extended. In this study for investigation of the spatial structure of cells, maximum temperature coefficient of dispersion country is evaluated. As well as for investigation the pattern of the maximum temperature of Iran, the new methods of spatial statistics, spatial autocorrelation (Moran World) and cluster analysis Clusters and Outlier, shot spots to study the temporal and spatial patterns and changes in the maximum temperature is used.

#### **Results and discussion**

The results showed that the highest maximum temperature of spatial anomalies in January, February and March has happened. The result of Moran World trend proves it that changes within a year of maximum temperatures of Iran followed the cluster model and on the bases of output of statistics follows the pattern of clustering of Iran with the maximum temperatures happened in March with the amount of 0/973876. Reviews of the structure's temperature using clusters and outlier showed that in the cold months decrease maximum temperature uniformity and homogeneity in the hot summer months the maximum temperature will increase. In the meantime, based on hot spots, the maximum temperature in the coastal shores of the Caspian Sea, parts of the West and North West and North East of the country has negative spatial autocorrelation (clusters of low value) and part of the central region as well as parts the South East and central regions has positive spatial autocorrelation (clusters with high-value, respectively).

#### Conclusion

Based on the analysis of spatial autocorrelation, spatial autocorrelation negative areas every 12 months relates on parts of the North West, North East and the Zagros bar. Areas with positive spatial autocorrelation often located on South coast of Persian Gulf and Oman respectively. Every 12 months, the upper area of the country has no significant pattern of spatial autocorrelation. Spatial analysis showed that the maximum temperature patterns of the two spots for the maximum temperature of in the form of an island faced the South East, South, South-West and Central Iran. Low cluster areas mostly based on Zagras bar area, North West, North East and the North Coast. These areas formed the core of certain areas of the country's hot and cold. The results showed that

<sup>\*</sup> Email: Kazemaliabadi\_msc@yahoo.com

latitude and altitude of the country is evident in the distribution of the maximum temperature. Results of this study with some of the previous findings in which the country's maximum temperatures were examined (Alijani, 2008: 50-64; Masoudian and kavyani, 2008: 53-66; Mojarad and Bastami, 2014: 129-152) have conformity.

**Keywords**: maximum temperatures, global Moran, hot spot analysis, spatial autocorrelation, Iran.

#### References

- 1. Ageena, I., Macdonald, N., & Morse, A. P. (2013). Variability of maximum and mean average temperature across Libya (1945–2009). Theoretical and Applied Climatology, 1-15.
- 2. AGHABOZORGI, s., SAYBANI m., WAH, t. (2012), Incremental Clustering of Time-Series by Fuzzy Clustering, JOURNAL OF INFORMATION SCIENCE AND ENGINEERING 28, 671-688 (2012).
- 3. Alijani, b. (2008), Climate of Iran, eighth edition, the publisher of Payam Noor University, Tehran (In Persian).
- 4. Alijani, b., Freedom, M. (1992), Principles of Climatology, the publisher, Tehran (In Persian).
- 5. Anderson, G. B., Bell, M. L., & Peng, R. D. (2013). Methods to calculate the heat index as an exposure metric in environmental health research. Environmental health perspectives, 121(10), 1111.
- 6. Anselin L (1995). Local indicators of spatial association: LISA. Geogr Anal 27(2):93-115.
- Anselin L, Syabri I, Kho Y (2009) GeoDa: an introduction to spatial data analysis. In Fischer MM, Getis A (eds) Handbook of applied spatial analysis. Springer, Berlin,Heidelberg and New York, pp.73-89.
- 8. Azizi, and., Roshani, d. (2008). Study of climate change on the southern coast of the Caspian Sea to the Mann-Kendall method, Geographical Research, No. 64, pp. 13-28 (In Persian).
- Bajat, B., Blagojević, D., Kilibarda, M., Luković, J., & Tošić, I. (2014). Spatial analysis of the temperature trends in Serbia during the period 1961–2010. Theoretical and Applied Climatology, 1-13.
- 10. Brovelli, M., A, and Liu, X (2009). GIS3 Exercise- Spatial Interpolation, GIS and Theory of Errors Group.
- Burns, D. A., Klaus, J., & McHale, M. R. (2007). Recent climate trends and implications for water resources in the Catskill Mountain region, New York, USA. Journal of Hydrology, 336(1), 155-170.
- 12. David, F. N., & Moore, P. G. (1954). Notes on contagious distributions in plant populations. Annals of Botany, 18(1), 47-53.
- 13. De Lucena, A. J., Rotunno Filho, O. C., de Almeida França, J. R., de Faria Peres, L., & Xavier, L. N. R. (2013). Urban climate and clues of heat island events in the metropolitan area of Rio de Janeiro. Theoretical and applied climatology, 111(3-4), 497-511.
- 14. Diffenbaugh, N. S., J. S. Pal, F. Giorgi, and G. Xuejie (2007), Heat stress intensification in the Mediterranean climate change hotspot, Geophys. Res. Lett., 34, L11706, doi: 10.1029/2007GL030000.
- 15. Diggle, P.J., (2003). Statistical Analysis of Spatial Point Patterns, second ed. Academic Press, London.GCOS, 2007. GCOS Upper-Air Network (GUAN): Justification, Requirements, Siting and Instrument Options. GCOS-112, WMOTD 1379.
- 16. Douglas, J. B. (1975). Clustering and aggregation. Sankhyā: The Indian Journal of Statistics, Series B, 398-417.
- 17. El Kenawy, A., López-Moreno, J. I., & Vicente-Serrano, S. M. (2012). Trend and variability of surface air temperature in northeastern Spain (1920–2006): linkage to atmospheric circulation. Atmospheric Research, 106, 159-180.
- 18. Founda, D., Papadopoulos, K. H., Petrakis, M., Giannakopoulos, C., & Good, P. (2004). Analysis of mean, maximum, and minimum temperature in Athens from 1897 to 2001

with emphasis on the last decade: trends, warm events, and cold events. Global and Planetary change, 44(1), 27-38.

- 19. Getis A, Aldstadt J (2004) Constructing the spatial weights matrix using a local statistic. Geogr Anal 36(2):90-104.
- 20. Getis A, Ord JK (1992), the analysis of spatial association by use of distance statistics. Geogr Anal 24(3):189-206.
- 21. Green, R. H. (1966). Measurement of non-randomness in spatial distributions. Researches on Population Ecology, 8 (1), 1-7.
- 22. Griffith, D., (1987), spatial Autocorrelation: A Primer. Resource Publication in Geography, Association of American geographers.
- 23. Hassanipak. a (2007). geostatistics (geostatistical), Tehran University Press, second edition, Tehran (In Persian).
- 24. Hengl, T (2006). Finding the right pixel size. Computers & geosciences, 32 (9).
- 25. Holdaway, M. R. (1996). Spatial modeling and interpolation of monthly temperature using kriging. Climate Research, 6(3), 215-225.
- 26. Homar V, Ramis C, Romero R, Alonso S (2010) Recent trends in temperature and precipitation over the Balearic Islands (Spain). Clim Change 98:199–211. Doi: 10.1007/s10584-009-9664-5.
- 27. Illian, J., Penttinen, A., Stoyan, H., and Stoyan, D. (2008). Statistical Analysis and Modelling of Spatial Point Patterns. John Wiley and Sons, Chichester.
- 28. Jacquez GM, Greiling DA (2003) Local clustering in breast, lung and colorectal cancer in Long Island, New York. Int J Health Geographics 2:3.
- 29. Johnston, K, Ver Hoef, J.M, Krivoruchko, K, Lucas, N, and Magri, A (2003). ArcGIS, Geostatistical Analyst, ESRI, USA.
- Kendall, W. S. (1998). Perfect simulation for the area-interaction point process. In L. Accardi and C.C. Heyde, editors, Probability Towards 2000, pages 218–234. Springer Lecture Notes in Statistics 128, Springer Verlag, and New York.
- Kim, S., & Singh, V. P. (2014). Modeling daily soil temperature using data-driven models and spatial distribution. Theoretical and Applied Climatology, 1-15.
- Kuglitsch, F. G., A. Toreti, E. Xoplaki, P. M. Della-Marta, C. S. Zerefos, M. Türkeş, and J. Luterbacher (2010), Heat wave changes in the eastern Mediterranean since 1960, Geophys. Res. Lett., 37, L04802, doi:10.1029/2009GL041841.
- Levine N (1996) spatial statistics and GIS: software tools to quantify spatial patterns. JAm Plann Assoc 62(3):381-391.
- 34. Lloyd, M. (1967). Mean crowding. The Journal of Animal Ecology, 1-30.
- 35. Masoodian, S. A (2008). On Precipitation Mapping in Iran, Journal of Humanities The University of Isfahan, Vol 30, No 2, 69-80.
- 36. Masoudian, SA, Kavyani, d. (2008), Climate of Iran, Isfahan University, Isfahan, Iran (In Persian).
- 37. Masoudian, SA. (2005), the trend of temperature in the past half century, Geographical Research, No. 54, Ss29-45 (In Persian).
- 38. Masoudian, SA. (2011), weather Iran, Mashhad by Sharia publishing, printing, Mashhad (In Persian).
- 39. Mitchell A (2005) The ESRI guide to GIS analysis, volume 2: spatial measurements and statistics. ESRI, Redlands [CA].
- 40. Mohammad Rezai, Sh. (2002), systematic approach to analyze ecosystem, Environmental Protection Agency, Tehran (In Persian).

- 41. Mojard, F., Bastami, s. (2014), Assessment temporal and spatial changes in maximum temperatures of Iran, professor of human Spatial Planning, Volume XVIII, No. 2, Ss129-159 (In Persian).
- 42. Moller, j. (2008) Handbook of Spatial Statistics, John Wiley and Sons, Chichester.
- 43. Moreno, J. I. L., and Bravo, D. N (2006). Interpolating Local Snow Depth Data: An Evaluation of Methods, Hydrological Processes, 20.
- 44. Morisita, M. (1959). Measuring of the dispersion of individuals and analysis of the distributional patterns. Mem. Fac. Sci. Kyushu Univ. Ser. E, 2(21), 5-235.
- 45. Nel W.P., Cooper C.J., (2009), Implications of fossil fuel constraints on economic growth and global warming, Energy Policy, 37, 166-180.
- 46. Nemec, J., Gruber, C., Chimani, B., & Auer, I. (2013). Trends in extreme temperature indices in Austria based on a new homogenised dataset. International Journal of Climatology, 33(6), 1538-1550.
- 47. Ord JK, Getis A (1995) Local spatial autocorrelation statistics: distributional issues and anapplication. Geogr Anal 27(4):287-306.
- 48. Rahim Zada, F., Askari, or. (2004). a view on the difference between the minimum and maximum rate of increase in temperature and decrease the temperature in the country Boarding, Geographical Research, No. 73, pp. 155-171 (In Persian).
- 49. Ripley, B. D. (1977). Modelling spatial patterns. Journal of the Royal Statistical Society. Series B (Methodological), 172-212.
- 50. Rogerson, P.A., (2006), Statistics Methods for Geographers: students Guide, SAGE Publications. Los Angeles, California.
- 51. Sonali, P., & Nagesh Kumar, D. (2013). Review of trend detection methods and their application to detect temperature changes in India. Journal of Hydrology, 476, 212-227.
- 52. Waagepetersenand, R., and Schweder, T. (2006). Likelihood-based inference for clustered line transect data. Journal of Agricultural, Biological, and Environ- mental Statistics, 11:264–279.
- 53. Wheeler D (2007) A comparison of spatial clustering and cluster detection techniques for childhood leukemia incidence in Ohio, 1996-2003. Int J Health Geographics 6(1):13.
- 54. Wheeler D, Paéz A (2009) Geographically Weighted Regression. In Fischer MM, Getis A (eds) Handbook of applied spatial analysis. Springer, Berlin, Heidelberg and New York, pp.461-486.
- 55. Wu, T., & Li, Y. (2013). Spatial interpolation of temperature in the United States using residual kriging. Applied Geography, 44, 112-120.
- 56. Zhang C, Luo L, Xu W, Ledwith V (2008) Use of local Moran's I and GIS to identify pollution hotspots of Pb in urban soils of Galway, Ireland. Sci Total Environ 398 (1-3):212-221.