

Original Article

Patterns of Glycoconjugate Distribution during Molar Tooth Germ Development in Mice

T. Kermani¹✉, AR. Varasteh¹

¹Assistant Professor, Department of Anatomy, School of Medicine, Birjand University of Medical Sciences, Birjand, Iran

²Associate Professor, Immunological Research Center, Bu-Ali Research Institute; Mashhad University of Medical Sciences, Mashhad, Iran

Abstract:

Objective: The aim of the present study was to evaluate the structure and distribution of Glycoconjugates during molar tooth germ development in mice

Materials and Methods: Sixteen tooth germs were obtained from BALB/c mice embryos 15 to 18 days post-gestation and fixed in 10% formalin. After routine tissue processing, 5 µm sections were cut and stained with BSA1-B4 and PNA using the lectin histochemical method. All slides were evaluated by light microscopy.

Results: Both lectins showed positive reaction in the tooth germ but with spatiotemporal differences. During bell stage, the reaction was strong with BSA1-B4 but moderate with PNA. Strong PNA uptake was observed in the odontoblastic and ameloblastic nuclei along with the apical cytoplasm of the ameloblasts.

Conclusion: Although the lectins that were used in the present study recognize the same terminal sugar residue, they reacted with different disaccharide sequences with various penaltomer sugars. Therefore it may be assumed that the pattern of affinity for different parts of the developing tooth germ such as ameloblasts and odontoblasts is different in various lectins.

Key Words: Development; Glycoconjugates; Lectin; Tooth

Journal of Dentistry, Tehran University of Medical Sciences, Tehran, Iran (2007; Vol: 4, No.3)

✉ Corresponding author:
T. Kermani, Department of
Anatomy, School of Medicine,
Birjand University of Medical
Sciences, Birjand, Iran.
kermani_t@bums.ac.ir

Received: 20 August 2006
Accepted: 20 December 2006

INTRODUCTION

The developing tooth is an excellent model to study the molecular mechanisms of morphogenesis. It is well known that tooth morphogenesis and differentiation of the dental cells are guided by interactions between epithelial and mesenchymal tissues [1-7]. Dental epithelium in mice is capable of inducing tooth formation just before the bud stage, whereas its tooth-forming potential relocates from the epithelium to the dental mesenchyme. [8-11]. However, the molecular mechanisms involved in dental cell differentiation are not well understood.

One of the critical molecules involved in em-

bryonic development is glycoconjugates that are detected by lectins. Glycoconjugates have important roles in cell-cell interaction, cell migration and proliferation [12-17]. Lectins are glycoproteins that can specifically react with terminal sugars. 'Lectin' comes from the Greek word 'legere', which means 'to select'. Most lectins act in a non-enzymatic manner and demonstrate a non-immune origin. Lectins are constantly encountered in nature [18]. These molecules may bind to a carbohydrate moiety occurring free in a solution or as a part of a protein/particulate body and can agglutinate cells and/or precipitate glycoconjugates. Lectins extracted from plants can conjugate




Fig 1. Cap stage; BSA-1 Reaction; EO; Enamel Organ, IEE; Inner Enamel Epithelium, OEC; Oral Epithelium Cells, SR; Stellate Reticulum × 200




Fig 2. Cap stage; PNA Reaction: EO; Enamel Organ, IEE; Inner Enamel Epithelium, OEC; Oral Epithelium Cells, SR; Stellate Reticulum × 200

with labels and are good tools to detect terminal sugars in different tissues [12-13]. These molecules have the potential to participate in both differentiation and maturation [19]. Many studies have been performed to describe the distribution of oligosaccharide chains in developing teeth [12-16,19,20]; however, the distribution pattern of these molecules shows a considerable amount of diversity in developing and tumoral tissues [12-16,21,22]. Therefore, it seems that further investigation can be helpful to elucidate the roles of these sugar residues in mouse models. The aim of the present study was to evaluate the structure and distribution of Glycoconjugates during molar tooth germ development in mice.

MATERIALS AND METHODS

BALB/c mice with an age range of 2-4 months, weighing approximately 25-30 grams were used in the present study. Males and females were caged together overnight. Observa-

tion of the vaginal plug was considered as day zero of gestation. The pregnant mice were kept at standard conditions with free access to water and food. Sixteen mice embryos were collected from the 15th to 18th days of gestation. Their heads were fixed, processed and sectioned serially.

Lectin Histochemistry:

The Lectin histochemical method was used for detecting glycoconjugates. Deparaffinized and dehydrated 5 µm sections were immersed in 0.1 M PBS containing 0.1mM CaCl₂, MgCl₂ and MnCl₂ (PBSc), followed by 1% H₂O₂ in methanol for 10-15 minutes. After rinsing with PBSc, they were incubated with peroxidase conjugated lectins including PNA (L7381; Sigma, USA) and BSA1-B4 (L5391; Sigma, USA) that bind galactose (Gal) and N acetyl galactoseamine (Gal-Nac) for two hours at room temperature, respectively. The lectins were diluted to a concentration of 10µg/ml [23,24].

Table 1. Lectin reactivity of the cells and ECM in the tooth germ at cap stage

Lectins	Oral Epithelial Cells	Basement Membrane	Enamel Organ	Ecto-mesenchyme	Dental Sac	Extracellular Matrix	Capillary Endothelium
BSA-1	+++ *	-	+++	+++	+++	-	-
PNA	-	-	+	-	-	-	-

* Evaluation of binding indicates staining intensity on a subjectively estimated scale: no staining (-), little staining (+), moderate staining (++), intense staining (+++)




Fig 3. Bell stage; BSA-1 Reaction: A; Ameloblasts, O; Odontoblasts, pd; predentine, DP; Dental Papilla, SR; Stellate Reticulum × 400




Fig 4. Bell stage; PNA Reaction: A; Ameloblasts, O; Odontoblasts, pd; predentine, DP; Dental Papilla, SR; Stellate Reticulum, Pe; Preenamel × 400

Following final washing with buffer, the binding sites were visualized by incubating sections in 0.03% diaminobenzidine containing 200µl H₂O₂ in PBS for 10 minutes. The sections were counterstained with alcian blue. To compare the intensity of positive staining, we used an arbitrary scale from 0 to 4, ranging from no- to strong reaction [25]. It has been shown that retinal epithelium and endothelial cells in different tissues can react with PNA [26] and BSA1-B4 [27], respectively; therefore a composite of several tissues including retinal pigmented epithelium, testis and ... were used as positive control. Negative control was obtained by omitting lectins in a number of slides in order to establish negative reaction. The study protocol was approved by the Ethics committee of Birjand University of Medical Sciences.

RESULTS

A positive reaction for both Lectins was identified in the studied tooth germ, but with spatiotemporal differences. BSA1-B4 reaction occurred during the cap stage with “strong” intensity in the oral epithelium, enamel organ and ectomesenchyme which was limited to the cells (Fig 1). The extracellular matrix (ECM) began to take-up lectin at the bell stage. The

predentin and odontoblasts of the apical region were more reactive to BSA1-B4 in this stage. In addition to the thick amorphous matrix under the ameloblasts, the dental papilla and the endothelial cells also demonstrated lectin uptake (Fig 2).

Oral epithelial cells showed positive reaction to PNA, but the basement membrane was negative. The enamel and ectomesenchyme were also positive but stained “weakly” for this lectin (Fig 3). At the bell stage a “strong” reaction to PNA was observed in the Golgi zone of the odontoblasts and ameloblasts in addition to the apical cytoplasm of the ameloblasts (Fig 4). Lectins have been suggested to label the Golgi compartment of cells in odontogenic tissues. The thick amorphous matrix under the ameloblasts was PNA positive but the predentin was negative except for a thin layer of newly formed dentin adjacent to the ameloblasts. Dispersed reaction was seen in the dental papillae. Lectin reactivity in different stages of tooth development is shown in Tables 1 and 2.

DISCUSSION

Glycoconjugates have been suggested to participate in many critical events of embryogenesis by establishing the dental morphologic

Table 2. Lectin reactivity of the cells and ECM in the tooth germ at bell stage.

Lectins	Ameloblasts			Odontoblasts			Pre-dentine	Dental Papilla	Capillary Endothelium
	Apical area	Nucleus	Base	Apical area	Nucleus	Base			
BSA-1	-	+	+++	+++	+++	-	+++	++	+++
PNA	+++	++	+++	-	++	-	-	+	+

no staining (-), little staining (+), moderate staining (++) , intense staining (+++)

pattern and ECM secretion [19]. They regulate cell-cell and cell-matrix interactions [8-17]. Our data indicated differential expression of lectin residues in various parts of the tooth germ that highlights the importance of these sugar residues in tooth development.

In the present study, PNA positive residues were found in ameloblasts which is in accordance with a study conducted by Lemous et al [14]. It has been shown that Peanut lectin influences epithelial proliferation in some cancers [21,22]. PNA can affect normal pancreatic growth [28] and PNA residues may also have the same function in regulation of cell proliferation during tooth development. In addition it has been suggested that PNA binding structures may be important for cellular interaction during morphogenetic processes. Ameloblasts have an inductive effect on the dental papilla resulting in the formation of odontoblasts. PNA reacted residues may be related to these types of interactions.

It has been demonstrated that an N-acetylgalactosamine rich matrix component is differentially expressed during odontogenesis [16-17]. However, according to the results obtained in the current investigation, predentin reaction to PNA was negative. Proteoglycans like D-galactose and N-acetylgalactosamine sugars bound to PNA are considered as important components of the dentin matrix. The production of these sugars in adult rat odontoblasts is related to PNA binding sites [29]. This was in contrast to the results obtained in the current investigation regarding PNA reaction in mouse tooth germs. A possible explanation could be that odontoblast secretion is dif-

ferent after tooth eruption; therefore considering that the present study was performed on mouse embryos, we did not find reactive residues of PNA in the predentin.

Growth hormone can regulate the glycoconjugate content of the extracellular matrix that is essential for normal tooth development [16], and becomes functional after birth. Various PNA expression patterns may be related to the effects of growth hormone.

According to our results, BSA1-B4 stained the cell surfaces in the cap and early bell stages. Moderate reaction to galactose has been reported in other studies [15]. The presence of glycoconjugates in calcified tissues like dentin may be related to mineral deposition. It has been suggested that highly charged plasma membranes produced by glycocalyxes are responsible for the transportation of mineral and/or organic materials between ameloblasts and extracellular fluid that is organized by glycoconjugates. [12]. Amelogenins are major enamel proteins and their properties are similar to lectins, hence they can bind to glycoconjugates during enamel formation [30].

CONCLUSION

The presence or absence of the glycoconjugates may have a critical role in tooth development because of their spatiotemporal distribution. However, more investigation is required in order to gain a better understanding of their molecular mechanisms.

ACKNOWLEDGMENT

The authors wish to thank the Research Deputy of Birjand University of Medical Sciences

and Immunology Research Lab of Bu-Ali Research Center of Mashhad University of Medical Sciences.

REFERENCES

- 1-Felszeghy S, Mészár Z, Prehm P, Módis L. The expression pattern of hyaluronan synthase during human tooth development. *Arch Oral Biol* 2005 Feb;50(2):175-9.
- 2-Lesot H, Kieffer-Combeau S, Fausser JL, Meyer JM, Perrin-Schmitt F, Peterková R et al. Cell-cell and cell-matrix interactions during initial enamel organ histomorphogenesis in the mouse. *Connect Tissue Res* 2002;43(2-3):191-200.
- 3-Hall RC, Embery G. The use of immunohistochemistry in understanding the structure and function of the extracellular matrix of dental tissues. *Adv Dent Res* 1997 Nov;11(4):478-86.
- 4-Thesleff I, Partanen AM, Vainio S. Epithelial-mesenchymal interactions in tooth morphogenesis: the roles of extracellular matrix, growth factors, and cell surface receptors. *J Craniofac Genet Dev Biol* 1991 Oct-Dec;11(4):229-37.
- 5-Slavkin HC. Molecular determinants during dental morphogenesis and cytodifferentiation: a review. *J Craniofac Genet Dev Biol* 1991 Oct-Dec;11(4):338-49.
- 6-Thesleff I, Stenman S, Vaheri A, Timpl R. Changes in the matrix proteins, fibronectin and collagen, during differentiation of mouse tooth germ. *Dev Biol* 1979 May;70(1):116-26.
- 7-Slavkin HC. Embryonic tooth formation. A tool for developmental biology. *Oral Sci Rev* 1974;4(0):7-136.
- 8-Thesleff I, Vaahtokari A, Partanen AM. Regulation of organogenesis. Common molecular mechanisms regulating the development of teeth and other organs. *Int J Dev Biol* 1995 Feb;39(1):35-50.
- 9-Yamamoto H, Kim EJ, Cho SW, Jung HS. Analysis of tooth formation by reaggregated dental mesenchyme from mouse embryo. *J Electron Microscop* (Tokyo). 2003;52(6):559-66.
- 10-Hu B, Nadiri A, Bopp-Kuchler S, Perrin-Schmitt F, Wang S, Lesot H. Dental epithelial histomorphogenesis in the mouse: positional information versus cell history. *Arch Oral Biol* 2005 Feb;50(2):131-6.
- 11-Lodish H, Baltimore D, Berk A, Zipursky SL, Matsudaiva P, Darnell S. *Molecular cell biology*. 3rd ed. Scientific American Books. 1995; p.1123-1200.
- 12-Nakamura H, Ozawa H. Lectin cytochemistry on the stratum intermedium and the papillary layer in the rat incisor enamel organ. *Arch Histol Cytol* 1990 Oct;53(4):351-69.
- 13-Lemus D, Lemus R, Romero S, Arancibia N, Fuenzalida M. Detection of sugar residues in rabbit embryo teeth with lectin-horseradish peroxidase conjugate: II. A light microscopical study. *J Morphol* 1997 Feb;231(2):175-84.
- 14-Lemus D, Romero S, Lemus R, Garcia J, Fuenzalida M. Light microscopic detection of sugar residues in rabbit embryo teeth with lectin-horseradish peroxidase conjugates. *J Morphol* 1996 Feb;227(2):185-95.
- 15-Jowett AK, Kimber SJ, Ferguson MW. Immunofluorescent lectin binding patterns and glycoprotein co-localization in the developing murine molar tooth. *Arch Oral Biol* 1992 Apr;37(4):303-14.
- 16-Zhang CZ, Young WG, Breipohl W, Doehn S, Li H, Waters MJ. Growth hormone regulates an N-acetylgalactosamine component in odontogenesis: a specific lectin-binding study in the Lewis dwarf rat. *J Oral Pathol Med* 1994 May;23(5):193-9.
- 17-Fazel AR, Schulte BA, Spicer SS. Glycoconjugate unique to migrating primordial germ cells differs with genera. *Anat Rec* 1990 Oct;228(2):177-84.
- 18-Goldstein IJ, Hughes RC, Monsigny M, Osawa T, Sharon N. What should be called a lectin? *Nature*. 1980 May;285:66.
- 19-Jiménez-Farfán D, Guevara J, Zenteno E, Hernández-Guerrero JC. Alteration of the sialylation pattern of the murine tooth germ after ethanol exposure. *Birth Defects Res A Clin Mol Teratol*. 2005 Dec;73(12):980-8.
- 20-Kermani T, Varasteh AR, Nicravesh MR, Moradi M. Study of glycosaminoglycans of extracellular matrix (ECM) in pulp of developing tooth.

- Journal of Dental Medicine Tehran University of Medical Sciences 2000; 13(2): 52-9.
- 21-Singh R, Subramanian S, Rhodes JM, Campbell BJ. Peanut lectin stimulates proliferation of colon cancer cells by interaction with glycosylated CD44v6 isoforms and consequential activation of c-Met and MAPK: functional implications for disease-associated glycosylation changes. *Glycobiology* 2006 Jul;16(7):594-601.
- 22-Valentiner U, Fabian S, Schumacher U, Leathem AJ. The influence of dietary lectins on the cell proliferation of human breast cancer cell lines in vitro. *Anticancer Res* 2003 Mar-Apr;23(2B): 1197-206.
- 23-Hounsell EF. Glycoprotein analysis in biomedicine series: Methods in molecular biology V.14, Humana Press Inc. Totowa Chapter 20, p. 247-75.
- 24-Fuenzalida M, Lemus S, Illanes J, Montiel E, Acuña O, Lemus D. Histochemical detection of sugar residues in lizard teeth (*Liolaemus gravenhorsti*): a lectin-binding study. *Biol Res* 2000;33(3-4):215-26.
- 25-Talaei-Khozani T, Aminizadeh N, Aliabadi E, Mesbah SF, Zolghadr J. Lectin reactivity of expanded mouse blastocysts after exposure to sera women with unexplained recurrent spontaneous abortion. *Reproductive Toxicology* 2005;20:531-7.
- 26-Alles AJ, Fazel AR, Spicer SS. Distribution of glycoconjugate in the optic vesicle and optic cup. *Anatomy & Embriology* 1990;182:611-6
- 27-Doherty CB, Lewis RM, Sharkey A, Burton GJ. Placental composition and surface area but not vascularization are altered by maternal protein restriction in the rat. *Placenta* 2003;24:34-8
- 28-Kelsall A, FitzGerald AJ, Howard CV, Evans RC, Singh R, Rhodes JM, Goodlad RA. Dietary lectins can stimulate pancreatic growth in the rat. *Int J Exp Pathol* 2002 Aug;83(4):203-8.
- 29-Yokose S, Zhungfeng C, Tajima Y, Fujieda K, Katayama I, Katayama T. The effects of estrogen deficiency on glycosylation of odontoblasts in rats. *J Endod* 1998 Oct;24(10):645-7.
- 30-Ravindranath RM, Basilrose RM. Localization of sulfated sialic acids in the dentinal tubules during tooth formation in mice. *Acta Histochem.* 2005;107(1):43-56.