

## Original Article

## The Phonemic Awareness Skills of Cochlear Implant Children and Children with Normal Hearing in Primary School

Aliakbar Dashtelei\* ; Hoda Mowzooni; Atiye Ashtari; Maryam Delfi; Gita Movallali  
Ahmad Salmani, Farzad Eshkenani  
University of Social Welfare and Rehabilitation Sciences, Tehran, Iran

**Objectives:** Phonemic awareness skills have a significant impact on children speech and language. The purpose of this study was investigating the phonemic awareness skills of children with cochlear implant and normal hearing peers in primary school.

**Methods:** Phonemic awareness subscales of phonological awareness test were administered to 30 children with cochlear implantation at the first to sixth grades of primary school and 30 children with normal hearing who were matched in age with cochlear implant group. All of children were between 6 to 11 years old. Children with cochlear implant had at least 1 to 2 years of implant experience and they were over 5 years when they receive implantation. Children with cochlear implant were selected from Special education centers in Tehran and children with normal hearing were recruited from primary schools in Tehran. The phonemic awareness skills were assessed in both groups.

**Results:** The results showed that the Mean scores of phonemic awareness skills in cochlear implant children were significantly lower than children with normal hearing ( $p < .0001$ ).

**Discussion:** children with cochlear implant, despite Cochlear implantation prosthesis, had lower performance in phonemic awareness when compared with normal hearing children. Therefore, due to importance of phonemic awareness skills in learning of literacy skills, and defects of these skills in children with cochlear implant, these skills should be assessed carefully in children with cochlear implant and rehabilitative interventions should be considered.

**Key words:** phonemic awareness, cochlear implant, children

Submitted: 19 September 2015

Accepted: 10 November 2015

### Introduction

Several factors contributing to language acquisition in children with CIs, such as auditory processing, non-verbal IQ, working memory, and phonological awareness (1). Phonological awareness refers to individual's understanding of the phonological structure of a word of his or her language (2). There are at least three forms of phonological awareness that include Syllabic awareness (including syllable Blending and segmentation), the rhyme awareness (including alliteration and rhyme) and Phonemic awareness (including phonemic blending, Recognizing words with the same initial phonemes, Recognizing Words with the same final phoneme, Phonemic segmentation, Naming and delete the final phoneme, Naming and delete the middle phoneme and naming and deletion of first phoneme) (1, 3-5). The most of the syllable awareness skills is formed as early as 3 years. With increasing age, children are

more aware of rhyme units such as the alliteration (vowel or consonant cluster before vowel in a syllable) and rhyme (vowel and consonant of the end of word). Finally, school-age children can be mastered in phonemic awareness skills. Phonemic awareness is the ability to identify phonemes as parts of syllables and words. Proficiency and skills in phonemic awareness appear at 6-7 years of age (3). Children with profound deafness are at risk for serious reading difficulties (7). Multiple factors affect their development of reading skills, including use of cochlear implants. The results from studies indicate that benefit of CIs is first observable at the syllable level (5). Children with cochlear implants have better phonemic skills than children having hearing aids (6, 8). Several factors influence the overall success of children experience with their cochlear implants. These factors include the age at which they receive an implant, duration of CIs,

\* All Correspondences to: Aliakbare dashtelei, Email: <Ali\_dashtelei@yahoo.com>

vocabulary skills, preoperative residual hearing, and socioeconomic status (9). A variety of recent studies have demonstrated benefits to hearing, language, and speech from implants, leading to assumptions that early implantation (age of CIs) and duration of implant should be associated with higher phonological and reading skills (1, 9-11). Individual differences in age at implant, duration of CIs and phonological development were all strongly associated with progress in phonemic skills and reading achievement in cochlear implants children for the deaf implanted children (1). To the extent that use of a cochlear implant is associated with greater use of phonological coding strategies that has a facilitative effect on the acquisition of literacy. Therefore reading scores are strongly correlated with measures of phonemic awareness (12, 13). So if the child has insufficient phonemic skills, his or her reading skills will face problem (1). Phonemic awareness needs a developed phonological system of phonemes representations(14). Thus, in children with cochlear implant, before the operation and in pre-lingual course, deprivation of auditory system has a negative effect on the development of phonological skills (14). Following this problem, children's literacy skills will be problematic (13, 15, 16).

Therefore understanding phonological features of this group, helps teachers and rehabilitation specialist for proper planning and intervention and because studies in investigating phonemic awareness in children with cochlear implants are rare; the aim of this study was to compare the phoneme skills in children with cochlear implant and normal children in primary school.

**Methods**

Present study employed a Cross-sectional design and was done on 30 children with cochlear implantation at the first to sixth grades of primary schools and 30 children with normal hearing who were matched in age with cochlear implant group. Children with cochlear implant were selected from Special education centers in Tehran and children with normal hearing were recruited from primary schools in Tehran. Inclusion criteria were being

monolingual, having cochlear implant, age of implantation is over 5 years, and at least 1 to 2 years benefited from cochlear implantations. Exclusion criteria were children with structural defect in the speech organs movement, neuromuscular problems, and history of epilepsy, seizures and brain injury.

In this study, phonological awareness test of Soleimani's et al. was used. This visual test includes three parts: syllabic awareness, rhyme awareness and phonemic awareness. In total, the test has 10 subscales and the subscales include 10 questions. Phonemic awareness includes seven subscales: phonemic blending, identify words with the same initial phoneme, identify words with the same final phoneme, phonemic segmentation, naming and deletion final phoneme, middle phoneme deletion, and naming and remove the initial phonemes. The demographic personal information questionnaire, inclusion and exclusion criteria were collected from the teachers. The test was performed for each subject in a relaxed and convenient environment. For each correct response, rating 1 was assigned and for each incorrect response, rating 0 was assigned, and the total numbers of test results were recorded. In some of subscales, normal distribution of data and homogeneity of variances were observed, so the parametric independent t-test was used. But in other subscales because data distribution was not normal and Homogeneity of variances was not observed, therefore nonparametric Mann-Whitney U test was used for comparing the phonemic awareness in subscales in both control and experimental groups.

**Results**

Totally 60 children, 30 children with cochlear implant and 30 normal hearing children were participated in the study. Most of them were in first grade of primary school (26.7 percent) and least of them were in second and sixth grade (13.3).the participants aged between 6 and 11 years. The mean age and standard deviation of participants were respectively 8.21 and 1.77. In table (1) mean, standard deviation and T test have been reported in two groups.

**Table 1.** Mean, standard deviation, and T-test of two independent groups

variable	Groups	numbers	mean	Standard deviation	T	P	Eta <sup>2</sup>
Phonemic awareness	Children with CI	30	22.87	18.902	8.62	0.000	0.56
	Children with NH	30	59.57	13.642			
Phonemic segmentation	Children with CI	30	2.47	2.801	9.323	0.000	0.60
	Children with NH	30	8.23	1.906			
Naming and deletion the middle phoneme	Children with CI	30	1.57	2.979	8.077	0.000	0.53
	Children with NH	30	7.70	2.902			
Naming and deletion the first phoneme	Children with CI	30	1.37	2.723	8.225	0.000	0.54
	Children with NH	30	7.57	3.104			

According to information obtained from the above table, we know that the mean of phonemic skill scores in normal hearing children were more than twice of cochlear implant children. The same situation observed between its normal subscales. Thus, the phonemic segmentation, naming and middle phoneme deletion, and naming and first phoneme deletion in normal hearing children are significantly more than cochlear implant children. T-test results show that phonemic skills score has statistically significant differences between two groups ( $T=8.623$ ,  $P<0.0001$ ). In other words,

phonemic skills were statistically significant difference between two groups. It is also a significant difference in the normal subscales of phonemic skills between two groups ( $P<0.0001$ ). According to the size effect scores (Eta square) can be concluded that the minimal difference between two groups is in the naming and deletion the middle phoneme subscales ( $0.53=Eta^2$ ) and the maximum difference is in the Phonemic segmentation subscales ( $0.60= Eta^2$ ).

The mean rating of phonemic awareness subscales and Mann-Whitney U were evaluated in the table (2):

**Table 2.** The mean rating and Mann-Whitney U test for comparison abnormal subscales phonemic awareness skills

Variable	group	number	Mean rating	Z	p
Phonemic blending	Children with CI	30	19.65	5.09	0.000
	Children with NH	30	41.35		
Recognizing words with the same initial phonemes	Children with CI	30	17.72	5.85	0.000
	Children with NH	30	43.28		
Recognizing words with the same final phoneme	Children with CI	30	18.88	5.26	0.000
	Children with NH	30	42.12		
Naming and deletion the final phoneme	Children with CI	30	20.18	4.73	0.000
	Children with NH	30	40.82		

As above table, mean rating of phonemic blending, recognizing words with the same initial phonemes, recognizing Words with the same final phoneme, and naming and deletion the final phoneme in normal hearing children are significantly more than children with cochlear implant. Therefor children with cochlear implant and normal hearing children in abnormal phonemic awareness subscales are statistically significant differences ( $p<0.0001$ ).

**Discussion**

The main aim of this study was to evaluate and compare the phonemic awareness in children with cochlear implant and normal hearing children who were in primary school. The results of this study (table 1) have been showed, generally, children with cochlear implant showed weaker performance on phonemic awareness tasks in comparison to normal hearing children. Also, results from seven subscales of phonemic awareness skills revealed that children with cochlear implantation lag those of normal hearing children. These findings are in agreement with previous studies (1, 8, 17-22). Consistent with previous studies, there is a strong relationship between phonemic awareness and literacy skills. Therefore weak performance of children with cochlear implant in phonemic awareness skills leads to the literacy problems. Also results of Table 3 showed that children with cochlear implant have

lower performance in comparison with normal hearing children in all subscales of phonemic awareness skills(1, 9, 11, 12, 14, 16, 18, 23, 24). Spencer and Oleson in their study examine phonemic awareness in cochlear implants children and normal ones and report that the performance of children with cochlear implant in a phonemic awareness tasks (phonemic blending task) is near to normal hearing children But in a phonemic segmentation tasks, children with cochlear implant did in a lower level than the normal hearing children (16). While our study showed that not only phonemic segmentations skills in cochlear implants children and normal hearing children, are significantly different (consistent with the Spencer and Oleson findings), but phonemic blending skills are also significantly different (not consistent with Spencer and Oleson findings). In fact, it can be argued that there is a significant difference in phonemic awareness skills between the children with cochlear implant and normal hearing children. Although according to previous studies children with cochlear implant had better performance compared with children with hearing loss who use hearing aids but they show lower performance in phonemic awareness when compared to normal hearing children (6, 8, 16). This may be due to factors such as age of implantation or duration of using the implant prosthesis. In fact, previous studies

have reported that if Cochlear implantation perform in early ages and duration of using the prosthesis be long, difficulties in phonemic awareness skills in cochlear implant children will be fewer and the difference in performance with normal children will be lower (10, 11, 25-27).

### Conclusion

Generally the results of this study showed that phonemic awareness skills in Children with Cochlear Implant are lower than normal hearing children. This weakness can lead to problems in the children's literacy. Therefore Assessment of phonemic awareness skills is essential to identify

defects in these skills and also to prevent literacy problems in cochlear implants children. The limitations of this study was the small number of Cochlear Implanted Children at primary school and lack of gender variable in the analysis due to limited number of each sex, and the hope is that these restrictions will be removed in future studies.

### Acknowledgment

We would like to thank the children, teachers, and support staff of all the schools and Department of Special Education in Tehran for their active and enthusiastic cooperation as well as everyone else who helped conducting this study.

### References

1. Johnson C, Goswami U. Phonological awareness, vocabulary, and reading in deaf children with cochlear implants. *Journal of Speech, Language, and Hearing Research*. 2010;53(2):237-61.
2. Gillon GT. *Phonological awareness: From research to practice*: Guilford Press; 2004.
3. Bernthal JB, N. Flipsen, P. *Articulation and Phonological Disorders*. SIXTH ed2009.
4. James D, Rajput K, Brinton J, Goswami U. Orthographic influences, vocabulary development, and phonological awareness in deaf children who use cochlear implants. *Applied Psycholinguistics*. 2009;30(04):659-84.
5. James D, Rajput K, Brown T, Sirimanna T, Brinton J, Goswami U. Phonological awareness in deaf children who use cochlear implants. *Journal of Speech, Language, and Hearing Research*. 2005;48(6):1511-28.
6. Jalilevand N. A review on the speech and language development of deaf children who receive cochlear implantation. 1392.
7. Pascoe M, Stackhouse J, Wells B. Persisting speech difficulties in children: children's speech and literacy difficulties: John Wiley & Sons; 2006.
8. Law ZW, So LK. Phonological abilities of hearing-impaired Cantonese-speaking children with cochlear implants or hearing aids. *Journal of Speech, Language, and Hearing Research*. 2006;49(6):1342-53.
9. Connor CM, Zwolan TA. Examining multiple sources of influence on the reading comprehension skills of children who use cochlear implants. *Journal of Speech, Language, and Hearing Research*. 2004;47(3):509-26.
10. DesJardin JL, Ambrose SE, Eisenberg LS. Literacy skills in children with cochlear implants: The importance of early oral language and joint storybook reading. *Journal of Deaf Studies and Deaf Education*. 2009;14(1):22-43.
11. Marschark M, Rhoten C, Fabich M. Effects of cochlear implants on children's reading and academic achievement. *Journal of Deaf Studies and Deaf Education*. 2007.
12. Dillon CM, de Jong K, Pisoni DB. Phonological awareness, reading skills, and vocabulary knowledge in children who use cochlear implants. *Journal of deaf studies and deaf education*. 2011:enr043.
13. Geers AE. Predictors of reading skill development in children with early cochlear implantation. *Ear and hearing*. 2003;24(1):59S-68S.
14. Weisi F, Shahbodaghi MR, Dadgar H, Moradi AR, Faghihzadeh S. Comparison of reading skills between cochlear implantation and normal hearing children in second and third grade elementary in Tehran. *Modern Rehabilitation*. 2012;6(3):13-9.
15. Archbold S, Harris M, O'Donoghue G, Nikolopoulos T, White A, Lloyd Richmond H. Reading abilities after cochlear implantation: The effect of age at implantation on outcomes at 5 and 7 years after implantation. *International journal of pediatric otorhinolaryngology*. 2008;72(10):1471-8.
16. Spencer LJ, Oleson JJ. Early listening and speaking skills predict later reading proficiency in pediatric cochlear implant users. *Ear and hearing*. 2008;29(2):270.
17. Spencer LJ, Tomblin JB. Evaluating phonological processing skills in children with prelingual deafness who use cochlear implants. *Journal of Deaf Studies and Deaf Education*. 2008:enn013.
18. Ambrose SE, Fey ME, Eisenberg LS. Phonological awareness and print knowledge of preschool children with cochlear implants. *Journal of Speech, Language, and Hearing Research*. 2012;55(3):811-23.
19. Harris M, Moreno C. Deaf children's use of phonological coding: Evidence from reading, spelling, and working memory. *Journal of Deaf Studies and Deaf Education*. 2004;9(3):253-68.
20. Soleymani Z, Saeedmanesh M, Dastjerdi M, Mehri A, Jahani Y. Relationship between phonological awareness, rapid automatized naming and reading in first grade students in Tehran, Iran. *Audiology*. 2010;19(1):18-25.
21. Weisi F, Rezaei M, Lotfi G, Valadbeigi A. Comparison of phonological awareness between children with cochlear implants and children with hearing aids. *Pajouhan Scientific Journal*. 2013;11(2):35-8.
22. Mamoodabadi N, Soleymani Z, Khodami M, Ajalloeian M, Jalaei S. A Comparative Study OF Performance of Normal and Cochlear Implanted Children in Two Phonological Awareness Tests. 2013.
23. Goswami UB, P. *Phonological Skills and Learning to Read*. Lawrence Erlbaum. 1990.
24. Tavakol S, Shafiei B. Identification of phonological awareness skills in down's syndrome children with Educable 7-12 Years old mental age in Isfahan. 2008.

25. Duchesne L, Sutton A, Bergeron F. Language achievement in children who received cochlear implants between 1 and 2 years of age: Group trends and individual patterns. *Journal of Deaf Studies and Deaf Education*. 2009:enp010.
26. Schorr EA, Roth FP, Fox NA. A comparison of the speech and language skills of children with cochlear implants and children with normal hearing. *Communication Disorders Quarterly*. 2008;29(4):195-210.
27. James D, Rajput K, Brinton J, Goswami U. Phonological awareness, vocabulary, and word reading in children who use cochlear implants: Does age of implantation explain individual variability in performance outcomes and growth? *Journal of Deaf Studies and Deaf Education*. 2008;13(1):117-37.