

Journal of Biodiversity and Ecological Sciences $(JBES^{\mathbb{C}})$

No.1, Issue3 ISSN: 2008-9287 Summer 2011 JBES

Orginal Article

MJ.Haar¹

J.Dekker^{2*}

Seed production in weedy Setaria spp.-gp.

Received Date:July/05/2011 Accepted Date:Sep/20/2011

Abstract

Seeds from Setaria faberi, S. viridis and S. pumila panicles in three Iowa crop fields were collected for the entire reproductive period. Seed number, panicle length, and seed number per panicle length varied among species, panicle types and sites. Greater numbers of seed per plant and per panicle were observed than previously reported. Setaria seed rain exhibited some stable, and many more plastic, responses. S. faberi panicles were consistently longer than those of S. viridis. S. viridis parameters were greater than S. pumila. Earlier panicles were greater than, or similar to, later ones for all parameters. More typically, tillers and panicles responded to local conditions in a plastic way, confounding the formulation of seed production generalizations. In S. faberi and S. viridis no consistent relationship between seed number and panicle length was observed among different tiller types. A more consistent relationship between parameters was observed for S. pumila compared to the others, making prediction possible for this species. The stability and plasticity of these relationships is partially due to the differences in S. faberi and S. viridis panicle, fascicle and spikelet morphology compared to S. pumila. These stable and plastic responses provide fine-scale adjustment to a locality, maximizing exploitation of local opportunity. Keywords: seed rain, tillering, panicle, plasticity, prediction, fecundity

INTRODUCTION

The success of the *Setaria* species group [*S. faberi* Herrm., giant foxtail; *S. viridis* [L.] Beauv., green foxtail; *S. pumila* [L.] Beauv., yellow foxtail] as weeds is due in part to seed production [the seed rain] under a wide range of environments and to high reproductive output under favorable conditions. The production of many dormant seed disperses germination over time and soil seed pools allow *Setaria* to persist at a locality. These adaptive traits provide *Setaria* the ability to seize and exploit opportunity in local agroecosystems.

Seed production is an emergent property of *Setaria* reproductive morphology. Seed numbers produced by a *Setaria* are a function of differences in plant architecture: shoot tilleringpanicle formation, panicle-fascicle branching, and fascicle spikelet-floret development. Inherent plastic differences in these reproductive structures among *Setaria* species determine the reproductive responses of species and populations to available opportunity in its immediate environment.

Setaria panicle inflorescences develop at the terminal ends of shoot tillers. Panicles that develop at the end of the main shoot are referred to as primary panicles [1°; Figure 1]. Secondary panicles [2°] arise at the nodes of the primary tiller, and tertiary panicles [3°] are those that laterally from secondary branch tillers. Developmentally, primary panicles flower first on a plant, followed by secondary, then tertiary. Setaria spp., panicles are composed of fascicles, which consist of spikelet [with florets] and bristle [seta only] shoots [Narayanaswami, 1956; figure 1]. The fascicle-spikelet structure differs among Setaria species. In S. faberi or S. viridis the number of fascicles within each panicle varies, a plastic response to a plant's immediate environment [Clark and Pohl, 1996]. Longer, earlier developing S. faberi or S. viridis panicles often have the most extensive fascicle branching, as well as more spikelets per fascicle [typically 4-6 or more per fascicle]. The plastic response of S. faberi or S. viridis to its immediate environment is also revealed in the number of spikelets-florets

1-2*Weed Biology Laboratory, Iowa State University, Ames, Iowa 50011 USA <u>jdekker@iastate.edu</u> that fully mature, which determines the seed number per panicle length [seed density]. Under favorable conditions more spikelets are able to develop into seeds, while under unfavorable conditions spikelets may abort. *S. pumila* panicle morphology is different from that of *S. faberi* or *S. viridis*. Only a single, terminal, spikelet-floret is found in each *S. pumila* fascicle [Clark and Pohl, 1996]. This stable, fixed morphology limits the ability of *S. pumila* to respond in a plastic way to its environment in terms of seed number relative to that of *S. faberi* or *S. viridis*.

In addition to morphological traits, genotype and environmental interact to influence the productivity of individual Setaria plants. Weedy Setaria species contain a relatively low degree of genetic variation compared to other species, but there exists significant genotypic and biotypic variation within and among S. viridis and S. pumila which may be associated with differences in seed production [Hubbard, 1915; Norris and Schoner, 1980; Santelmann and Mead, 1961; Schoner, 1978; Schreiber and Oliver, 1971; Wang et al., 1995a; Wang et al., 1995b]. The local environment also affects Setaria growth and seed production [Lee and Cavers, 1981; Nadeau and Morrison, 1986; Santelmann and Meade, 1961; Santelmann et al., 1963; Schreiber, 1965; Van den Born, 1971].

Assessment and prediction of Setaria seed production, additions to the local agricultural soil seed pool, is of considerable importance to weed management systems [Freckleton and Stephens, 2009; Holst et al., 2007; Dekker, 2011]. Rapid estimation of seed production of weedy infestations has stimulated considerable research to discover the relationship of panicle size and The relationship between seed seed numbers. number and panicle length has been described for three Setaria species as stable across environments [Barbour and Forcella, 1996; Forcella et al., 2000] and has been used to estimate Setaria seed production [Defelice et al., 1989; Fausey et al., 1997]. This relationship has also been shown to be stable in E. crus-galli [Norris, 1992]. The values reported for Setaria seed production may not be reliable due to both the methodologies used to determine seed production and environmental variability. Although Setaria panicles and seeds develop over a period of time [e.g., ca. 60 d; Haar and Dekker, 1995], it is common to find reports of seed production based on a single or periodic harvests [Biniak and Aldrich, 1986; Defelice et al., 1989; Fausey et al., 1997; Kawano and Miyake, 1983; Knake, 1972; Nadeau and Morrison, 1986;

Santelmann and Meade, 1961; Schreiber, 1965; Wall, 1993]. Values determined by such methods represent only a fraction of total seed production. There is little consistency in the degree of maturity of plants from which seed is harvested or the units of measure for Setaria seed production. Seed production has been reported on per panicle [Biniak and Aldrich, 1986; Santelmann et al., 1963], per panicle length [Fausev et al., 1997], per plant [Kawano and Miyake, 1983; Nadeau and Morrison, 1986; Schreiber, 1965; Wall, 1993] or per area basis [Defelice et al., 1989]. The relationship between panicle length and seed number may also be compromised by the lack of information about whether this relationship pertains under a variety of environmental conditions.

We have made the relationship between seed number and panicle length the subject of further investigation, in particular its validity and consistency among *Setaria* species, panicle types and environments. An accurate measure of seed production could improve the understanding of Setaria population dynamics and assist in the development of more efficient weed management systems. Conversely, highly variable relationships between Setaria panicle length and seed production may provide us some understanding of stability and plasticity in Setaria reproduction.

The first objective of this study was to provide an accurate measure of *Setaria* seed production by collecting the entire seed output of individual panicles throughout the seed rain period. Secondly, seed production and panicle length were compared among the three *Setaria* species, developmental panicle types and sites. We hypothesized that although differences in seed production and panicle length may exist among *Setaria* species, panicle types and sites, the relationship between seed number and panicle length would be stable across environments. Evaluating this hypothesis was the third objective.

MATERIAL & METHODS Species and sites

The three weedy *Setaria* species in this study frequently occur together in Iowa agricultural fields; their presence and relative proportion varies widely by locality [Wang et al., 1995a; Wang et al., 1995b]. *S. faberi, S. viridis* and *S. pumila* were selected for experimental study at three sites near Ames, Iowa, in 1995. Site selection criteria included a diversity representative of central Iowa agricultural production, and a representative sample of individual plant sizes occurring at those sites [table 1].

Seed Collection

Panicle type was determined at anthesis and bagged. Bagging occurred between July 25 and September 8, 1995 when the panicles were covered with 7.6 by 25.4 cm mesh pollination bags [Delnet non-woven fabrics, Applied Extrusion Technologies Inc. Middletown, DE 19899, USA] held in place with wire. No more than two bags were attached to a plant. Panicles were harvested on October 10 and 12, 1995. A killing frost on September 21 prevented a few late tertiary S. glauca panicles from completing maturation.

Data Collection

Panicles and seed were removed from bags after harvest, and length was determined by measuring from the panicle tip to the point of attachment for the most basal fascicle [Norris, 1992]. Seed that remained attached to the panicle was removed. Seed was cleaned with an air flow cleaner to remove aborted and sterile spikelets and debris, weighed, and electronically counted.

Statistical Analysis

Means were calculated for seed number per panicle, panicle length and seed density [seed number per unit panicle length] for *Setaria* species, panicle types and sites. Paired t-tests $[\Box=0.05]$ were used to separate means and slopes for all parameters among panicle types within a species and site, among sites within a species and panicle type, among species within a site and among sites within a species. Linear models best described [highest R2 value] the relationships between panicle length and seed number, and of panicle length and seed number per panicle length. The linear regression procedure of SAS [1989] was used for analysis.

RESULTS

Setaria Species

Setaria faberi. S. faberi seed number [SN], panicle length [PL] and seed density [SPD] was greater for primary [1°] than for tertiary [3°] panicles, with a single exception in SPD at a site A [Tables 2, 3, 4; comparison 1]. For all parameters, secondary [2°] panicles were either similar to or greater than 3°. When differences occurred among sites within a panicle type, SN, PL and SPD were greater at site B than at the other sites, with the exception of 3° panicles, which usually did not differ among sites [Tables 2, 3, 4; comparison 2]. Averaged over all panicle types, SN, PL and SPD were similar at all sites, with a single exception in which seed density was greater at site B than C [Tables 2, 3, 4; comparison 4].

The ability of the linear model to describe the relationship between PL and SN or SPD [R2] varied widely among sites and panicle types. The change in SN with PL was greater in 2° panicles than that observed for 3° panicles at two of the three sites [Table 5, comparison 1]. The secondary and 3° panicles at site C did not show a change in SN with panicle length. No difference in the relationship between SN and PL was observed between 1° and 2° panicle types at any of the three sites. Differences in the SPD per panicle length relationship were not found among panicle types at any site.

Comparisons within individual *S. faberi* panicle types among sites revealed that differences were largely due to changes in secondary and 3° but not 1° panicles [Table 5, comparison 2]. Changes in SN and SPD with panicle length were similar in 1° tillers at all three locations. Changes in 2° *S. faberi* SN and SPD with panicle length were greater at site B compared to both other sites. Changes in 3° *S. faberi* SN and SPD with panicle length were greater at site B compared to site C.

When averaged over all three panicle types [Table 4, total] *S. faberi* SN per panicle was correlated with panicle length. Greater change in SN with changes in PL were observed at site B compared to the other two sites [Table 5, comparison 3]. Site B also revealed a greater change in SD with PL. Unlike at sites A and C, *S. faberi* SD at site B increased as PL increased.

Setaria viridis. Primary panicles were greater in SN, PL and SPD than 3° tillers at both sites and secondary panicles were either greater or similar to tertiary panicles [Tables 2, 3, 4; comparison 1]. Whether averaged over all panicle types or compared by individual panicle type, SN, PL and SPD in green Setaria were similar at both sites [Tables 2, 3, 4; comparisons 2 and 4].

Changes in SPD with panicle length were similar for all panicle types [Table 4 comparison 1] at site C, but lower for 3° panicles at site A. Comparisons within individual S. viridis panicle types between sites revealed differences for 3° panicles, but not 1° or 2° panicles [Table 5, comparison 2]. The change in SN or SPD with 3° panicle length was greater for site A than site C. S. viridis SN and SPD changed with panicle length at both sites when averaged over all three panicle types [Table 5, comparison 3]. The rates of increase were similar at both sites for either the SD or SN to panicle length relationship. Setaria pumila. Seed number, PL and SPD in 1° S. pumila tillers were either greater than or similar to those in 2° tillers, depending on the site in which the comparison was made [Tables 2, 3, 4; comparison 1]. Differences between 1° and 2° panicles were not observed in S. pumila at site C. For all three parameters, 1° and 2° panicles at site B were either greater than or similar to comparable types at the other sites. Tertiary S. pumila panicles, however, were similar for all parameters at all sites in which they were sampled [Tables 2, 3, 4; comparison 2]. Inferences about S. pumila were compromised because no 3° panicles occurred at site A. High plant density is thought to be responsible for the absence of S. pumila 3° panicles. When averaged over panicle types, the SN, PL and SPD for S. pumila at site B were either similar to or greater than those at the other sites [Tables 2, 3, 4; comparison 4].

Comparisons among sites for individual S. pumila panicle types revealed no differences in terms of changes in SN or SPD with changes in panicle length [Table 5, comparisons 1 and 2]. Seed number did not change with PL in 1° S. pumila panicles at sites B and C, in 2° at site C, or in 3° at either site [Table 5]. The only change in SD with changes in PL occurred in 2° panicles at site A. When averaged over all three panicle types, the number of S. pumila seed per panicle increased with PL at all three sites, while SD increased at two sites, [Table 5 comparison 3]. The degree of change was similar at all three sites for both parameters.

Setaria Species-Group

Relative differences in mean PL, SN per panicle and SPD between species often changed among sites. When averaged over panicle types, variation was large; however, some consistent.

DISCUSSION

Setaria Seed Production

Previous reports of Setaria seed production may have underestimated the seed We found a higher number of seed per rain panicle at each site [Table 2] compared to that previously reported for S. faberi, [Biniak and Aldrich, 1986: 207 seeds seeds per panicle] or [Defelice et al., 1989: 110 to 280 seeds per panicle]; for S. viridis, [Wall, 1993: 437 to 577 seeds per panicle] or [Van den Born, 1971: 350 to 500 seed per panicle]. The mean number of S. faberi seed per panicle in this study is greater than that reported for the entire S. faberi [220, 730 and 2,423 seeds per plant], S. viridis [234] or S. pumila [199] plant in two studies by Kawano and

Miyake [1983]. These differences in seed number could be due to differences in collection technique, duration of time over which seed was gathered, genotypic responses or differences in resource availability and conditions at individual sites. Conversely, our data may still be an underestimate because the pollenation bags used to capture seed may have decreased the final seed yield by shading the panicle.

Plasticity and Stability in Setaria Seed Production

In this study all observed seed production parameters were plastic. This variability was revealed at several levels of plant organization: *Setaria* species, tiller branches and individual panicles. In some instances seed production was stable, and independent of the interaction between the biological factors and the site they grew on. More typically, individual *Setaria* species, tillers and panicles responded to the site they grew on in a plastic way, confounded the formulation of generalizations about panicle length and seed number.

Few consistent observations were made about seed productivity between the three Setaria Some species and parameters were species. consistent [stable] in different environments, while others were plastic in response to site S. faberi and S. pumila seed conditions. production usually was greater at site B within each of the individual panicle types. But, S. pumila seed and panicle parameters were usually greater at site B when averaged over types, unlike the other two species. The productivity of tiller panicle types in S. viridis was similar at both sites in which it was evaluated. Earlier-maturing panicle types [e.g., primary] of all species usually were more productive than later maturing panicles [e.g., tertiary].

As conditions changed, each *Setaria* species adjusted its seed productivity in a plastic manner, but the way this plasticity was expressed differed in each of the individual species. Within individual panicles of any type, seed production plasticity was expressed in two ways. First, the seed number per panicle could change with changing panicle length [Table 5, seed number per panicle length]. Second, seed density could change with changing panicle length].

For *S. faberi*, intra-panicle plasticity among sites was indicated by differences in the relationship [slope] between secondary and tertiary panicle seed number and density with changes in length. For *S. viridis*, intra-panicle plasticity was only observed in tertiary panicles, in which the relationship between seed number and density changed with changing panicle length. A stable relationship [slope] between seed number and density with changing panicle length was observed more frequently among primary *S. faberi* and *S. viridis* panicles.

S. pumila was more stable than S. faberi and S. viridis in response to changing site conditions, and generalizations about panicle length and seed number may be possible. Changes in S. pumila seed number with panicle length was more stable than in the others, and even less plastic than S. viridis or S. faberi in response to changes in seed density. These species differences may be a function of differences in panicle branching and fascicle organization [Clark and Pohl, 1996]. S. pumila panicle fascicles contain a single fertile spikelet, while in S. viridis and S. faberi fascicles can support one or more fertile spikelets depending on resources and conditions. S. pumila stability also occurs at the level of panicle type. No differences between primary, secondary or tertiary panicles were observed within a site in changes in seed number and density with panicle length.

Plasticity and Prediction in *Setaria* Seed Production

This study indicates that *Setaria* traits such as seed number per panicle, panicle length, and seed density possess both stable and variable attributes. These traits were often plastic, but there appeared to be a limit to the range within which these plastic responses occur. The relative differences in panicle length among species, sites and panicle types revealed these limits to plasticity.

Although many of these characteristics were variable, some consistent observations occurred. *S. faberi* had longer panicles than *S. viridis. S. viridis* panicles had a greater number of seed and higher seed density than yellow. Earlier-developing panicle types were always greater than or similar to the later developing panicle type for each of the parameters measured. Where a difference among sites was found, the values from site B were always the greatest. All other comparisons varied.

The correlation between seed number and panicle length was not constant across environments or panicle types. It is evident that estimates of seed production based on the panicle length must be population and panicle-type specific for *S. faberi* and perhaps for *S. viridis*. Comparatively little change in seed density with length for *S. pumila* was observed, indicating generalizations and predictions of seed rain may be possible in this species as it is in others [Norris, 1992]. The observed degree of variation among these characteristics calls into question the accuracy of estimating seed production based on panicle length for *S. faberi* or *S. viridis* as proposed by some [Barbour and Forcella, 1993; Fausey et al., 1997; Forcella et al., 2000].

Plastic responses to changing conditions allows an individual *Setaria* species, tiller and plant to finely adjust its seed production to highly localized conditions. Although this plasticity confounds our ability to develop quick, accurate predictive tools about the seed rain, it allows the individual weed to exploit the conditions and resources available to maximum advantage. While selection over many generations has resulted in a degree of phenotype stability, plasticity can be highly advantageous to these plants and is preserved [Sultan, 1987].

REFERENCE

- BARBOUR, J. C.; FORCELLA, F., (1993). Predicting seed production by foxtails (*Setaria spp*). Proceeding North Central Weed Science Society., 48, 100 pp.
- BINIAK, B. M.; ALDRICH, R. J., (1986). Reducing velvetleaf (Abutilon theophrasti) and giant foxtail (*Setaria faberi*) seed production with simulated-roller herbicide applications. Weed Science., 34, 256-259.
- CLARK, L. G.; POHL, R. W., (1996). Agnes Chase's first book of grasses, 4th ed. Smithsonian Institution Press., Washington, London.
- DEFELICE, M.S.; et al., (1989) Weed control in soybeans (Glycine max) with reduced rates of postemergence herbicides. Weed Science., 37, 365-374.
- DEKKER, J., (2011). Evolutionary models of weed life history population dynamics, In: Daniels, J. A. (Ed.), Advances in environmental research (Vol. 10). Hauppauge, NY: Nova
- Science Publishers.
- FAUSEY, J.C.; et al., (1997). Giant foxtail (*Setaria faberi*) interference in nonirrigated corn (Zea mays). Weed Science., 45, 256-260.
- FORCELLA, F.; et al., (2000). Estimating seed production of three *Setaria* species in row crops. Weed Science., 48. 436-444.
- FRECKLETON, R. P.; STEPHENS, P. A., (2009). Predictive models of weed population
- dynamics. Weed Research., 49, 225-232.
- HAAR, M.; DEKKER, J., (1995). Correlation of giant (*Setaria faberi*), green (*S. viridis*) and yellow (S. glauca) foxtail seed production with panicle length, branching and tillering. Proceedings North Central Weed Science Society., 50, 102.
- HOLST, N.; et al., (2007). Field weed population dynamics: a review of model approaches and applications. Weed Research., 47, 1-14.
- HUBBARD, F. T., (1915) A taxonomic study of *Setaria* italica and its immediate allies. American Journal Botany., 2, 169-198.
- KAWANO, S.; MIYAKE, S., (1983). The productive and reproductive biology of flowering plants X. Reproductive energy allocation and propagule output of

five congeners of the genus *Setaria* (Gramineae). Oecologia., 57, 6-13.

- KNAKE, EL., (1972). Effect of shade on giant foxtail. Weed Science., 20, 588-592.
- LEE, S. M.; CAVERS, P. B., (1981). The effects of shade on growth, development and resource allocation patterns of three species of foxtail (*Setaria*). Canadian Journal Botany., 59, 1776-1785.
- NADEAU, L. B.; MORRISON, I. N., (1986). Influence of soil moisture on shoot and root growth of green and yellow foxtail (*Setaria* viridis and S. lutescens). Weed Science., 34, 225-232.
- NARAYANASWAMI, S., (1956). Structure and development of the caryopsis in some Indian millets(VI. *Setaria* italic). Botanical Gazette., 118, 112-122.
- NORRIS, R. F.; SCHONER, C. J., (1980). Yellow foxtail (*Setaria* lutescens) biotype studies: Dormancy and germination. Weed Science, 28, 159-163.
- NORRIS, R. F. (1992). Relationship between inflorescence size and seed production in barnyardgrass (Echinochloa crus-galli). Weed Science., 40, 74-78.
- SANTELMANN, P.W.; MEADE, J.A., (1961). Variation in morphological characters and Dalapon susceptibility within the species (*Setaria* lutescens and *S. faberi*). Weeds., 9, 407-410.
- SANTELMANN, P.W.; et al., (1963). Growth and development of yellow foxtail and giant foxtail. Weeds., 11, 139-142.
- Statistical Analysis Systems., (1989). SAS/STATS User's Guide, version 6, 4th ed. SAS Institute, Inc., Cary, NC.
- SCHONER, C. A., (1978). Yellow foxtail (*Setaria* lutescens) biotype studies: Growth and morphological characteristics. Weed Science., 26, 632-636.
- SCHREIBER, M. M., (1965). Effect of date of planting and stage of cutting on seed production of giant foxtail. Weeds.,13, 60-62.
- SCHREIBER, M. M.; OLIVER, L. R., (1971). Two new varieties of *Setaria* viridis. Weed Science., 19, 424-427.
- SULTAN, S. E., (1987). Evolutionary implications of phenotypic plasticity in plants. Evolutionary Biology., 21, 127-178.
- VAN DENBORN, W. H., (1971). Green foxtail: Seed dormancy, germination and growth. Canadian Journal Plant Science., 51, 53-59.
- WALL, D. A., (1993). Comparison of green foxtail (*Setaria* viridis) and wild oat(Avena fatua) growth, development and competitiveness under three temperature regimes. Weed Science., 41, 369-378.
- WANG, R. L., (1995). Weedy adaptation in *Setaria spp.* I. Isozyme analysis of genetic diversity and population genetic structure in(*Setaria* viridis). American Journal Botany., 82, 308-317.
- WANG, R. L., (1995). Weedy adaptation in *Setaria spp.* II. Genetic diversity and population genetic structure in S. glauca, S. geniculata and *S. faberi* (Poaceae). American Journal Botany., 82, 1031-1039.

5

Site	Crop	Weed control	Setaria spp.	Weed population size
	er op		S. faberi	······································
А	soyabean	none	S. viridis	large (277-2903 plants per m ²
	-		S. pumila	
D	maize	inter-row cultivation	S. pumila	low
D	soyabean	inter-row cultivation	S. faberi	low
			S. faberi	
С	maize	inter-row cultivation	S. viridis	low
_			S. pumila	

Table 1 . Site clopping (clop, weed control) and weed (species, population size) experimental mornation.

Table 2. Mean (m) seed number and standard error (SE) for *Setaria* species (*S. faberi*, *S. viridis*, *S. pumila*) panicle types (P; 1°, primary; 2°, secondary; 3°, tertiary) and panicle number (n) at three Ames, Iowa sites (A, B, C).

^a statistical comparisons: 1 = within a species and site, among panicle types (within a column); 2 = within a species and panicle type, among sites (within a row); 3 = within a site, among species (within a column); 4 = among sites, within a species (within a row). Means within a comparison followed by the same letter are not significantly different (P=0.05) as determined by t-tests.

									See	d numb	er pe	er pai	nicle										
			Si	ite A							Sit	e B							Sit	te C			
			(Compar	ison	a					C	omp	ariso	n ^a					(Com	bariso	n ^a	n
Р	No.	SE	1	2	3	4	n		No.	SE	1	2	3	4	n		No.	SE	1	2	3	4	
Seta	iria fab	eri																					
1°	725	133	а	В			14		2127	110	а	Α			22		427	54	а	В			7
2°	578	58	ab	В			22		1063	123	b	Α			29		330	61	а	В			9
3°	317	57	b	AB	Ì		4	İİ	365	37	с	Α			31		165	20	b	В			8
m	540	119	ĺ		а	А	3		1185	512	i .		a	Α	3	$\mathbf{\nabla}$	355	37		Ī	ab	А	3
Seta	iria vir	idis																					
1°	725	105	а	А			16										685	69	a	А			1 3
2°	592	71	а	А			9	İİ					V		7		413	58	b	А			1 1
3°	172	56	b	Α	İ		7				7 ,						144	24	c	А			1 9
m	496	167			а	А	3									İ	414	156		İ	а	А	3
Seta	iria pu	nila																					
1°	105	16	а	В			9		213	17	а	Α			11		139	35	а	В			5
2°	63	11	b	В			15		167	12	b	Α			26		147	13	а	Α			3
3°	-	-	Ì		ĺ		0		54	9	с	Α	ĺ		20	ÍÍ	64	12	а	Α			3
m	84	17			b	В	2		145	47			а	Α	3		117	26		Ī	b	AB	3

Sig	mincantly o	IIIIereni	(P=0	(US) a	s dete	rmine	d by t-te	ests.															
										M	ean par	nicle le	ngth (c	<u>m</u>									
			Site	Þ				_			Site	B				_				Site	εC		
				Comp	arison ^a	Ĵ						Compa	rison ^a							2	omparison ^a		,
P	cm	SE	1	2	з	4	,	┝	cm	SE	-	2	ω	4	3	⊢	cm	SE	N		3 4	\vdash	
Setaria	faberi																						
1°	13.3	3 0.5	а	в			14		16.3	0.2	a	⊳			22		12.6	0.7	2	в			7
2*	12.0	0.6	в	>			22		12.0	0.7	ъ	⊳			29		11.6	1.1	ച	⊳			و
ů	7.5	0.7	в	>			4		7.3	0.4	0	⊳			31		7.6	0.9	ъ	⊳			8
в	11.1	1 1.6			a	A	3		11.9	2.6			a	⊳	з		10.6	1.5			۵	>	з
Setaria	viridis																						
1°	5.6	0.5	a	A			16										10.0	0.4	a		A		12
2°	9.1	0.5	<u>،</u>	>			9										7.5	0.6	ь		<u>></u>		11
3,	4.0	5 0.7	в	A			7										5.3	0.4	c		A		15
з	7.5	9 1.7				⊳	ω	┝								┝	7.6	1.4	Ļ.	L	р- в	>	ω
Setaria	pumila																						
1°	6./	1 0.5	a	в			9		10.9	0.8	ല	Þ			11		7.8	1.0	<u>a</u>	AB			5
2*	5.6	0.3	a				15		8.9	0.4	ь	>			26		7.3	0.6	ab	АВ			ω
ů							0		5.5	0.3	0	⊳			20		4.5	0.5	ь	Þ			ω
в	6.(0.4			c	в	2		8.4	1.6			ച	Þ	ω		6.5	1.0			в	в	з

Journal of Biodiversity and Ecological Sciences SID. ir

з	ů	2°	1°	Setan	з	ů	2°	1°	Setan	з	ů	2°	1°	Setan	P				
14		11	16	ia pumila	55	31	64	69	ia viridis	48	40	48	55	ia faberi	cm	No./			^a statis within signific
ω		1	2		12	7	6			4	4	4	9	1	SE				a site, cantly c
		σ	a			ъ	۵	۵]		ъ	a	a		1				mparis among liffere
		в	Þ			Þ	Þ	Þ			Þ	в	в		2	Com	Site		sons: 1 3 speci nt (P=(
ъ					a					۵					ω	parisonª	A		= with es (wit).05) as
Þ					Þ					AB					4				in a sp hin a c detern
-														1	5				ecies ai olumn) nined b
2	0	15	9		ω	7	9	16		ω	4	22	14						nd site ; 4 = y t-tes
																			e, amo amoi its.
16	10	19	20							28	46	80	130		cm	No./		See	ong pai ng site:
4	1	1	1							24	ω	<u>б</u>	б		SE			d numbe	nicle ty s, with
	σ	۵	۵								0	ъ	۵]	1			er per pa	/pes (w in a sp
	Þ	A	Þ	1							<u>ہ</u>	Þ	Þ	1	2	Comp	Site B	anicle ler	/ithin a ecies (
ъ.				1						۵				1	ω	arisonª		ngth (no	u colum (within
A										A					4			./cm)	1n); 2 = a row
з	20	26	11							ω	31	29	22	1	5). Me
				1					1					1					n a spe ans wit
17	14	20	17	1	50	26	54	69	1	28	23	28	33	1	cm	No./			hin a c
2	2	2	ω	1	13	ω	6	6	1	ω	ω	4	ω	1	SE				ompar
	0.			1					1			ab		1	1				cle typ ison fo
		 >			\vdash	 >									2	Com	Sit		e, amc llowe
										۵					ω	parison	е С		ong site d by th
в					a					6					4	•			es (wit 1e sam
A					A									{	\vdash				hin a le lette
ω	ω	ω	s		ω	19	11	13		ω	00	9	7		∍				row); 3 er are n

Table 5	. Line	ar regres	ssion and co	omparis	ons of	f slopes (±	s.e.) and	l coeffic	cients of	determina	tion (R	²) describin	ng the rela	tionshi	ps betwee	n panicle
length	and	seed	number,	and	of	panicle	length	and	seed	number	per	panicle	length,	for	Setaria	species
(S. faber	i, S. vi	ridis, S. p	<i>pumila</i>) and	panicle	types	(P; 1°, pri	mary; 2°,	second	ary; 3°, 1	tertiary) at	three A	mes, Iowa	sites (A, B	, C).		
^a statisti	cal con	parisons	s (within a	column): 1 =	slopes wit	hin a spe	cies and	d site an	nong panic	le type	s; $2 = slop$	es within a	a speci	es and pan	icle type
among s	ites; ar	d 3 = slc	opes for tota	l panicl	e type	s within a	species an	mong si	tes.							

			D 11		<u> </u>	Lin	ear regres	sion and	statist	ical compariso	ons	4.0			1 4	
0.1	D	C1	Panicle le	ength-	Seed 1	num	ber \mathbf{p}^2	1.0		1	Panicle lei	igth-S	eed num	ber p	er length	1.0
Site	P	Slope	SE	I"	2	3	R-	b=0		Slope	SE	1	2	3	R ²	b=0
Setaria j	faberi								-							
А	1°	38.7	95.0	a	а		0.01	Y		-1.5	6.7	а	a		0.00	Y
	2°	48.2	19.7	a	b		0.23	Ν	ļ	0.4	1.6	a	b		0.00	Y
	3°	75.9	23.9	a	ab		0.83	Y		4.7	3.1	a	ab		0.54	Y
	Tot	56.8	20.4			b	0.17	Ν	ļ	1.1	1.5			b	0.01	Y
В	1°	149.7	113.8	ab	a		0.08	Y		0.8	6.9	a	а		0.00	Y
	2°	153.4	12.8	a	a		0.84	Ν		6.8	0.8	a	а		0.70	Ν
	3°	86.4	6.0	b	a		0.88	Ν		5.5	0.8	a	a		0.60	Ν
	Tot	172.3	8.6			а	0.83	Ν		8.1	0.5			a	0.75	Ν
С	1°	64.6	15.0	a	a		0.79	Ν		2.6	1.3	a	a		0.45	Y
	2°	33.8	17.8	a	b		0.34	Y		0.6	1.3	a	b		0.03	Y
	3°	12.5	8.3	b	b		0.27	Y		-1.4	1.1	a	b		0.23	Y
	Tot	39.5	7.2			b	0.58	Ν		1.0	0.6			b	0.11	Y
Setaria	viridis										-					
А	1°	163.3	33.2	a	a		0.63	Ν		9.4	3.0	a	a		0.41	Ν
	2°	121.9	33.7	a	a		0.65	Ν		7.2	4.3	a	a		0.28	Y
	3°	82.2	8.2	b	a		0.95	Ν	Ì	9.5	1.7	a	a		0.85	Ν
	Tot	118.5	13.5	-		а	0.72	Ν	ĺ	7.9	1.2			a	0.58	Ν
С	1°	49.8	53.5	a	a		0.07	Y	Ì	-1.8	4.9	a	a		0.01	Y
	2°	68.1	24.0	a	a		0.47	Ν	1	2.8	3.2	a	a		0.08	Y
	3°	50.1	8.4	a	b		0.68	Ν	Ì	3.1	1.4	а	b		0.21	Y
	Tot	92.0	10.1			а	0.67	Ν	Ì	6.4	1.2		ĺ	a	0.43	Ν
Setaria	pumila															
A	1°	21.4	8.6	a	a		0.47	Ν		1.0	1.2	a	a		0.09	Y
	2°	24.0	5.8	a	a		0.56	Ν		2.5	0.9	a	a		0.37	Ν
	Tot	25.3	4.7		ĺ	а	0.57	Ν	i i	2.3	0.7	İ	İ	a	0.31	Ν
В	1°	12.4	5.7	a	a		0.35	Y	ΪJ.	-0.5	0.5	a	a	İ	0.12	Y
	İ	ii						Ν	1	0.5	0.5	a	a		0.04	Y
	2°	22.9	3.9	a	а		0.58		í.	ĺ			ĺ	İ	ĺ	
	30	13.6	72	а	a		0 17	Y	Ì	0.4	12	а	а		0.01	Y
	Tot	24.8	2.1			а	0.71	N	i	1.2	0.3			а	0.24	N
С	1°	24.0	14.4	а	a		0.48	Y	i	1.0	14	а	а		0.13	Y
e	2°	10.1	18.8	а 2	a		0.22	Ŷ	ł	-1 3	2.8	а а	a		0.18	Ŷ
	3°	63	22.5	a	a		0.07	Ŷ	Ì	-2.3	4.0	a a	a	ļ	0.10	Ŷ
	Tot	22.8	6-2	"	a		0.60	N	1	0.8	4.0 0.7	u	a	а	0.12	Y
	100	22.0	0.2			a	0.00	19	1	0.8	0.7			a	0.12	1

Linear regression and statistical compariso

Figure 1. Schematic diagram of weedy *Setaria* species-group reproductive shoot architecture and panicle structure; tiller and panicle types (1°, primary; 2°, secondary; 3°, tertiary): left; fascicle branching ($F_{1.5}$) on panicle axis: top, right; fascicle structure and arrangement of bristle (seta) shoots (BS) and spikelet shoots (SS) along rachilla axis: *S. pumila* (bottom, middle); *S. viridis* and *S. faberi* (bottom, right).



Journal of Biodiversity and Ecological Sciences (JBES[©])

Publish Your Work in This Journal



Submit your manuscript here: http://www.jbes.ir