

## Phytochemical response of hyssop (*Hyssopus officinalis* L.) to foliar application of jasmonic acid

Abdollah Ghasemi Pirbalouti<sup>1,2\*</sup>, Abolghasem Gorgij<sup>2</sup>, Mehdi Rahimmalek<sup>3</sup>, Behzad Hamedi<sup>2</sup>

<sup>1</sup>Medicinal Plants Program, Plant and Soil Science Department, College of Natural, Massachusetts University, Amherst, 01003, MA, USA;

\* E-mail: [aghasemipir@psis.umass.edu](mailto:aghasemipir@psis.umass.edu)

<sup>2</sup>Department of Medicinal Plants, Researches Centre of Medicinal Plants & Ethno-veterinary, Shahrekord Branch, Islamic Azad University, PO Box: 166, Shahrekord, Iran;

<sup>3</sup>Department of Agronomy and Plant Breeding, College of Agriculture, Isfahan University of Technology, Isfahan 84156 83111, Iran;

### ARTICLE INFO

**Type:** Original Research

**Topic:** Agronomy; Plant Physiology

**Received:** October 11<sup>th</sup> 2012

**Accepted:** December 27<sup>th</sup> 2012

### Key words:

- ✓ *Hyssopus officinalis* L.
- ✓ Jasmonic Acid
- ✓ Essential oil
- ✓ GC-MS

### ABSTRACT

**Background & Aim:** Jasmonic acid (JA) is regarded as endogenous regulators that play important roles in regulating stress responses, plant growth and development. Effect of foliar application of JA on the essential oil and oil chemical components of hyssop (*Hyssopus officinalis* L.) were investigated.

**Experimental:** This study conducted in a CRD with three replications and in experimental greenhouse, I.A.U., Shahrekord Branch, Iran at 2012. Experimental treatments included (I) water foliar application (control), (II) water + acetone foliar application (as a solvent), (III-V) 50, 100 and 200 JA  $\mu$ L. The essential oils obtained by hydro-distillation were analyzed by Gas Chromatography-Mass Spectrometry.

**Results & Discussion:** The results of GC-MS showed that the major components of the oil were *cis*-3-pinane (10-39%), *trans*-3-pinane (4-28%) and  $\beta$ -pinene (27-34%). The results of analysis of variance of the experiment showed that different levels of the foliar application of JA do have significant impacts on chemical constituents in the essential oil. Finally, foliar application of 200  $\mu$ L JA on some of secondary metabolite production in *H. officinalis* oil could be partially changed.

### Recommended applications/industries:

**Abbreviation:** JA: Jasmonic Acid, MJ: Methyl Jasmonate, GC-MS: Gas Chromatography-Mass Spectrometry  $\mu$ L: Micro liter, M: Molar; ANOVA: Analysis of Variance, CRD: Completely Randomized Design.

### 1. Introduction

Hyssop (*Hyssopus officinalis* L.) belongs to the family Lamiaceae family and is a native plant from southern

Europe and Near East to the region surrounding the Caspian Sea and cultivated in central and south European countries, including Russia, Spain, France, Yugoslavia, Netherland, Hungary and Italy (Mitić & Đorđević, 2000). This plant grows on dry banks and

among rocks and ruins with a height ranging from 50 to 120 cm (Le Strange, 1977; Könemann, 1999). The leaves are mainly used as an aromatic condiment (Baytop, 1997). *H. officinalis* has been known as a culinary and medicinal herb for hundreds of years. Hyssop oil finds its greatest use in flavoring preparations for alcoholic beverages, meat products and seasonings. It is used in tea blends for cough relief, antispasmodic effects, and relieving catarrh (Khazaie *et al.*, 2008).

*H. officinalis* var. *angustifolius* (Persian name: “Zoofa”) is grown and cultivated in some parts of Iran. The aerial parts of hyssop are used in Iranian folk medicine for their asthma, bronchitis, ulcers and wounds, carminative, antiseptic and antimicrobial (Zargari 1990; Ghasemi Pirbalouti, 2009). The essential oil and extracts isolated from *H. officinalis* have been shown to have biological and pharmacological activities, including anti-bacterial (Michalczyk *et al.*, 2012), anti-fungal (Fraternali *et al.*, 2004), anti-oxidant (Fernández-López *et al.*, 2003; Kizil *et al.*, 2010), sedative (Churl *et al.*, 2005), spasmolytic (Mazzanti *et al.*, 1998), anti-viral (Herrmann Jr & Kucera, 1967; Kreis *et al.*, 1990), cytotoxic (Renzini *et al.*, 1999), insecticidal (Pavela, 2004), and antiplatelet (Tognolini *et al.*, 2006). Kazazi *et al.* (2007) reported the main components of the extracts under different SFE conditions from *H. officinalis* cultivated in Iran were sabinene (4.2–17.1%), *iso*-pinocamphe (0.9–16.5%) and pinocamphe (0.7–13.6%).

There are several commercially available chemical compounds that could be used as elicitors to modify plant secondary metabolites and subsequently the bioactivity of medicinal plants. The most well-known chemical elicitors include salicylic acid (Métraux *et al.*, 1990; Van Wees *et al.*, 2000), jasmonic acid and its derivatives (Yazaki *et al.*, 1997; Sharan *et al.*, 1998; Vazquez-Flota and De Luca, 1998; Palazon *et al.*, 2003; Arimura *et al.*, 2005; Zheljazkov and Astatkie, 2012). Jasmonic acid and its volatile methyl ester, MJ, collectively termed jasmonates, are regarded as endogenous regulators that play important roles in regulating stress responses, plant growth and development (Creelman *et al.*, 1997).

The hypothesis of this study was that JA would have bioactivity or influence plant growth and development; hence, the JA could be used to promote or to limit plant growth. Few studies have been done to

investigate the effects of JA foliar application on the accumulation of secondary metabolites in medical plants in agricultural systems. Therefore, this study was done to evaluate the effect of various concentrations of JA on essential oil content and its composition of hyssop (*H. officinalis*) cultivated in greenhouse conditions.

## 2. Materials and Methods

### 2.1. Chemicals

Alkan standard solution C<sub>5</sub>-C<sub>24</sub> and jasmonic acid (JA) were purchased from Sigma–Aldrich Co. (Steineheim, Germany). Acetone and anhydrous sodium sulphate were bought from Merck Co. (Darmstadt, Germany).

### 2.2. Plant material and field site description

*H. officinalis* seeds were obtained from the Pakan Seed Company, Isfahan, Iran. In the third week of February 2012, seeds were sown in each plot (0.5 m x 0.5 m) with plant density of 12 (plants / m<sup>2</sup>). This experiment was conducted under plastic greenhouse conditions at Research Field, I.A.U., Shahrekord Branch, Chaharmahal va Bakhtiari province, southwestern Iran (latitude 32° 20' N, longitude 50° 51' E, altitude 2071 m above sea level). Plants were grown in a greenhouse in the relative humidity of 60-65% and, daily and nightly temperatures of 27±2 °C and 20±3 °C, respectively. No inorganic fertilizer and systemic pesticide were used during the experiment. For the first month (established phase), a watering level equivalent to 60-65% of soil field capacity was applied once a day. For the subsequent months, the plots were irrigated when 50% of soil available water was depleted (irrigation intervals varied from 2 or 3 days). The areal parts of hyssop were collected from each pot before flowering stage (10 June 2012), all of which were dried at 40 °C in the dark until it reached to a constant weight. Oil yield based matter dried (v/w) and chemical composition of essential oils were measured.

### 2.3. Treatments

Experimental treatments included (I) water foliar application (control), (II) water + acetone foliar application (as a solvent), (III-V) 50, 100 and 200 JA µL. Treatments were foliar-sprayed once at ten days before harvesting.

#### 2.4. Essential oil isolation

Fifty gram dried plant material were powered and subjected to hydro-distillation for 3 h using a Clevenger- type apparatus according to the method recommended in British pharmacopoeia ([British Pharmacopoeia, 1988](#)). Samples were dried with anhydrous sodium sulphate and kept in amber vials at  $4\pm 1$  °C prior to use.

#### 2.5. GC/MS analysis

The essential oils were analyzed using an Agilent 7890, a gas chromatograph (Agilent Technologies, Palo Alto, CA, USA) with a HP-5MS 5% phenylmethylsiloxane capillary column (30 m x 0.25 mm, 0.25  $\mu$ m film thickness). Oven temperature was kept at 60 °C for 4 min initially, and then, raised at the rate of 4 °C/min to 260 °C. Injector and detector temperatures were set at 290 and 300 °C, respectively. Helium was used as carrier gas at a flow rate of 2 mL/min, and 0.1  $\mu$ L samples were injected manually in the split mode. Peaks area percentages were used for obtaining quantitative data. The gas chromatograph was coupled to an Agilent 5975 C (Agilent Technologies, Palo Alto, CA, USA) mass selective detector. The EI-MS operating parameters were as follows: ionization voltage, 70 eV; ion source temperature, 200 °C. Retention indices were calculated for all components using a homologous series of *n*-alkanes (C<sub>5</sub>-C<sub>24</sub>) injected in conditions equal to samples ones. Identification of oil components was accomplished based on comparison of their retention times with those of authentic standards and by comparison of their mass spectral fragmentation patterns (WILLEY/ChemStation data system ([Adams, 2007](#); [McLafferty, 2009](#))).

#### 2.6. Statistical analysis

The data was statistically analyzed using a CRD by the program SPSS (17.0). Means of the characteristics were compared by Duncan's multiple range test at  $P < 0.05$  level.

### 3. Results and Discussion

The color of extracted oil was yellow. The oil yields of studied treatments ranged between 0.77 to 0.86% (v/w), based on dry weight. Statistical analysis indicated that there was no significant difference

between treatments for oil yield (Table 1). There also was no significant difference between treatments for total dry matter of hyssop.

The chemical constituents identified by GC-MS, are presented in Table 1. Totally, thirty eight components were identified representing more than 85-99% of the oil composition. The analysis of essential oils detected major compounds, viz. *cis*-3-pinanone, *trans*-3-pinanone,  $\beta$ -pinene,  $\beta$ -phellandrene, *trans*-caryophyllene, myrtenol, germacrene D, elemol, bicyclogermacrene,  $\alpha$ -pinene,  $\beta$ -myrcene, sabinene and pinocarvone (Table 1 and Fig 1-3). In this study, the essential oils obtained from hyssop contained oxygenated monoterpenes, monoterpene hydrocarbons and sesquiterpenes. Similarly, Ozer *et al.* (2005) reported that the major constituents in the essential oil obtained by hydro-distillation from the aerial parts of *H. officinalis* L. subsp. *angustifolius* (Bieb.) collected from Turkey were pinocarvone (36.3%), pinocamphone (19.6%),  $\beta$ -pinene (10.6%), 1,8-cineole (7.2%), and isopinocampone (5.3%). The results of earlier studies indicated the major volatile constituents obtained from the aerial parts of *H. officinalis* were sabinene, *iso*-pinocampone, pinocamphone, camphor and  $\beta$ -pinene,  $\alpha$  and  $\beta$ -phellandrene, germacrene D, myrtenol and pinocampone ([Garg \*et al.\*, 1999](#); [Mitić & Đorđević, 2000](#); [Chalchat \*et al.\*, 2001](#); [Fraternale \*et al.\*, 2004](#); [Khazaie \*et al.\*, 2008](#)).

The result of analysis of variance indicated that different levels of the foliar application of JA had a significant impact on the main constituents in the essential oils of hyssop (Table 1). Percentages *cis*-3-pinanone, *trans*-3-pinanone,  $\beta$ -pinene,  $\beta$ -phellandrene, myrtenol, and  $\alpha$ -pinene amounts in oils of *H. officinalis* under foliar application 200  $\mu$ L JA decreased (Fig 1-3). Decreased amount of these constituents in plants grown under JA (200  $\mu$ L) foliar application might be attributed to stress condition, which would activate the synthesis of secondary metabolites. On the other hand, content of sesquiterpenes group including *trans*-caryophyllene, germacrene D, elemol and bicyclogermacrene increased under JA 200  $\mu$ L treatment (Table 1). The results of present study revealed that the foliar application of JA 200  $\mu$ L caused significantly increased sesquiterpenes in comparison with monoterpenes.


JA and MJ are known as transducers of elicitor signal transduction that results in the biosynthesis of plant secondary metabolites. They act as inducers of

phytoalexins (Zhao *et al.*, 2005). Accumulation of isoflavonoids, as phytoalexins, was enhanced in vitro cultures of *Pueraria lobata* (Willd.) Ohwi (Thiem and Krawczyk, 2010). Ashrafi *et al.* (2012) reported that the foliar application of JA (100  $\mu$ L) have increased thymol and carvacrol contents in essential oil obtained from *Thymus daenensis* aerial parts. JA and its derivatives are known to stimulate production of secondary metabolites in plants (Sanz *et al.*, 2000). JAs have been shown to regulate the synthesis of various secondary metabolites including caffe-oylputrescine in tomato leaves (Chen *et al.*, 2006), anthocyanins (Zhao *et al.*, 2005; Uppalapati *et al.*, 2005) and defense-related volatiles (Thaler *et al.*, 2002; Ament *et al.*, 2004). JA induces the production of ajmalicine and catharanthine in *Catharanthus roseus* (Vazquez-Flota and De Luca, 1998), rosmarinic acid and shikonin in *Lithospermum erythrorhizon* (Yazaki *et al.*, 1997), scopoletin and scopolin in *Nicotiana tabacum* (Sharan *et al.*, 1998) and taxol and paclitaxel in *Taxus* sp. cell suspensions (Palazon *et al.*, 2003). Induction of plant secondary metabolite accumulation by the JA signaling pathway is not limited to certain types of metabolites, but includes a wide variety of plant secondary products including terpenoids, flavonoids, alkaloids, and phenylpropanoids plus many other types of secondary metabolites in most plants (Zhao *et al.*, 2005). Therefore, the JA signaling pathway is generally regarded as an integral signal for biosynthesis of many plant secondary products (Zhao *et al.*, 2005). Increasing evidence indicates that JA-induced changes in secondary metabolism constitute a ubiquitous plant defense response (Gundlach *et al.*, 1992; Memelink *et al.*, 2001; Goossens *et al.*, 2003; Zhao *et al.*, 2005).

JA has also been implicated as the signal molecule responsible for increased synthesis of nicotine, an insecticidal compound produced by *Nicotiana sylvestris* upon leaf wounding (Zheng *et al.*, 1997). JAs are produced and accumulated in plants, but exogenous application of JA and MJ can elicit secondary metabolite accumulation in defense response induction (Thiem and Krawczyk 2010).

#### 4. Conclusion

Plant secondary metabolites are unique sources for pharmaceuticals, food additives, flavors and other industrial materials. Accumulation of such metabolites often occurs in plants subjected to stresses including various elicitors or signal molecules. Jasmonic and salicylic acids compounds have long been observed to be transducers of elicitor signals for the production of plant secondary metabolites. JA (200  $\mu$ L) could be used as foliar application in *H. officinalis* for increasing phenolic components including carvacrol and thymol. Phenolic components production in lemon balm could be partially changed by supplementation of different elicitors.




**Fig 1.** Effect of foliar application of JA on  $\beta$ -pinene content


**Table 1.** Effect of JA foliar application on some of main components contents in *H. officinalis* oil

Treatments	$\beta$ -phellandrene	$\beta$ -Myrcene	Pinocarvone	Myrtenol	$\beta$ -Caryophyllene	Germacrene D	Bicyclo-germacrene	Elemol
JA50	2.36	2.29	4.16	3.82	1.49	3.11	0.00	0.00
JA100	2.54	2.44	2.01	3.77	1.57	3.24	2.73	0.00
JA200	0.56	0.99	2.01	0.73	5.82	13.19	10.17	17.26
Water	6.10	1.95	0.00	2.95	1.45	3.10	2.60	0.00
Acetone	0.00	1.15	0.00	3.17	1.03	0.00	0.00	2.98





**Fig 2.** Effect of foliar application of JA on *trans*-3-pinanone content



**Fig 3.** Effect of foliar application of JA on *cis*-3-pinanone content

## References

- Abbaszadeh, B., Sharifi Ashourabadi, E., Ardakani, M.R. and Paknejad, F. 2005. Effect of spraying of nitrogen fertilizer on *Melissa officinalis* L yield in the greenhouse condition. *Iran J. Med Aromat Plants.*, 21: 207-216.
- Adams, R.P., 2007. *Identification of essential oil components by gas chromatography/mass spectrometry*, 4th Ed., p. 456, Allured Publishing Corporation, Carol Stream, IL.
- Akhondzadeh, S., Noroozian, M., Mohammadi, M., Ohadinia, M., Jamshidi, A.H. and Khani, K. 2003. *Melissa officinalis* extract in the treatment of patients with mild to moderate Alzheimer's disease: a double blind, randomised, placebo controlled trial. *J Neurol Neurosurg Psychiatry.*, 74: 863-866.
- Allahverdiyev, A., Duran, N., Ozguven, M. and Koltas, S. 2004. Antiviral activity of the volatile oils of *Melissa officinalis* L. against Herpes simplex virus type-2. *Phytomedicine.*, 11: 657-661.
- Ament, K., Kant, M.R., Sabelis, M.W., Haring, M.A. and Schuurink, R.C. 2004. Jasmonic acid is a key regulator of spider mite-induced volatile terpenoid and methyl salicylate emission in tomato. *Plant Physiol.*, 135: 2025-2037.
- Amin, G.R., 1991. *Popular medicinal plants of Iran*, Ministry of Health Publication, Tehran.
- Arimura, G. and Kost Boland, W. 2005. Herbivore-induced, indirect plant defences. *Biochimica et Biophysica Acta.*, 1734: 91-111.
- Asgari, F. and Sefidkon, F. 2004. Essential oil composition of *Melissa officinalis* L. From different region. *Iran J. Med Aromatic Plants.*, 20: 229-237.
- Ashrafi, M., Ghasemi Pirbalouti, A., Rahimmalek, M. and Hamed, B. 2012. Effects of foliar application of jasmonic acid (JA) on the essential oil chemical composition of *Thymus daenesis* Celak. *J Herbal Drugs.*, 3: 75-90.
- Bayat, M., Azami Tameh, A., Ghahremani, M., Akbari, M., Ejtemaei Mehr, S., Khanavi, M. and Hassanzadeh, G. 2012. Neuro-protective properties of *Melissa officinalis* after hypoxic-ischemic injury both in vitro and in vivo. *DARU J Pharm Sci.*, 20: 42.
- Bisset, N.G. and Wichtl, M. 2001. *Herbal Drugs and Phytopharmaceuticals*, CRC Press: Boca Raton, London, New York, and Washington DC.
- British Pharmacopoeia Commission, 1988. *British Pharmacopoeia*. Her Majesty's Stationery Office, London, pp: 137-138.
- Canadanovic-Brunet, J., Cetkovic, G., Djilas, S., Tumbas, V., Bogdanovic, G., Mandic, A., et al. 2008. Radical scavenging, antibacterial, and antiproliferative activities of *Melissa officinalis* L. Extracts. *J Med Food.*, 11: 133-143.
- Carnat, A.P., Carnat, A., Fraisse, D. and Lamaison, J.L. 1998. The aromatic and polyphenolic composition of lemon balm (*Melissa officinalis* L, subsp *officinalis*) tea. *Pharm Acta Helv.*, 72: 301-305.
- Chen, H., Jones, A.D. and Howe, G.A. 2006. Constitutive activation of the jasmonate signaling pathway enhances the production of secondary metabolites in tomato. *FEBS Lett.*, 580: 2540-2546.
- Chung, M.J., Cho, S.Y., Bhuiyan, M.J., Kim, K.H. and Lee, S.J. 2010. Anti-diabetic effects of lemon balm (*Melissa officinalis*) essential oil on glucose- and lipidregulating enzymes in type 2 diabetic mice. *Brit J Nutr.*, 104: 180-188.
- Cohen, R.A., Kucera, L.S. and Herrmann Jr, E.C. 1964. Antiviral activity of *Melissa officinalis* (lemon Balm) extract. *P Soc Exp Biol Med.*, 117: 431-434.
- Creelman, R., Tierney, M and Mullet, J. 1997. Jasmonic acid/methyl Jasmonate accumulate in wounded soybean hypocotyle and modulate wound

- gene expression. *Proc. Natl. Acad. Sci. U.S.A.*, 89: 4938–4941.
- Fraternale, D., Ricci, D., Epifano, F. and Curini, M. (2004). Composition and antifungal activity of two essential oils of hyssop (*Hyssopus officinalis* L.). *J. Essent Oil Res.*, 16(6), 617-622.
- Dastmalchi, K., Dorman, H.J.D., Oinonen, P.P., Darwis, Y., Laakso, I. and Hiltunen, R. 2008. Chemical composition and *In vitro* antioxidative activity of a lemon balm (*Melissa officinalis* L.) extract. *LWT-Food Sci Technol.*, 41: 391-400.
- Dastmalchi, K., Ollilainen, V., Lackman, P., Boije af Gennäs, G., Dorman, H.J. et al. 2009. Acetylcholinesterase inhibitory guided fractionation of *Melissa officinalis* L. *Bioorg. Med. Chem.*, 17: 867–871.
- de Sousa, A.C., Gattass, C.R., Alviano, D.S., Alviano, C.S., Blank, A.F. and Alves, P.B. 2004. *Melissa officinalis* L. essential oil: antitumoral and antioxidant activities. *J. Pharm. Pharmacol.*, 56: 677–681.
- Fernández-López, J., Sevilla, L., Sayas-Barberá, E., Navarro, C., Marín, F. and Pérez-Alvarez, J.A. 2003. Evaluation of the Antioxidant Potential of Hyssop (*Hyssopus officinalis* L.) and Rosemary (*Rosmarinus officinalis* L.) extracts in Cooked Pork Meat. *J. Food Sci.*, 68: 660–664.
- Renzini, G., Scazzocchio, F., Lu, M., Mazzanti, G. Salvatore, G. 1999. Antibacterial and cytotoxic activity of *Hyssopus officinalis* L. oils. *J. Essent Oil Res.*, 11(5), 649-654.
- Garg, S.N., Naqvi, A.A., Singh, A., Ram, G. and Kumar, S. 1999, Composition of essential oil from an annual crop of *Hyssopus officinalis* grown in indian plains. *Flavour Fragr. J.*, 14: 170–172.
- Gharib, F.A.E., 2006. Effect of salicylic acid on the growth, metabolic activities and oil content of basil and marjoram. *Int J Agr Biol.*, 4: 485–492.
- Gharib, F.A.E., 2007. Effect of salicylic acid on the growth, metabolic activities and oil content of basil and marjoram. *Int J Agr Biol.*, 9: 294-301.
- Ghassemi-Dehkordi, N, 2002. *Iranian Herbal Pharmacopoeia*. Ministry of Health and Medical Education, Tehran, Iran, pp. 468-469.
- Goossens, A., Häkkinen, S.T., Laakso, I., Seppanen-Laakso, T., Biondi, S., et al. 2003. A functional genomics approach toward the understanding of secondary metabolism in plant cells. *Proc Nath Acad Sci USA.*, 100: 8595–8600.
- Gundlach, H., Müller, M.J., Kuchan, T.M. and Zenk, M.H. 1992. Jasmonic acid is a signal transducer in elicitor-induced plant cell cultures. *Proc Nath Acad Sci USA.*, 89: 2389–2393.
- Heitz, A., Carnat, A., Fraisse, D., Carnat, A.P. and Lamaison, J.L. 2000. Luteolin 3'-glucuronide, the major flavonoid from *Melissa officinalis* subsp. *officinalis*. *Fitoterapia.*, 71: 201-202.
- Hohmann, J., Zupko, I., Redei, D., Csanyi, M., Falkay, G., Mathe, I. and Janicsak, G. 1999. Protective effects of the aerial parts of *Salvia officinalis*, *Melissa officinalis* and *Lavandula angustifolia* and their constituents against enzyme-dependent and enzyme-independent lipid peroxidation. *Planta Med.*, 65: 576-578.
- Ivanova, D., Gerova, D., Chervenkov, T. and Yankova, T. 2005. Polyphenols and antioxidant capacity of Bulgarian medicinal plant. *J Ethnopharmacol.*, 96: 145–150.
- Khazaie HR, Nadjafi F. and Bannayan M. 2008. Effect of irrigation frequency and planting density on herbage biomass and oil production of thyme (*Thymus vulgaris*) and hyssop (*Hyssopus officinalis*). *Ind Crops Prod.*, 27: 315-321.
- Koliopoulos, G., Pitarokili, D., Kioulos, E., Michaelakis, E. and Tzakou, O. 2010. Chemical composition and larvicidal evaluation of *Mentha*, *Salvia*, and *Melissa* essential oils against the West Nile virus mosquito *Culex pipiens*. *Parasitol Res.*, 107: 327–335.
- Koytchev, R., Alken, R.G. and Dundarov, S. 1999. Balm mint extract (Lo-701) for topical treatment of recurring herpes labialis. *Phytomedicine.*, 6: 225–230.
- Kreis, W., Kaplan, M.H., Freeman, J., Sun, D.K., Sarin, P.S. 1990. Inhibition of HIV replication by Hyssop officinalis extracts. *Antiviral. Res.*, 14: 323-337.
- Leung, A.Y. and Foster, S. 1996. *Encyclopedia of common natural ingredient used in food, drugs and cosmetics*. John Wiley & Sons, Inc, New York.
- Michalczyk, M., Macura, R., Tesarowicz, I., and Banaś, J. 2012. Effect of adding essential oils of coriander (*Coriandrum sativum* L) and hyssop (*Hyssopus officinalis* L.) on the shelf life of ground beef. *Meat Science*, 90(3), 842-850.
- Malatyali, E., Tepe, B., Degerli, S. and Berk, S. 2012. In vitro amoebicidal activities of *Satureja cuneifolia*

- and *Melissa officinalis* on *Acanthamoeba castellanii* cysts and trophozoites. *Parasitol Res.*, 110: 2175–2180.
- Masakova, N.S., Tseevatuy, B.S., Trofimenko, S.L. and Remmer, G.S. 1979. The chemical composition of volatile oil in lemon-balm as an indicator of therapeutic use. *Planta Med.*, 36: 274.
- Mazzanti, G., Lu, M., Salvatore, G., 1998. Spasmolytic action of the essential oil from *Hyssopus officinalis* L. var. *decumbens* and its major components. *Phytothera. Res.*, 12: S92-S94.
- McLafferty, F.W. 2009. *Wiley Registry of Mass Spectral Data*, 9<sup>th</sup> ed. Wiley, Inc., Hoboken, NJ.
- Memelink, J., Verpoorte, R. and Kijne, J.W. 2001. ORCANisation of jasmonate-responsive gene expression in alkaloid metabolism. *Trends Plant Sci.*, 6: 212–219.
- Métraux, J.P., Signer, H., Ryals, J., Ward, E., Wyss-Benz, M., Gaudin, J., et al. 1990. Increase in salicylic acid at the onset of systemic-acquired resistance in cucumber. *Science.*, 250: 1004–1006.
- Mimica-Dukic, N., Bozin, B., Sokovic, M. and Simin, N. 2004. Antimicrobial and antioxidant activities of *Melissa officinalis* L. (Lamiaceae) essential oil. *J Agric Food Chem.*, 52: 2485-2489.
- Miroliaei, M., Khazaei, S., Moshkelgosha, S. and Shirvani, M. 2011. Inhibitory effects of Lemon balm (*Melissa officinalis* L.) extract on the formation of advanced glycation end products. *Food Chem.*, 129: 267–271.
- Mozaffarian, V. 2008. *A dictionary of Iranian plant names*. Farhang Mosavar Press, Tehran, Iran.
- Ozer, H., Şahin, F., Kılıç, H. and Güllüce, M. 2005. Essential oil composition of *Hyssopus officinalis* L. subsp. *angustifolius* (Bieb.) Arcangeli from Turkey. *Flavour Fragr. J.*, 20: 42–44.
- Palazon, J., Cusidó, R.M., Bonfill, M., Morales, C. and Piñol, M.T. 2003. Inhibition of paclitaxel and baccatin III accumulation by mevinolin and fosmidomycin in suspension cultures of *Taxus baccata*. *J Biotechnol.*, 101: 157–163.
- Pavela R. 2004. Insecticidal activity of certain medicinal plants. *Fitoterapia.*, 75: 745-749.
- Sadraei, H., Ghannadi, A. and Malekshahi, K. 2003. Relaxant effect of essential oil of *Melissa officinalis* and citral on rat ileum contractions. *Fitoterapia.*, 74: 445–452.
- Sanz, M.K., Hernandez, X.E., Tonn, C.E. and Guerreiro, E. 2000. Enhancement of tessaric acid production in *Tessaria absinthioides* cell suspension cultures. *Plant Cell Rep.*, 19: 821–824.
- Sharafzadeh, S., Khosh-Khui, M. and Javidnia, K. 2011. Effect of nutrients on essential oil components, pigments and total phenolic content of lemon balm (*Melissa officinalis* L.). *Adv Environ Res.*, 5: 639-646, 2011.
- Sharan, M., Taguchi, G., Gonda, K., Jouke, T., Shimosaka, M., Hayashida, N. and Okazaki, M. 1998. Effects of methyl jasmonate and elicitor on the activation of phenylalanine ammonialyase and the accumulation of scopoletin and scopolin in tobacco cell cultures. *Plant Sci.*, 132: 13–19.
- Szabo, E., Thelen, A. and Petersen, M. 1999. Fungal elicitor preparations and methyl jasmonate enhance rosmarinic acid accumulation in suspension cultures of *Coleus blumei*. *Plant Cell Rep.*, 18: 485–489.
- Taherpoura, A., Maroofi, H., Rafie, Z. and Larijani, K. 2012. Chemical composition analysis of the essential oil of *Melissa officinalis* L. from Kurdistan, Iran by HS/SPME method and calculation of the bio-physicochemical coefficients of the components. *Nat Prod Rep.*, 26: 152-160.
- Thaler, J.S., Farag, M.A. Pare, P.W. and Dicke, M. 2002. Jasmonate-deficient plants have reduced direct and indirect defences against herbivores. *Ecol Lett.*, 5: 764–774.
- Thiem, B. and Krawczyk, A. 2010. Enhanced isoflavones accumulation in methyl jasmonate-treated *in vitro* cultures of kudzu (*Pueraria lobata* Ohwi). *Herba Polon.*, 56: 48-56.
- Triantaphyllou, K., Blekas, G. and Boskou, D. 2001. Antioxidant properties of water extracts obtained from herbs of the species Lamiaceae. *Int J Food Sci Nutr.*, 52: 313–317.
- Uppalapati, S.R., Ayoubi, P., Weng, H., Palmer, D.A., Mitchell, R.E., Jones, W. and Bender, C.L. 2005. The phytotoxin coronatine and methyl jasmonate impact multiple phytohormone pathways in tomato. *Plant J.*, 42: 201–217.
- Van Wees, S.C.M. and Glazebrook, J. 2003. Loss of non-host resistance of *Arabidopsis NahG* to *Pseudomonas syringae* pv. *phaseolicola* is due to degradation products of salicylic acid. *Plant J.*, 33: 733–742.
- Vazquez-Flota, A. and De Luca, V. 1998. Jasmonate modulates development and light-regulated alkaloid biosynthesis in *Catharanthus roseus*. *Phytochemistry.*, 49: 395–402.

- Venkutonis, P.R., Gruzdiene, D., Trizite, D. and Trizite, G. 2005. Assessment of antioxidant activity of plant extracts by different methods. *Acta Horticulturae.*, 677: 99–107.
- Walker, T.S., Bais, H.P. and Vivanco, J.M. 2002. Jasmonic acid-induced hypericin production in cell suspension cultures of *Hypericum perforatum* L. (St. John's wort). *Phytochemistry.*, 60: 289–293.
- Yazaki, K., Takeda, K. and Tabata, M. 1997. Effects of methyl jasmonate on shikonin and dihydro-echinofuran production in *Lithospermum* cell cultures. *Plant Cell Physiol.*, 38: 776–782.
- Zargari, A., 1992. *Medicinal plants*. Tehran University Press, Tehran, Iran.
- Zhao, J., Davis, L.C. and Verpoorte, R. 2005. Elicitor signal transduction leading to production of plant secondary metabolites. *Biotechnol Adv.*, 23: 283-333.
- Zheljazkov, V.D. and Astatkie, T. 2012. Distillation waste water can modify peppermint (*Mentha × piperita* L.) oil composition. *Ind Crop Prod.*, 36: 420–426.
- Zheng, Z.P., Krumm, T. and Baldwin, I.T. 1997. Structural requirements of jasmonates and mimics for nicotine induction in *Nicotiana sylvestris*. *J Cheml Ecol.*, 23: 777–2789.

Archive of SID