



Effect of the essential oil of thyme on intestinal morphology in Rat

Heshmat Sepehri Moghadam*, Mozhdeh Emadi, Elham Molashahi

Department of Agricultural Science, Payam Noor University, P.O. Box 19395-4697 Tehran, Iran;

*Email: he_sepehri@yahoo.com

ARTICLE INFO

Type: Original Article

Topic: Ethnopharmacology

Received 25th May 2014

Accepted 9th June 2014

Key words:

- ✓ Indomethacin
- ✓ Small intestine
- ✓ *Thymus vulgaris*
- ✓ Volatile oil

ABSTRACT

Background & Aim: Non-steroidal anti-inflammatory drugs (NSAIDs) are commonly used in clinical medicine. The adverse effects they produce in the gastrointestinal tract however, often limit their utility. Oxidative stress has been shown to occur in the small intestine in response to the oral administration of indomethacin. In view of this, the effect of thyme essential oil, an agent with anti-oxidant properties, was evaluated on indomethacin-induced small intestinal damage in a rat model.

Experimental: The chemical composition of thyme essential oil was identified by gas chromatography mass spectrometry. Tissues morphology affected by indomethacin, with and without treatment with thyme, was measured.

Results & Discussion: The small intestine morphology (villus height, villus width, and crypt depth and crypt width of duodenum, jejunum and ilea) from indomethacin-treated animals with thyme oil was significantly improved than those from indomethacin animals. Treatment with thyme oil was found to ameliorate these drug-induced changes.

Recommended applications/industries: The essential oil of thyme appears to hold promise as an agent that can potentially reduce non-steroidal anti-inflammatory drugs induced small intestinal damage.

1. Introduction

Thymus vulgaris L. (thyme) is an aromatic plant belonging to the Lamiaceae family that has been used traditionally as condiments and flavoring agents. Thyme is also used in folk medicine as remedies for different ailments of the digestive tract. Most of its bioactive effects come from essential oils, thymol and carvacrol. In addition to its flavouring role, thymol is commonly used in food and cosmetic industry as preservative and antioxidant. It is also used in dentistry as a protective agent against caries and plaques and in anesthesiology as a preservative and stabilizer of halothane (Boudry & Perrier, 2008). Thymol is also a

potent scavenger of reactive oxygen species (Kruk et al., 2000) and has anti-inflammatory properties by inhibiting COX1 activity or release by neutrophils (Braga et al., 2006). Thyme essential oil could be used as a growth promoter due to its mode of action, including: improving feed flavor and intake, stimulating the secretion of digestive enzymes, increasing gastric and intestinal motility, gut development and antimicrobial activity (Jamroz et al., 2006; Tipu et al., 2006). Carvacrol, an active component of thyme essential oil, has been noted to have a positive effect on intestinal morphology and secretion of mucin in the intestine (Jamroz et al., 2006). Non-steroidal anti-inflammatory drugs (NSAID) are a class of drugs that usually used for the treatment

of acute and chronic conditions where pain and inflammation are present. Non-steroidal anti-inflammatory drug inhibit the activity of both cyclooxygenase-1 (COX-1) and cyclooxygenase-2 (COX-2) and thereby, the synthesis of prostaglandins and thromboxans. It is thought that inhibiting COX-2 leads to the anti-inflammatory analgesic and antipyretic effects and that these NSAID also inhibiting COX-1 may cause gastrointestinal bleeding and ulcers. The main adverse drug reactions associated with NSAID uses related to direct and indirect irritation of the gastrointestinal tract. Nonsteroidal anti-inflammatory drug cause a dual assault on the gastrointestinal tract: the acidic molecules directly irritate the gastric mucosa and inhibition of COX-1 and COX-2 reduces the levels of protective prostaglandins. Inhibition of prostaglandin synthesis in the gastrointestinal tract causes increased gastric acid secretion, diminished bicarbonate secretion, diminished mucus secretion and diminished tropic effects on epithelial mucosa. Nonsteroidal anti-inflammatory drug induced injury on gastrointestinal tract is well documented, and jejunal inflammation caused by indomethacin in rats is a broadly used experimental model of enteritis (Menozzi *et al.*, 2009). The pathogenesis of non-steroidal anti-inflammatory drugs induced small intestinal lesions remains unclear, although it is considered to be quite different from that of upper gastrointestinal tract ulcers due to the absence of acid and the presence of bacteria and bile in the small intestine (Yamada *et al.*, 2012). The aim of this study was to show the effect of indomethacin on small intestinal morphology, and to assess the role of thyme in the indomethacin-induced changes.

2. Materials and Methods

2.1. Animals and design

In this study, 16 rats were chosen that treatment groups were comprised of eight groups of animals treated with thyme volatile oil dissolved in distilled water to the desired concentration. Thyme volatile oil dissolved in distilled water was administered once daily (1 ml). Rats were divided randomly into four groups: Group A: received intramuscular injection of indomethacin in male rat, Group B: was received intramuscular injection of indomethacin in female rat, Group C: received intramuscular injection of indomethacin and thyme volatile oil in male rat and

Group D: received intramuscular injection of indomethacin and thyme volatile oil in female rat. Animals were housed in individual cages and kept in a ventilated room with temperature regulated at 23 °C, with a 12 h light and 12 h dark cycle. They were fed with commercial rat pellets and water was supplied *ad libitum* and were acclimatized for 2 days prior to experimentation.

2.2. Extraction of thyme volatile oils

Fresh plants were collected at the flowering stage and processed immediately after harvest. Volatile oil was distilled from the ground plant material using Clevenger distillation apparatus (Herbal Exir Co., Mashhad, Iran). The samples were distilled for 2 h and the oils obtained, dried with anhydrous sodium sulphate, and stored in dark sealed glass vials at 4 °C until required. The main active compounds of the thyme were determined by GC-MS.

2.3. Anesthetic protocol

The animals were anesthetized using 0.05 ml inspiration chloroform and after those animals injected with intramuscular of indomethacin. The dose of Indomethacin (8 mg/kg) was determined from a previous study (Abimosleh *et al.*, 2013).

2.4. Data collection

At end of experiment, rats were sacrificed by CO₂ asphyxiation followed by cervical dislocation and then the mid part of duodenum, jejunum and ileum were excised for histomorphometric analysis. Briefly, the small intestine was divided into three segments: the duodenum, the jejunum and the ileum. The duodenum is recognizable as the first stretch of the intestine leading from the stomach, it is mostly straight. The jejunum and ileum are both curly parts of the intestine, with the ileum being the last section before the small intestine becomes the large intestine. Samples of duodenum, jejunum and ileum (0.5 cm × 0.5 cm segments) were obtained at its midpoint and immersed in a 10% buffered formalin solution for 72 h. Then they were excised and washed by physiological saline. The samples were treated in tissue processor apparatus and embedded in paraffin wax (Bancroft & Gamble, 2002).

Transverse sections were cut (6 μm) using a rotary microtome (LEICA RM 2145), placed on a glass slide and stained with hematoxylin and eosin, and analyzed

under a light microscope to determine morphometric indices. Morphological parameters were measured using the Image Pro Plus v 4.5 software package. The measured morphometric variables (Sakamoto *et al.*, 2000; Aptekmann *et al.*, 2001) included: villus height (VH) measured from the villus-crypt junction; villus width (VW) measured at midvillus height; crypt depth (CD) and crypt width (CW). The mean from 10 villus per sample was used as the average value for further analysis.

2.5. Statistical analysis

All data was checked for normality before analysis. The data was subjected to ANOVA using SAS (SAS, 2000). Differences between treatment means were evaluated by Duncan's multiple range tests. A value of $p < 0.05$ was considered significant.

3. Results and discussion

Results of the essential oil analysis indicated that thyme oil contained thymol and its isomer, carvacrol at the rate of 21.9 and 31.9%, respectively. The concentrations of two predominate components of thyme volatile oils; thymol and carvacrol have been reported to range from as low as 3% to as high as 60% of total essential oils (Khaksar *et al.*, 2013).

Effects of feeding diets containing thyme volatile oils on small intestinal morphology (duodenum, jejunum and ilea) are shown in Table 1, 2 and 3. The thyme volatile oil significantly improved small intestinal morphology in indomethacin-treated with thyme rats when compared to the indomethacin rat groups. Results of duodenum and jejunum in rats were affected by inclusion of thyme volatile oil into the diets. VH, VW, CD and CW significantly improved in rats fed thyme volatile oil in duodenum and jejunum. However, the only significant difference in the ilea was in VH and CD. Intestinal villi are the main site of nutrient absorption and their better development could be the reason for higher nutrient absorption (Hanczakowska & Swiatkiewicz, 2012). Velazques *et al.* (2005) reported positive effects of herb extract on the development of the intestinal tract. As mentioned above, there is a relationship between villi height and nutrient absorption and digestibility (Hanczakowska & Swiatkiewicz, 2012).

Table 1. Effect of thyme volatile oil on duodenum morphology

Treatment*	VH	VW	CD	CW
A	129.21 ^c	27.72 ^b	68.24 ^a	42.29 ^a
B	118.59 ^c	20.52 ^b	59.46 ^b	34.74 ^a
C	198.41 ^a	46.53 ^a	46.93 ^b	14.62 ^b
D	142.94 ^b	35.62 ^a	37.57 ^b	16.00 ^b
P value**	0.001	0.020	0.002	0.001
SEM	0.80	0.82	1.12	0.60

*A: injection of indomethacin in male rat

B: injection of indomethacin in female rat

C: injection of indomethacin and thyme volatile oil in male rat

D: injection of indomethacin and thyme volatile oil in female rat

**a,b,c Means in each column with different superscripts are significantly different ($p < 0.05$).

Table 2. Effect of thyme volatile oil on jejunum morphology

Treatment*	VH	VW	CD	CW
A	111.87 ^b	33.25 ^c	58.89 ^c	25.73 ^b
B	104.26 ^b	33.53 ^c	53.47 ^c	36.17 ^a
C	179.09 ^a	44.08 ^a	87.88 ^a	38.37 ^a
D	161.87 ^a	39.66 ^b	70.46 ^b	23.42 ^b
P value**	0.001	0.003	0.001	0.001
SEM	15.03	0.43	0.57	2.71

*A: injection of indomethacin in male rat

B: injection of indomethacin in female rat

C: injection of indomethacin and thyme volatile oil in male rat

D: injection of indomethacin and thyme volatile oil in female rat

**a,b,c Means in each column with different superscripts are significantly different ($p < 0.05$).

Table 3. Effect of thyme volatile oil on ilea morphology

Treatment*	VH	VW	CD	CW
A	158.31 ^b	35.42	74.27 ^b	36.14
B	151.63 ^b	35.14	68.12 ^b	34.87
C	208.98 ^a	39.95	109.83 ^a	41.00
D	196.87 ^a	39.56	94.55 ^a	36.14
P value**	0.018	0.520	0.011	0.309
SEM	1.32	0.38	1.14	0.58

*A: injection of indomethacin in male rat

B: injection of indomethacin in female rat

C: injection of indomethacin and thyme volatile oil in male rat

D: injection of indomethacin and thyme volatile oil in female rat

**a,b,c Means in each column with different superscripts are significantly different ($p < 0.05$).

Pappenheimer & Michel (2003) proved a decisive role of villi in the intestinal absorption of nutrients. Increasing villi height suggests an increased surface area for greater absorption of available nutrients. In addition, increase in villi to crypt ratio is related to an increase in digestion and absorption (Montagne *et al.*, 2003).

Antimicrobial agents such as essential oils or their active components are known to reduce the intestinal microbial load, which in turn reduces the presence of toxins that are associated with changes in intestinal histomorphology, such as shorter villi and deeper crypts (Xu *et al.*, 2003). As regards that productive potential of the cell cycle in small intestine is resultant of cell proliferation, cell migration in pivot crypt-villi, death and cell ablation, then, digestive material with high viscosity in lumen can be caused loss of cell's villi, decrease villi high and increase production of cell crypt and finally increase in crypt death and produce deeper crypts. However, hydro-alcoholic plant extracts from sage, thyme, and rosemary leaves did not exert any influence on intestinal villus height, villus surface area and crypt depth. Longer villi increase the absorptive surface of intestine, while smaller crypt indicates a decrease of enterocyte replacement and tissue turnover, and lower demand for tissue development as well. It can be stated that, increments in villus height and crypt death as directly correlated with enhanced epithelial cell turnover (Fang *et al.*, 1997). In contrary, some studies reported that there was no significant effect in the villus height, crypt death and villus surface area due to feeding 200 ppm of plant extract, based on a blend of oregano, cinnamon and pepper essential oils and 5000 ppm of hydro-alcoholic extract from sage, thyme and rosemary leaves. Guo *et al.* (2004) found no significant differences in intestinal morphology among a medicinal plant group and control group. Garcia *et al.* (2007) also reported no significant differences in villus height and crypt death between birds in the control group and a group receiving a blend of plant extracts. However, little information is available on how phytochemical compounds may affect gastrointestinal histomorphology and functionality. Observations were reported by Jamroz *et al.* (2006) who observed qualitative increase in the number of goblet cells and in mucin secretion at the surface of the jejunum villi when feeding broilers a mixture of 5 mg/kg of carvacrol. The positive effect of extracts from medicine plants on piglet villi height was reported also by Fang *et al.* (2009). Now we do not have a satisfactory explanation of the beneficial effect of herbal extracts and their antioxidative activity on ileal structure. It is possible that it is due to free radical-scavenging activities of polyphenolic compounds

(Asfar *et al.*, 2003; Hanczakowska & Swiatkiewicz, 2012).

Perhaps an increased villus height is paralleled by an increased expression of brush border enzymes and improved nutrient transport system (Hanczakowska & Swiatkiewicz, 2012). Demir *et al.* (2003) showed that crypt death in the ileum was significantly reduced by including garlic and thyme in the diet compared with including antibiotics and oregano.

4. Conclusion

Summing up the results obtained, it can be stated that thyme volatile oils can have positive changes in small intestine morphology in rats due to NSAID-induced small intestinal damage.

5. References

- Abimosleh, S.M., Tran, C.D. and Howarth, G.S. 2013. Emu oil reduces small intestinal inflammation in the absence of clinical improvement in a rat model of indomethacin-induced enteropathy. *Evidence-Based Complementary and Alternative Medicine*, Article ID 429706, 10 pages.
- Aptekmann, K.P., Baraldi, A.S.M., Stefanini, M.A. and Orsi, M.A. 2001. Morphometric analysis of the intestine of domestic quails (*Coturnix coturnix japonica*) treated with different levels of dietary calcium. *Anatomia, Histologia, Embryologia.*, 30: 277-280.
- Asfar, A., Abdeen, S., Dashti, H., Khoursheed, M., Al-Sayer, H., Mathew, T. and Al-Bader, A. 2003. Effect of green tea in the prevention and reversal of fasting-induced intestinal mucosal damage. *Nutrition.*, 19: 536-540.
- Bancroft, J.D. and Gamble, M.N. 2002. Theory and Practice of Histological Techniques. 5th Ed, Churchill Livingstone, London.
- Boudry, G. and Perrier, C. 2008. Thyme and cinnamon extracts induce anion secretion in piglet small intestine via cholinergic pathways. *Journal of Physiology and Pharmacology.*, 59(3): 543-552.
- Braga, P.C., Dalsasso, M., Culici, M., Bianchi, T., Bordoni, L. and Marabini, L. 2006. Anti-inflammatory activity of thymol: inhibitory effect of the release of human neutrophil elastase. *Pharmacology.*, 77:130-136.

- Demir, E., Sarica, Ş., Özcan, M.A. and Suiçmez, M. 2003. The use of natural feed additives as alternative for an antibiotic growth promoter in broiler diets. *British Poultry Science.*, 45(Supp. 1):44 (Abstract.).
- Duncan, D. B. 1955. Multiple range test and multiple *F* tests. *Biometrics.*, 11:1-42.
- Fang, J., Yan, F.Y., Kong, X.F., Ruan, Z., Liu, Z.Q., Huang, R.L., Geng, M.M., Yang, F., Zhang, Y.Z., Li, P., Gong, J., Wu, G.Y., Fan, M.Z., Liu, Y.L., Hou, Y.Q. and Yin, Y.L. 2009. Dietary supplementation with *Acanthopanax senticosus* extract enhances gut health in weaning piglets. *Livestock Science.*, 123: 268-275.
- Fang, W.F., Broughton, A. and Jacobson, E.D. 1997. Indomethacin induced intestinal inflammation. *Digestive Diseases.*, 22: 749-60.
- Garcia, V., Catala-Gregori, V., Hernandez, F., Megias, M.D. and Madrid, J. 2007. Effect of formic acid and plant extracts on growth, nutrient digestibility, intestine mucosa morphology and meat yield of broilers. *The Journal of Applied Poultry Research.*, 16:555-562.
- Guo, F., Kwakkel, C.R.P., Soede, J. and Williams, B.A. 2004. Effect of a Chinese herb medicine formulation as an alternative for antibiotics on performance of broiler. *British Poultry Science.*, 45: 793-797.
- Hanczakowska, E. and Swiatkiewicz, M. 2012. Effect of herbal extracts on piglet performance and small intestinal epithelial villi. *Czech Journal of Animal Science.*, (9): 420-429.
- Jamroz, D., Wartecki, T., Houszka, M. and Kamel, C. 2006. Influence of diet type on the inclusion of plant origin active substances on morphological and histochemical characteristics of the stomach and jejunum walls in chicken. *Journal of Animal Physiology and Animal Nutrition.*, 90: 255-268
- Khaksar, V., Golian, A. and Raji, A. 2013. Effect of feed additives on intestinal histomorphology of broilers fed wheat-based diet. *Iranian Journal of Applied Animal Science.*, 3: 725-731.
- Kruk, I., Michalska, T., Lichszeld, K., Kladna, A. and Aboul-Enein, H.Y. 2000. The effect of thymol and its derivate on reactions generating reactive oxygen species. *Chemosphere.*, 41:1054-1064.
- Menozzi, A., Sozzoli, C., Poli, E., Martelli, M., Martelli, L., Zullian, C. and Bertini, S. 2009. Effects of oral curcumin on indomethacin-induced small intestinal damage in the rat. *Drug Discoveries and Therapeutics.*, 3(2): 71-76.
- Montagne, L., Pluske, J.R. and Hampson, D.J. 2003. A review of interactions between dietary fibre and the intestinal mucosa, and their consequences on digestive health in young non-ruminant animals. *Animal Feed Science and Technology.*, 108: 95-117.
- Pappenheimer, J.R. and Michel, C.C. 2003. Role of villus microcirculation in intestinal absorption of glucose: coupling of epithelial with endothelial transport. *Journal of Physiology.*, 2: 561-574.
- Sakamoto, K., Hirose, H., Onizuka, A., Hayashi, M., Futamura, N., Kawamura, Y. and Ezaki, T. 2000. Quantitative study of changes in intestinal morphology and mucus gel on total par-enteral nutrition in rats. *Journal of Surgical Research.*, 94: 99-106.
- SAS. 2002. SAS User's Guide: Statistics. Software Release 9.1. SAS Inst. Inc., Cary, NC.
- Tipu, M.A., Akhtar, M.S., Anjum, M.I. and Raja, M.L. 2006. New dimension of medicinal plants as animal feed. *Pakistan Veterinary Journal*, 26:144-148.
- Velazques, G., Borbolla, A.G., Mariscal-Landin, G., Reis de Souza, T. and Pinelli, A. 2005. Intestinal morphology of weaned pigs fed diets containing herbal extracts as growth promoters. *Journal of Animal Science.*, 83 (Suppl. 1), M87.
- Xu, Z.R., Hu, C.H., Xia, M.S., Zhan, X.A. and Wang, M.Q. 2003. Effects of dietary fructooligosaccharide on digestive enzyme activities, intestinal microflora and morphology of male broilers. *Poultry Science.*, 82: 1030-1036.
- Yamada, S., Naito, Y., Takagi, T., Mizushima, K., Horie, R., Fukumoto, K., Inoue, K., Harusato, A., Uchiyama, K., Handa, O., Yagi, N., Ichikawa, H. and Yoshikawa, T. 2012. Rebamipide ameliorates indomethacin-induced small intestinal injury in rats via the inhibition of matrix metalloproteinases activity. *Journal of Gastroenterology and Hepatology.*, 27(12): 1816-1824.