

Oral Mucositis Prevention and Management by Therapeutic Laser in Head and Neck Cancers

Reza Fekrazad¹, Nasim Chiniforush²

¹Laser Research Center in Medical Sciences (LRCMS), AJA University of Medical Sciences- Laser Research Center of Dentistry (LRCD), School of Dentistry, Tehran University of Medical Sciences, Tehran, Iran

²Laser Research Center of Dentistry (LRCD), School of Dentistry, Tehran University of Medical Sciences, Tehran, Iran

Abstract:

Introduction: Oral mucositis is considered a severe complication in cancer patients receiving radiotherapy or chemotherapy for head and neck cancer. The aim of this review study was to assess the effect of low level laser therapy for prevention and management of oral mucositis in cancer patients.

Methods: The electronic databases searched included Pubmed, ISI Web of Knowledge and Google scholar with keywords as “oral mucositis”, “low level laser therapy” from 2000 to 2013.

Results: The results of most studies showed that photobiomodulation (PBM) reduced the severity of mucositis. Also, it can delay the appearance of severe mucositis.

Conclusion: Low level laser therapy is a safe approach for management and prevention of oral mucositis.

Keywords: low level laser therapy; mucositis; prevention; management.

Please cite this article as follows:

Fekrazad R, Chiniforush N. Oral Mucositis Prevention and Management by Therapeutic Laser. *J Lasers Med Sci* 2014;5(1):1-7

Corresponding Author: Nasim Chiniforush, DDS, PhD Candidate; Laser Research Center of Dentistry (LRCD), School of Dentistry, Tehran University of Medical Sciences, Tehran, Iran. Tel: +98-2188015017; Fax: +98-2188687471; E-mail: n-chiniforush@farabi.tums.ac.ir

Introduction

Oral mucositis is considered as a painful condition in patients with cancer after radiotherapy and chemotherapy. It appears as small patches to big ulceration and hemorrhage in oral mucosa¹.

The prevalence of mucositis is 40-100% dependent on the type cancer and therapy used. In 40% of chemotherapy patients and in 100 % of radiotherapy treated patients, mucositis happened^{2,3}.

Oral mucositis not only damages the epithelial cells but also the local tissue reaction causes damage from reactive oxygen species and inflammatory cytokines which finally results in submucosal tissue damage. Oral mucositis appears in oral mucosa after the 5th to 10th day after chemotherapy or radiotherapy. It manifests as a burning sensation to ulcer formation that affect the quality of life of patients by producing pain and discomfort for

swallowing which ultimately leads to malnutrition and dehydration⁴.

Treatment modalities can be non-pharmacological and by pharmacological agents like lidocaine, morphine which is used as palliative care and pain relief. Non pharmacological treatments include mouthrinse, cryotherapy, etc⁵. Recently, photobiomodulation has been used in different aspects of dentistry. photobiomodulation (PBM) is a safe, non pharmacological method which can modulate various metabolic processes via absorption of energy by chromophores⁶.

Low level laser therapy changes the mitochondrial respiration and leads to increased ATP production which produces intracellular reactive oxygen species (ROS). These changes result in fibroblast proliferation, collagen synthesis, adjustment of inflammatory response, as well as motivation of angiogenesis and tissue repair⁷.

There are several explanations for pain reduction

caused by PBM including modulation of the inflammatory process, alteration of excitation and nerve conduction in peripheral nerves and stimulation of release of endogenous endorphins⁸. So, laser therapy can be considered as a palliative treatment for this problem.

The aim of this review study was to assess the effect of low level laser therapy for prevention and management of oral mucositis in cancer patients.

Methods

The electronic databases searched included Pubmed, ISI Web of Knowledge and Google scholar with keywords as “oral mucositis”, “low level laser therapy” from 2000

to 2013.

The papers which are relevant were included in this review of literature.

74 articles were found, but 27 of them were selected for evaluation due to accessibility to their full texts. Also, systematic reviews and meta analysis were excluded.

Results

Among the 27 articles assessed, three of them were animal studies and 24 were human studies. The studies are summarized in Table 1 and Table 2. The results of most studies showed that PBM reduced the severity of mucositis. Also, it can delay the appearance of severe

Table 1. Human studies which evaluated the effect of low level laser therapy on management of oral mucositis.

First author/ year	Title	participants	wavelength	Parameters	Results	Final
Antunes HS 2013 ⁹	Phase III trial of low-level laser therapy to prevent oral mucositis in head and neck cancer patients treated with concurrent chemoradiation	94 HNSCC patients	InGaAlP 660nm	P:100mW E:1J Energy density: 4J/cm ²	Preventive Low-level laser therapy (LLLT) in HNSCC patients receiving chemoradiotherapy is an effective tool for reducing the incidence of grade 3-4 oral mucositis (OM).	+
Arbabi-Kalati F 2013 ¹⁰	Evaluation of the effect of low level laser on prevention of chemotherapy-induced mucositis	24 HNC patients	Diode 630 nm	output power: 30 mW. energy dose : 5 J/cm ²	means of pain severity and mucositis grades were significantly lower in the case group compared to the control group	+
Gautam AP 2013 ¹¹	Effect of low-level laser therapy on patient reported measures of oral mucositis and quality of life in head and neck cancer patients receiving chemoradiotherapy--a randomized controlled trial.	220 HNC patients	He-Ne, 632.8 nm	power density = 24 mW/cm(2), dosage = 3.0 J at each point, total dose/ session = 36-40 J, spot size 1 cm(2), irradiation time/point 125 s)	LLLT was effective in improving the patient's subjective experience of OM and quality of life (QOL) in HNC patients receiving chemoradiotherapy.	+
Gautam AP 2012 ¹¹	Low level laser therapy for concurrent chemoradiotherapy induced oral mucositis in head and neck cancer patients - a triple blinded randomized controlled trial	221 HNC patients	HeNe, 632.8 nm	Power density=24 mW, dosage=3.0 J/ point, total dosage/ session=36-40 J, spot-size=1cm(2), 5 sessions/week)	LLLT decreased the incidence of chemoradiotherapy induced severe OM and its associated pain, dysphagia and opioid analgesics use.	+
Hodgson BD 2012 ¹²	Amelioration of oral mucositis pain by NASA near-infrared light-emitting diodes in bone marrow transplant patients.	80 HSCT patients	GaAlAs 670 nm	for 80 s at ~50 mW/cm(2) energy density : 4 J/cm(2).	Phototherapy demonstrated a significant reduction in patient-reported pain as measured by the WHO criteria in this patient population included in this study.	+
Gouvêa de Lima A 2012 ¹³	Oral mucositis prevention by low-level laser therapy in head-and-neck cancer patients undergoing concurrent chemoradiotherapy: a phase III randomized study.	75 HNSCC patients	GaAlAs	Energy density:2.5 J/cm(2)	LLL therapy was not effective in reducing severe oral mucositis, although a marginal benefit could not be excluded.	=
Carvalho PA, 2011 ¹⁴	Evaluation of low-level laser therapy in the prevention and treatment of radiation-induced mucositis: a double-blind randomized study in head and neck cancer patients.	70 patients with malignant neoplasms in the oral cavity or oropharynx	660nm	Group 1: (15mW/3.8J/cm(2)/spot size 4mm(2)) Group 2: (660nm/5mW/1.3J/cm(2)/spot size 4mm(2))	Low-level laser therapy during radiotherapy was found to be effective in controlling the intensity of mucositis and pain.	+

Treatment of Oral Mucositis by Laser

Table 1. Continue

Cauwels RG 2011 ¹⁵	Low level laser therapy in oral mucositis: a pilot study.	16 children	GaAlAs 830 nm	Power: 150 mW.	Immediate pain relief and improved wound healing resolved functional impairment that was obtained in all cases.	+
Silva GB 2011 ¹⁶	The prevention of induced oral mucositis with low-level laser therapy in bone marrow transplantation patients: a randomized clinical trial.	42 HSCT patients	InGaAlP 660 nm	Power:40 mW, Energy density: 4 J/cm	Our results indicate that the preventive use of LLLT in patients who have undergone HSCT is a powerful instrument in reducing OM incidence.	+
Lima AG 2010 ¹⁷	Efficacy of low-level laser therapy and aluminum hydroxide in patients with chemotherapy and radiotherapy-induced oral mucositis.	25 HNC patients	830 nm	Twelve patients received LLLT (15 mW, 12 J/cm ²) daily from the 1st day until the end of RT before each sessions during 5 consecutive days	The prophylactic use of both treatments proposed in this study seems to reduce the incidence of severe OM lesions. However, the LLLT was more effective in delaying the appearance of severe OM.	+
Khourri VY 2009 ¹⁸	Use of therapeutic laser for prevention and treatment of oral mucositis.	22 HSCT patients	InGaAlP 660 nm GaAlAs 780 nm	Power: 25 mW dose: 6.3J/cm ² 10 s	Laser reduced the frequency and severity of oral mucositis.	+
Zanin T, 2010 ¹⁹	Use of 660-nm diode laser in the prevention and treatment of human oral mucositis induced by radiotherapy and chemotherapy.	72 patients HNC patients	diode 660 nm	Power : 30 mW, spot size = 2 mm, energy = 2 J per point	Laser therapy was effective in preventing and treating oral effects induced by radiotherapy and chemotherapy, thus improving the patient's quality of life.	+
Simões A, 2009 ²⁰	Laser phototherapy as topical prophylaxis against head and neck cancer radiotherapy-induced oral mucositis: comparison between low and high/low power lasers.	39 HNC patients	InGaAlP laser (660 nm) GaAlAs laser (808 nm)	LPT: 40 mW/6Jcm ⁻² /0.24 J per point). high power LPT : 1 W/cm ² .	Low-intensity laser therapy appears to reduce the severity of mucositis, at least in part, by reducing cyclooxygenase-2 levels and associated inhibition of the inflammatory response.	+
Kuhn A 2009 ²¹	Low-level infrared laser therapy in chemotherapy-induced oral mucositis: a randomized placebo-controlled trial in children.	21 patients (86%) patients had a diagnosis of leukemia or lymphoma and 3(14%) had solid tumors	GaAlAs 830 nm	Power: 100 mW, dose: 4 J/cm ² ,	laser therapy in addition to oral care can decrease the duration of chemotherapy-induced OM.	+
Campos L, 2009 ²²	Improvement in quality of life of an oncological patient by laser phototherapy.	a 15-year-old girl diagnosed	Diode 660 nm	Power: 40 mW, energy density: 6 J/cm ² , 0.24 J per point, spot size of 0.04 cm ² A high-power diode laser at 1 W,10 J/cm ² defocused mode	Normal oral function and consequent improvements in the quality of life of this oncologic patient were observed with laser phototherapy (LPT).	+
Eduardo FP 2009 ²³	Severity of oral mucositis in patients undergoing hematopoietic cell transplantation and an oral laser phototherapy protocol: a survey of 30 patients.	30 patients undergoing HCT	Diode 660 nm	Power: 40 mW	The low grades of OM observed in this survey show the beneficial effects of laser phototherapy.	+
Abramoff MM 2008 ²⁴	Low-level laser therapy in the prevention and treatment of chemotherapy-induced oral mucositis in young patients.	13 Patients	GaAlAs 685 nm	Power:35 mW 600-m spot. energy: 2 J per point, dose: 72 J/cm ² .	The ease of use of LLLT, high patient acceptance, and the positive results achieved, make this therapy feasible for the prevention and treatment of OM in young patients.	+
Genot-Klastersky MT 2008 ²⁵	The use of low-energy laser (LEL) for the prevention of chemotherapy- and/or radiotherapy-induced oral mucositis in cancer patients: results from two prospective studies.	First study:26 patients Second study: 36 patients	Visible IR	visible 100 mW laser and an IR laser with power from 50, 250, and 500 mW	LEL is an effective and safe approach to prevent or treat oral mucositis resulting from cancer chemotherapy.	+

Treatment of Oral Mucositis by Laser

Table 1. Continue

Antunes HS 2008 ²⁶	The Impact of low power laser in the treatment of conditioning-induced oral mucositis: a report of 11 clinical cases and their review.	11 HSCT patients	InGaAlP, 660 nm	Power: 50 mW dose: 8 J/cm ²	The use of low power laser therapy in HSCT patients is a powerful instrument in the treatment of overt oral mucositis.	+
Arora H 2008 ²⁷	Efficacy of He-Ne Laser in the prevention and treatment of radiotherapy-induced oral mucositis in oral cancer patients.	24 HNC patients	He-Ne 632.8nm	Power: 10 mW energy density: 1.8 J/cm ² .	Laser therapy applied prophylactically during radiotherapy can reduce the severity of oral mucositis, severity of pain, and functional impairment.	+
Jaguar GC 2007 ²⁹	Low-energy laser therapy for prevention of oral mucositis in hematopoietic stem cell transplantation.	24 HSCT patients	GaAlAs 660 nm	power :10 mW energy density: 2,5 J/cm ²	This study suggests that laser therapy can be useful in oral mucositis to HSCT patients and improve the patient's quality of life.	+
Schubert MM 2007 ⁵	A phase III randomized double-blind placebo-controlled clinical trial to determine the efficacy of low level laser therapy for the prevention of oral mucositis in patients undergoing hematopoietic cell transplantation.	70 HCT patients	650 nm 780 nm	energy density: 2 J/cm ² .	LLLT reduced the severity of oral mucositis and pain scores	+
Cruz LB 2007 ²⁹	Influence of low-energy laser in the prevention of oral mucositis in children with cancer receiving chemotherapy.	60 HSCT patients	780 nm wavelength	Power: 60mW, Dose: 4 J/cm ² .	This study showed no evidence of benefit from the prophylactic use of low-energy laser in children and adolescents with cancer treated with chemotherapy when optimal dental and oral care was provided.	=
Arun Maiya G 2006 ³⁰	Effect of low level helium-neon (He-Ne) laser therapy in the prevention & treatment of radiation induced mucositis in head & neck cancer patients	20 patients with carcinoma of oral cavity with stages II-IV	He-Ne 632.8 nm	Power: 10mW	The low-level He-Ne laser therapy during the radiotherapy treatment was found to be effective in preventing and treating the mucositis in head and neck cancer patients.	+
Nes AG 2005 ³¹	Patients with moderate chemotherapy-induced mucositis: pain therapy using low intensity lasers.	13 adult patients	AsGaAl 830 nm	Power: 250 mW. Dose: 35 J cm ⁻² .	There was a significant decrease in the daily average experience of pain felt before and after each treatment.	+

Table 2. Animal studies which evaluated the effect of low level laser therapy on management of oral mucositis.

Author/year	Title	participants	wavelength	Parameters	Results	Final
Lopes NN 2010 ³²	Effects of low-level laser therapy on collagen expression and neutrophil infiltrate in 5-fluorouracil-induced oral mucositis in hamsters.	72 hamster cheek	InGaAlP 660 nm	power of 35 or 100 mW	LLLT promotes wound healing and appears to have an anti-inflammatory effect, as evidenced by the reduction in neutrophil infiltrate.	+
Lopes NN 2009 ³³	Cyclooxygenase-2 and vascular endothelial growth factor expression in 5-fluorouracil-induced oral mucositis in hamsters: evaluation of two low-intensity laser protocols.	72 hamster cheek	InGaAlP 660 nm	Power: 35 mW or 100 mW laser	The increase in collagen organization in response to the 35 mW laser indicates that LLLT promotes wound healing. Also, LLLT also appears to have an anti-inflammatory effect.	+
França CM 2009 ⁴	Low-intensity red laser on the prevention and treatment of induced-oral mucositis in hamsters.	30 hamsters	Diode 660 nm	Power= 30mW, Dose=1.2J/cm ² , 40s, spot size 3mm ² ,	The results suggest that laser phototherapy had a positive effect in reducing mucositis severity, and a more pronounced effect in treating established mucositis.	+

HNSCC: head and neck squamous cell carcinoma

HNC: head and neck cancer

HSCT: hematopoietic stem cell transplant

GaAlAs: gallium-aluminum-arsenide

InGaAlP: Indium Gallium Aluminum Phosphorus

+: effective response

=: no effective response

mucositis.

Discussion

Oral mucositis is considered as a side effect of radiotherapy or chemotherapy of head and neck cancer patients.

It's necessary to standardize some protocols for prevention and management of chemotherapy or radiotherapy induced oral mucositis to prevent reduction or discontinuation of treatments²⁰. Recently, researches have focused on the effects of low level laser therapy as a new modality for managing this problem via photobiomodulation effect^{9, 10}. Oral mucositis exhibit different biologic phases including initiation, inflammatory response, amplification/ signaling, ulceration and healing. Acceleration of wound healing happens through enhanced release of growth factors, increased neovascularization and collagen formation. Some studies stated that laser can increase the production of vascular endothelial growth factor (VEGF) which results in angiogenesis and improved microcirculation and subsequently to wound healing^{33, 34}.

Mostly, low level laser therapy was performed by diode lasers including red and infra-red wavelengths. Shorter wavelengths (632.8-660 nm) can affect the superficial layer of the epithelium but longer wavelengths penetrate deeper and affect sub-epithelial tissues. It seems that shorter wavelengths are more effective than longer ones in treatment of oral mucositis which was approved by Schubert et al. Red laser are absorbed in superficial layer of oral mucosa which is favorable for wound healing⁵.

The effect of laser on oral mucositis showed some beneficial effects such as it delayed the time of onset, reduced the peak severity and shortened the duration of this problem. These advantages can be attributed to anti-inflammatory and analgesic effects of laser which are followed by increased local vascularity and re-epithelization of tissue³⁰. Oral wound healing can be achieved by transformation of fibroblast to myofibroblasts which promote mucosal healing. On the other hand, analgesic effects can happen through release of endorphin and eukephalin. The other mechanism of laser induced pain relief is through membrane cell depolarization, blocking the nervous impulse and fast axonal flow³⁵.

Most of studies showed the positive effect of PBM on oral mucositis but there were just two studies that reported negative effect of laser therapy. Curz et al. in assessing low level laser therapy in prevention of oral mucositis in children came to this conclusion that low level laser therapy had no beneficial effect as prophylactic option for prevention of oral mucositis²⁹.

Bjordal et al. conducted meta-analysis of the effect of

Low-level laser therapy (LLLT) in cancer therapy-induced oral mucositis. Despite, various laser wavelengths, parameters and dosage, they demonstrated moderate to strong evidence for efficacy of LLLT in oral mucositis management³⁵. Also, Migliorati et al. stated that LLLT can become a routine practice in management of oral mucositis to control its severity³⁶.

In assessing animal studies, anti-inflammatory effect of LLLT and wound healing was seen due to reduction in neutrophil infiltrate and cyclooxygenase-2 expression^{32, 33}.

Other factors such as oral hygiene had influenced on prevention of oral mucositis. So, patients should be educated about oral hygiene and given instructions. Some researchers showed positive effects of good oral health on reducing the risk of severe mucositis due to decrease of oral bacteria²⁹.

One of the main factors that can enhance the prevention of oral mucositis is the ideal time of starting laser therapy which has not been clear and needs clinical trials which would evaluate and compare the different startpoints of treatment to clarify it³⁵.

According to two meta-analysis conducted about laser phototherapy in management of oral mucositis in head and neck cancers, LLLT can be used with doses of 1-6 J per point in oropharyngeal area by existence of moderate to strong evidence. For radiotherapy induced oral mucositis, a suggestion of laser therapy was possible with wavelength of 632 nm. No guideline was possible for application of LLL in management of oral mucositis in patients treated with a combined protocol of radiochemotherapy for head and neck cancer. Also, there was no possible guideline for chemotherapy induced oral mucositis due to low level of evidence^{35, 36}.

Several studies showed positive results of LLLT for reducing the incidence and severity of mucositis accompanied by other advantages such as improved quality of life of patients. More standard investigations are needed to clarify which parameters including wavelength, energy density, time of exposure, have the best interactions with tissues to produce favorable outcomes.

Conclusion

Low level laser therapy is a noninvasive modality for prevention and management of oral mucositis with to some advantages like analgesic effect, inflammation reduction and atraumatic repeated use.

References

1. Avritscher E, Cooksley L. Scope and epidemiology of cancer therapy-induced oral and gastrointestinal mucositis. *Semin Oncol Nurs* 2004;20:3-8.
2. Trotti A, Bellm LA, Epstein JB, Frame D, Fuchs HJ, Gwede CK, et al. Mucositis incidence, severity and associated outcomes in patients with head and neck cancer receiving radiotherapy with or without chemotherapy: a systematic literature review. *Radiother Oncol* 2003;66(3):253-62.
3. Bjordal JM, Bensadoun RJ, Tuner J, Frigo L, Gjerde K, Lopes-Martins RA. A systematic review with meta-analysis of the effect of low-level laser therapy (LLLT) in cancer therapy-induced oral mucositis. *Support Care Cancer* 2011;19(8):1069-77.
4. França CM, França CM, Núñez SC, Prates RA, Noborikawa E, Faria MR, et al. Low-intensity red laser on the prevention and treatment of induced-oral mucositis in hamsters. *J Photochem Photobiol B* 2009 9;94(1):25-31.
5. Schubert MM, Eduardo FP, Guthrie KA, Franquin JC, Bensadoun RJ, Migliorati CA, et al. A phase III randomized double-blind placebo-controlled clinical trial to determine the efficacy of low level laser therapy for the prevention of oral mucositis in patients undergoing hematopoietic cell transplantation. *Support Care Cancer* 2007;15(10):1145-54.
6. Fekrazad R, Chiniforush N, Bouraima SA, Valipour M, Aslani M, Zare M, et al. Low Level Laser Therapy in Management of Complications after Intra Oral Surgeries. *J Lasers Med Sci* 2012;3(4):135-40.
7. Mohkovska T, Mayberry J. It is time to test low level laser therapy in Great Britain. *Postgrad Med J* 2005;81(957):436-41.
8. Chow RT, Johnson MI, Lopes-Martins RA, Bjordal JM. Efficacy of low-level laser therapy in the management of neck pain: a systematic review and meta-analysis of randomised placebo or active-treatment controlled trials. *Lancet* 2009 5;374(9705):1897-908.
9. Antunes HS, Herchenhorn D, Small IA, Araújo CM, Viégas CM, Cabral E, et al. Phase III trial of low-level laser therapy to prevent oral mucositis in head and neck cancer patients treated with concurrent chemoradiation. *Radiother Oncol* 2013.
10. Arbabi-Kalati F, Arbabi-Kalati F, Moridi T. Evaluation of the effect of low level laser on prevention of chemotherapy-induced mucositis. *Acta Med Iran* 2013 ;51(3):157-62.
11. Gautam AP, Fernandes DJ, Vidyasagar MS, Maiya AG, Nigudgi S. Effect of low-level laser therapy on patient reported measures of oral mucositis and quality of life in head and neck cancer patients receiving chemoradiotherapy-a randomized controlled trial. *Support Care Cancer* 2013;21(5):1421-8.
12. Hodgson BD, Margolis DM, Salzman DE, Eastwood D, Tarima S, Williams LD, et al. Amelioration of oral mucositis pain by NASA near-infrared light-emitting diodes in bone marrow transplant patients. *Support Care Cancer* 2012;20(7):1405-15.
13. Gouvêa de Lima A, Villar RC, de Castro G Jr, Antequera R, Gil E, Rosalmeida MC, et al. Oral mucositis prevention by low-level laser therapy in head-and-neck cancer patients undergoing concurrent chemoradiotherapy: a phase III randomized study. *Int J Radiat Oncol Biol Phys* 2012;82(1):270-5.
14. Carvalho PA, Jaguar GC, Pellizzon AC, Prado JD, Lopes RN, Alves FA. Evaluation of low-level laser therapy in the prevention and treatment of radiation-induced mucositis: a double-blind randomized study in head and neck cancer patients. *Oral Oncol* 2011;47(12):1176-81.
15. Cauwels RG, Martens LC. Low level laser therapy in oral mucositis: a pilot study. *Eur Arch Paediatr Dent* 2011;12(2):118-23.
16. Silva GB, Mendonça EF, Bariani C, Antunes HS, Silva MA. The prevention of induced oral mucositis with low-level laser therapy in bone marrow transplantation patients: a randomized clinical trial. *Photomed Laser Surg* 2011;29(1):27-31.
17. Lima AG, Antequera R, Peres MP, Snitcosky IM, Federico MH, Villar RC. Efficacy of low-level laser therapy and aluminum hydroxide in patients with chemotherapy and radiotherapy-induced oral mucositis. *Braz Dent J* 2010;21(3):186-92.
18. Khouri VY, Stracieri AB, Rodrigues MC, Moraes DA, Pieroni F, Simões BP, et al. Use of therapeutic laser for prevention and treatment of oral mucositis. *Braz Dent J* 2009;20(3):215-20.
19. Zanin T, Zanin F, Carvalhosa AA, Castro PH, Pacheco MT, Zanin IC, et al. Use of 660-nm diode laser in the prevention and treatment of human oral mucositis induced by radiotherapy and chemotherapy. *Photomed Laser Surg* 2010;28(2):233-7.
20. Simões A, Eduardo FP, Luiz AC, Campos L, Sá PH, Cristóforo M, et al. Laser phototherapy as topical prophylaxis against head and neck cancer radiotherapy-induced oral mucositis: comparison between low and high/low power lasers. *Lasers Surg Med* 2009;41(4):264-70.
21. Kuhn A, Porto FA, Miraglia P, Brunetto AL. Low-level infrared laser therapy in chemotherapy-induced oral mucositis: a randomized placebo-controlled trial in children. *J Pediatr Hematol Oncol* 2009;31(1):33-7.
22. Campos L, Simões A, Sá PH, Eduardo Cde P. Improvement in quality of life of an oncological patient by laser phototherapy. *Photomed Laser Surg* 2009;27(2):371-4.
23. Eduardo FP, Bezinelli L, Luiz AC, Correa L, Vogel C, Eduardo CP. Severity of oral mucositis in patients undergoing hematopoietic cell transplantation and an oral laser phototherapy protocol: a survey of 30 patients. *Photomed Laser Surg* 2009;27(1):137-44.
24. Abramoff MM, Lopes NN, Lopes LA, Dib LL, Guilherme A, Caran EM, et al. Low-level laser therapy in the prevention and treatment of chemotherapy-induced oral mucositis in young patients. *Photomed Laser Surg* 2008;26(4):393-400.
25. Genot-Klastersky MT, Klastersky J, Awada F, Awada A, Crombez P, Martinez MD, et al. The use of low-energy laser (LEL) for the prevention of chemotherapy- and/or radiotherapy-induced oral mucositis in cancer patients: results from two prospective studies. *Support Care Cancer*

- 2008;16(12):1381-7.
26. Antunes HS, Ferreira EM, de Matos VD, Pinheiro CT, Ferreira CG. The Impact of low power laser in the treatment of conditioning-induced oral mucositis: a report of 11 clinical cases and their review. *Med Oral Patol Oral Cir Bucal* 2008;13(3):E189-92.
 27. Arora H, Pai KM, Maiya A, Vidyasagar MS, Rajeev A. Efficacy of He-Ne Laser in the prevention and treatment of radiotherapy-induced oral mucositis in oral cancer patients. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2008;105(2):180-6.
 28. Jaguar GC, Prado JD, Nishimoto IN, Pinheiro MC, de Castro DO Jr, da Cruz Perez DE, et al. Low-energy laser therapy for prevention of oral mucositis in hematopoietic stem cell transplantation. *Oral Dis* 2007;13(6):538-43.
 29. Cruz LB, Ribeiro AS, Rech A, Rosa LG, Castro CG Jr, Brunetto AL. Influence of low-energy laser in the prevention of oral mucositis in children with cancer receiving chemotherapy. *Pediatr Blood Cancer* 2007;48(4):435-40.
 30. Arun Maiya G, Sagar MS, Fernandes D. Effect of low level helium-neon (He-Ne) laser therapy in the prevention & treatment of radiation induced mucositis in head & neck cancer patients. *Indian J Med Res* 2006;124(4):399-402.
 31. Nes AG, Posso MB. Patients with moderate chemotherapy-induced mucositis: pain therapy using low intensity lasers. *Int Nurs Rev* 2005;52(1):68-72.
 32. Lopes NN, Plapler H, Lalla RV, Chavantes MC, Yoshimura EM, da Silva MA, et al. Effects of low-level laser therapy on collagen expression and neutrophil infiltrate in 5-fluorouracil-induced oral mucositis in hamsters. *Lasers Surg Med* 2010;42(6):546-52.
 33. Lopes NN, Plapler H, Chavantes MC, Lalla RV, Yoshimura EM, Alves MT. Cyclooxygenase-2 and vascular endothelial growth factor expression in 5-fluorouracil-induced oral mucositis in hamsters: evaluation of two low-intensity laser protocols. *Support Care Cancer* 2009;17(11):1409-15.
 34. Zhang WG, Wu CY, Pan WX, Tian L, Xia JL. Low-power Helium-Neon laser irradiation enhances the expression of VEGF in murine myocardium. *Chin Med J (Engl)* 2004;117(10):1476-80.
 35. Bjordal JM, Bensadoun RJ, Tuner J, Frigo L, Gjerde K, Lopes-Martins RA. A systematic review with meta-analysis of the effect of low-level laser therapy (LLLT) in cancer therapy-induced oral mucositis. *Support Care Cancer* 2011;19(8):1069-77.
 36. Migliorati C, Hewson I, Lalla RV, Antunes HS, Estilo CL, Hodgson B, et al. Systematic review of laser and other light therapy for the management of oral mucositis in cancer patients. *Support Care Cancer* 2013;21(1):333-41.

Archive of SID