

## Evaluation of the effects of hydroalcoholic extract of *Berberis vulgaris* root on the activity of liver enzymes in male hypercholesterolemic rats

Soheila Taheri<sup>1</sup>, Ali Zarei<sup>2</sup>, Saeed Changizi Ashtiyani<sup>1\*</sup>, Azam Rezaei<sup>3</sup>, Saeed Zaheiri<sup>4</sup>

### Abstract

**Objectives:** Hyperlipidemia can cause a variety of diseases such as atherosclerosis, diabetes, and fatty liver which is followed by increased liver enzymes. Since *Berberis vulgaris* (*B. vulgaris*) root possesses antioxidant properties, the present study was conducted to investigate the effect of its extract on the activity of liver enzymes in rats.

**Materials and Methods:** In this experimental study, sixty Wistar rats were selected and allocated to six groups of ten each. The control group received a normal diet and the sham group received a fatty diet while the other groups including experimental groups received a fatty diet and the alcoholic extract of *B. vulgaris* at minimum (75 mg/kg), moderate (150 mg/kg), and maximum (300 mg/kg) doses by intraperitoneal injection (i.p.) or oral atorvastatin (10 mg/kg) with a fatty diet. At the end of this 21-day period, blood samples were drawn and the levels of the intended factors were measured. Data were analyzed using SPSS software version 11.5.

**Results:** The comparison of the obtained results showed that the levels of alanine transaminase (ALT) and alkaline phosphatase (ALP) enzymes in the sham group that only received fatty food increased ( $p < 0.05$ ), whereas in the treatment groups receiving *B. vulgaris* extract as well as in the group receiving Atorvastatin, these enzymes significantly decreased; however, no significant changes were observed in aspartate transaminase (AST) levels.

**Conclusion:** Noticing the antioxidant properties of *B. vulgaris* root extract and its effects on reducing the activity of liver enzymes, the extract of this plant can be a good choice for improving the function of liver.

**Keywords:** Alanine Transaminase, Alkaline Phosphatase, Aspartate Transaminase, Atorvastatin, *Berberis vulgaris*, Liver Enzyme, Rat

1- Department of Physiology, Arak University of Medical Sciences, Arak, I. R. Iran

2- Islamic Azad University of Damghan, Damghan, I. R. Iran

3- Islamic Azad University of Arsanjan, Arsanjan, I. R. Iran

4- Department of Biology, Payame Noor University of Abadeh, Abadeh, I. R. Iran

\*Corresponding Author: Tl/Fax: +988614173526

E-mail: dr.ashtiyani@arakmu.ac.ir

## Introduction

Researchers with interest in natural products have intensified their efforts towards scientific evaluation of traditional medicines. Barberry has played a prominent role in herbal healing for more than 2500 years. *B. vulgaris* (European barberry)/(Jaundice berry)/(Ambarbaris)/(Barberry) is a shrub in the Berberidaceae family, native to central and southern Europe, northwest Africa, and western Asia. Seedless barberry (*B. vulgaris* var. *asperma*) is one of the most important medicinal fruit plants in the south of Khorasan province in Iran. Zereshk or Sereshk is the Persian name for the dried fruit of *B. vulgaris*, which is widely cultivated in Iran (Mohammadi et al., 2011). India's Ayurveda healers have used it for dysentery (Jellin et al., 2000).

During the early middle ages, European herbalists used it to treat liver and gallbladder ailments (Jellin et al., 2000). Russian healers used it for inflammations, high blood pressure, and abnormal uterine bleeding (Jellin et al., 2000). In folk medicine, European barberry root bark has been used for various conditions including liver dysfunction, gallbladder diseases, diarrhea, indigestion, and urinary tract diseases (Jellin et al., 2000).

The stem, root bark, and fruit of barberry contain isoquinoline alkaloids (e.g. berberine), which are the main active ingredients of barberry (Arayne et al., 2007). The bitter compounds in barberry, including alkaloids, stimulate digestive functions following meals (Arayne et al., 2007). Barberry and goldenseal (*Hydrastis canadensis*) have very similar therapeutic uses because both contain similar active substances (berberine alkaloids (Arayne et al., 2007)). These substances have been shown to combat infections and bacteria, stimulate the activity of the immune system, and lower fever (Arayne et al., 2007). Pharmacological studies suggest that these

substances have antimicrobial and antiparasitic, anti-inflammatory, immune-stimulant, fever reducing, hypotensive (causing a reduction in blood pressure), sedative, anticonvulsant, and smooth muscle relaxant effects. Smooth muscles line the gastrointestinal tract; therefore, this last effect may help improve digestion and reduce stomach pain (Arayne et al., 2007).

Mammals' cells possess enzymatic and non-enzymatic capability against the formation of bonds between them and free radicals. Non-enzymatic capability consists of beta-carotene, vitamin C, and vitamin E. The involved enzymes include superoxide dismutase, catalase, and glutathione peroxidase. In case the anti-oxidant defense mechanism is lost in the body, an increase in the formation of free radicals can cause cellular oxidative stress. Lipid peroxidation and oxidant stress increase in diabetes and hyperlipidemia (Kelly and Husband, 2003; Nazari et al., 2005). Recently, the inhibitory effect of *B. vulgaris* root extract on oxidative stress has been reported (Noori et al., 2004). Hyperlipidemia can also play an indirect role by activating the production of free radicals such as monocytes and neutrophils (Pourghassem et al., 2009).

Dietary habits have a major effect on coronary risk factors and hyperlipidemia (Zarei et al., 2011; Heidarian et al., 2008). These diseases have developed across a wide range of developed and underdeveloped societies. Although using lipid lowering anti-lipid chemical drugs such as lovastatin and atorvastatin can reduce hypercholesterolemia and cardiovascular diseases, such drugs have side effects (Zarei et al., 2011; Heidarian et al., 2008). Factors, such as patients' dissatisfaction with chemical drugs, presence of side effects over long-term use and overuse, and financial burdens imposed on the patients, have led to an increased tendency towards alternative and traditional medicine. Using medicinal

plants and fruits, in addition to decreasing health care costs in many societies, has had satisfactory results.

*B. vulgaris* is a medicinal plant that is commonly applied to the treatment of liver and biliary diseases in traditional medicine. *B. vulgaris* has secondary metabolites such as berberin, berlambine, oxyberberine, oxycanthine, chlorumamine, anthocyanine, bervulcine, lambertine, and magniflorine, which are frequently used in the pharmaceutical industry (Arayne et al., 2007; Mohammadi et al., 2011) and plays a major role in the treatment of gastrointestinal diseases, hemorrhages, gum inflammation, sore throat, biliary fevers, malaria, leishmaniasis, hepatitis, inflammation, diarrhea, and high blood cholesterol (Arayne et al., 2007; Mohammadi et al., 2011; Vrzal et al., 2005). Previous studies have shown that the aqueous extract of *B. vulgaris* fruit activates liver function and is effective in dissolving and possibly moderating blood cholesterol levels. Moreover, the extract of this plant reduces blood cholesterol and triglyceride levels (Farhadi, Gavadifar, 2008). In fact, *B. vulgaris* roots possess hypoglycemic properties (Golfaraz, Ahmad, 2008). Based on the studies done on alkaloids in *B. vulgaris*, such as berberin, anthocyanins, and its phenolic compounds, the antioxidant properties of the compounds present in this plant have been proven (Mohammadi et al., 2011; Hwong 2006; Motalleb, 2005).

Therefore, noticing the previous studies on the antioxidant properties of *B. vulgaris* and with regard to the hypoglycemic and triglyceride reducing effects of this plant and the relationship between these factors and liver enzymes, this study was carried out to investigate the effects of *B. vulgaris* root extract on the activity level of liver enzymes in hypercholesterolemic rats and to compare its effect with that of a hypoglycemic drug (atorvastatin) (Farhadi et al., 2008).

## Materials and Methods

This experimental study was done on sixty male rats supplied from the Razi Animal Breeding Center of Fars province. The rats were kept in standard conditions of temperature and light ad libitum. Animal care and handling were performed according to the guidelines set by the Iranian Ministry of Health and Medical Education for lab animals. Before launching the project, all of the rats were weighed to make sure that they are within the same range of weight ( $170 \pm 5$  g). The rats were assorted into six groups of ten each in the following way:

A) Control group: The rats did not receive any vehicle or drugs and had a normal diet. B) Sham group: Hypercholesterolemic rats (2% cholesterol was added to their food to render them hypercholesterolemic) were daily administered 0.2 ml of normal saline as intraperitoneal injection (i.p.) for 21 days. C) Treatment group 1: Hypercholesterolemic rats were administered 75 mg/kg (minimum dose) of the alcoholic extract of *B. vulgaris* (i.p.) for 21 days. D) Treatment group 2: Hypercholesterolemic rats were administered 150 mg/kg (moderate dose) of the alcoholic extract of *B. vulgaris* (i.p.) for 21 days. E) Treatment group 3: Hypercholesterolemic rats were administered 300 mg/kg (maximum dose) of the alcoholic extract of *B. vulgaris* (i.p.) for 21 days. F) Atorvastatin group: In this group, 10 mg/kg of atorvastatin in the form of oral emulsion was administered to hypercholesterolemic rats through gavage for 21 days (Gosain S et al. 2010).

### Preparation of high cholesterol 2% food

For preparing high cholesterol 2% food, 20 g of pure cholesterol powder (Fluka Chemika) was solved in 5 ml of heated olive oil and was mixed with 1 kg of the rats' food. The food was kept in a refrigerator only for two days. (Rafati et al., 2006)

### Extraction method

For preparation of the alcoholic extract of *B. vulgaris*, after obtaining the roots from the Shiraz University Herb Farm, Iran, and removing the impurities, 500 mg of the roots were milled and mixed with ethylic alcohol 98% at the ratio of 1:5. The mixture was kept in a capped container for 4 hours in lab conditions and was filtered by small and big filter papers. The filtered liquid was placed in water bath for solvent evaporation. Eventually, different doses (mg/kg bodyweight) of the extract (10 g per 100 g of the milled roots) were prepared by normal saline. All of the treatment groups received a fatty diet during the experiment. The experiment was carried out over a 21-day period and the injections were done through gavage everyday at 9 a.m. as i.p. injections were made using insulin syringe and atorvastatin (10 mg/kg) (Shafa Pharmaceuticals Co., Iran) was administered as oral emulsion. After this period, mild anesthesia using ether was done for obtaining blood samples from heart to examine the concentration of plasma biochemical factors. After centrifuging blood at 3000 rounds per minute, serum samples were isolated and transferred to the lab for measuring the intended factors. The amount of liver

enzymes contains: alanine transaminase (ALT), alkaline phosphatase (ALP), and aspartate transaminase (AST) were assessed through radio immunoassay (RIA) and Pars Azmoon kit using RIA1000 (USA). All obtained values are expressed as mean±SD and data analyses were done using SPSS ver. 11.5. The group's data means were compared by one-way analysis of variance and Tukey's post hoc test. The level of significance was set at 0.05.

## Results

The comparison between the results of statistical analysis on the effects of *B. vulgaris* root extract and atorvastatin on the activity level of liver enzymes in hypercholesteremic rats (Table 1) indicated that the amount of ALP in the sham group receiving a fatty diet had significantly increased in comparison with the control group, whereas changes in all of the treatment groups receiving the extract and the group receiving atorvastatin presented significant decreases compared with the sham group (p=0.00).

Table 1. The comparison between effect of different doses of *Berberis vulgaris* (*B. vulgaris*) and atorvastatin on the parameters of liver activity in hypercholesterolemic rats.

Group Parameters	Control	Sham	Atorvastatin	<i>B. vulgaris</i> Maximum Dose	<i>B. vulgaris</i> Moderate Dose	<i>B. vulgaris</i> Minimum Dose
Cholesterol	85.00±0.98	97.25±3.13 *	80.62±2.02 †	76.12±5.11 †	86.00±3.95	86.00±5.35
ALT(U/l)	55.87±3.7	80.37±6.7 *	65.28±2.4 †	41.71±6.8 #†	56.57±5.8 †	59.12±4.7
AST(U/l)	234.83±15.4	277.71±38.7	177.43±5.5	244.29±41.5	300.38±34.9	286.83±29.9
ALP(U/l)	1017.30±71.8	2400 ±44.5 *	673.62±12.1 †	914±10.6 †	1023.90±14.0 †	1027.16±10.3 †

† Comparison with the sham group, \* Comparison with the control group, # Comparison with the atorvastatin group, Comparison between the minimum and maximum doses.

## Evaluation of the effects of hydroalcoholic extract of *Berberis vulgaris* root

However, none of the treatment groups compared with one another and compared with the atorvastatin group presented significant changes. In terms of ALT, changes in the sham group were significantly higher than the control group, whereas its amounts in all of the treatment groups receiving *B. vulgaris* root extract as well as the atorvastatin group had significantly decreased. Moreover, the treatment group receiving the maximum dose in comparison with the minimum extract dose and the atorvastatin group showed a significant decrease ( $p=0.001$ ).

This, in fact, indicates that the effect of the maximum dose on the amount of ALP decrease is much stronger than other doses and even atorvastatin. Regarding AST, changes in the sham group are not more significant in comparison with the control group and its amounts in the treatment group does not indicate significant changes ( $p=0.090$ ). There were not shown any significant changes in liver enzymes between moderate dose of *B. vulgaris* extract with minimum and maximum doses (Table 1).

### Discussion

The comparison of the results of statistical analyses showed that ALT and ALP enzymes amounts in the sham group that only received a fatty diet had increased, whereas these enzymes in the treatment groups receiving *B. vulgaris* extract and the atorvastatin group had decreased; however, no significant changes were observed in AST. Non-alcoholic fatty liver disease (NAFLD) is an insulin-resistant pathogenic factor in type II diabetes. Liver enzymes levels in circulation, including ALT, AST, and gamma glutamyl transferase, on the other hand, in asymptomatic with NAFLD are high.

Following reduction in the amount of insulin the metabolism of acids in the liver is disrupted and eventually fatty liver is

resulted which is followed by increases in ALT, AST, and ALP levels (Bush 1991; Jelodar, Nazifi, 1997). Normally, following reduction in serum cholesterol, liver lipidosis decreases and the activity of liver enzymes gets significantly decreased (Jelodar, Nazifi, 1997; Yu, Keeffe, 2003). In this study, fatty liver diseases are defined as the deposition of fat (mainly neutral fats such as triglyceride) in the liver which are usually accompanied by increased liver enzymes (ALT and AST) that may result in liver inflammation. This issue can cause liver hardening and scarring that result in liver cirrhosis. With progression of the disease, scarred and hardened tissues are develop around liver cells and gradually cause nodules and cysts in liver which, in turn, by creating pressure on biliary ducts, lead to obstruction in these ducts, and, eventually, increased bilirubin and cholesterol.

Free radicals destroy cell membranes, such as hepatic cells, which result in increasing liver enzymes. This causes the entrance of enzymes that are normally located inside cell cytosol into blood circulation and increased activity of these enzymes indicates the degree and type of liver damage (Shariati and Zarei, 2006). Therefore, noticing the aforementioned studies and the findings of the present study, increase in the amount of ALT in the sham group that only received fatty food is predictable. *B. vulgaris* has anti-oxidant properties and the berberin in this plant effectively reduces lipid peroxidation with its anti-oxidant properties. This way, it protects antioxidant enzymes, against damage due to oxygen free radicals and peroxide hydrogen (Thirupurasundari et al., 2009; Ziai et al., 2010). Antioxidant activities of the ethanolic extracts of roots, twigs, and leaves of common barberry (*B. vulgaris* L.) were studied. All the extracts were found to possess some radical-scavenging and antioxidant activities (Zovko et al., 2010).

In their analysis of *B. vulgaris* root extract, Tomosaka *et al.* found three phenolic compounds with antioxidant and anti-inflammatory properties (Tomosaka *et al.*, 2008). Therefore, considering the antioxidant properties of *B. vulgaris*, reduction in the activity of liver enzymes in the treatment groups receiving the plant extract is predictable.

Increase in HDL-C lipoprotein results in reduction of cholesterol level and improvement of liver function (Hu *et al.*, 2003). *B. vulgaris* plays a significant role in reducing cholesterol (Arayne *et al.*, 2007; Mohammadi *et al.*, 2011; Vrzal *et al.*, 2005) and its extract decreases cholesterol and triglyceride levels (Farhadi *et al.*, 2008); therefore, it can be applied to the treatment of liver diseases. Noticing this cholesterol-reducing property which, in turn, decreases liver lipodosis that has a significant relationship with reduction of liver enzymes activity, in the present study, the decrease in liver enzymes in the treatment groups receiving the extract seems logical. There is a direct relationship between the amount of lipids and leptin, whereas there is an indirect relationship between the level of thyroid hormones and lipids. Furthermore, with increase in the amount of lipids and their deposition in the liver, the level of liver enzymes increases (Bush, 1991; Jelodar, Nazifi, 1997; Saeb *et al.*, 2010; Tohidi, 2007). By comparing the aqueous and alcoholic root extracts of *B. vulgaris*, Shahid *et al.* demonstrated that both extracts possess significant anti-inflammatory effects. The alcoholic extracts possibly play their role through blocking mediators which are released in the delaying phase of inflammation such as prostaglandin and the aqueous extract through blocking mediators, such as bradykinin, histamine, and serotonin, which are released in the primary phase and partly blocking the mediators in the delaying

phase, such as prostaglandin (Shahid *et al.*, 2009).

As it was mentioned before, increased lipids and liver inflammation result in an increase in liver enzymes and since *B. vulgaris* has anti-inflammatory properties and reduces cholesterol level, the decrease in liver enzymes seems reasonable (Noori *et al.*, 2004). Ivanovska *et al.*'s study on the effective substances in the alcoholic root extract of *B. vulgaris* indicated the presence of effective substances such as berberine and oxyacanthine with anti-inflammatory effects which their degree of influence depends, to a great extent, on their duration of use (Arayne *et al.*, 2007). Fallah Huseini *et al.*'s study on the effects of *B. vulgaris* L. Root extracts on carbon tetrachloride induced liver toxicity shows that *B. vulgaris* root extracts prevented CCl<sub>4</sub> induced hepatotoxicity in rats (Fallah Huseini *et al.*, 2010). Hyperlipidemia also triggers the production of free radicals (Heidarian *et al.*, 2008; Sharma *et al.*, 2009). In addition, in patients with liver toxicity, active metabolites formed through P<sub>450</sub> increase and result in liver necrosis. Here, the toxic amounts depend on glutathione level (Chiou, Tzeng, 2000; Callberg, Mannervik, 1985). TNF- and IL-6, on the other hand, are inflammatory cytokines that have a close relationship with leptin and lipid percentage (Boghrabadiet *et al.*, 2009).

Alkaline phosphatase is a trans-peptidase that increases in bone and liver diseases. Research has shown that phenolic compounds in medicinal plants can prevent the toxic effects of drugs on the liver and cause the release of glutamic pyruvic transaminase (SGPT) and alkaline phosphatase into blood (Nazari *et al.*, 2005).

### Conclusion

The findings of the present study show that alcoholic extract of *B. vulgaris* root could play an important role in lowering ALP

and ALT. However, more studies are needed to determine the underlying mechanisms.

### Acknowledgment

This article was part of the research project of the research committee and ethics committee No: 90-123-12 of Arak University of Medical Sciences, Arak, Iran. The authors would like to thank the moral and financial support provided by the Vice-presidency of Research of Arak University of Medical Sciences.

### Conflict of interest

There is not any conflict of interest in this study.

### References

- Arayne MS, Sultan N, Bahadur SS. 2007. The berberis story: *Berberis vulgaris* in therapeutics. *Pak J Pharm Sci*, 20: 83-92.
- Ashtiyani SC, Zarei A, Shariati M. 2011. [Certain plasma biochemical factors in rats.] In Persian. *Arak University of Medical Journal*, 14: 18-25.
- Boghrabadi V, Piri M, Sadeghi H, Sankian M. 2009. [Effects of aerobic training on leptin, tumor necrosis factor- and interleukin-6 levels in obese and lean men] In Persian. *Koomesh*, 11: 33-40.
- Bush BM. Interpretation of laboratory result for small animal clinicians. 1991. 1<sup>st</sup> ed., London, Blackwell Scientific Publication, 408-410.
- Callberg I, Mannervik B. 1985. Glutathione Reductase. *MethodS Enzymol*, 113: 484- 490.
- Chiou TJ, Tzeng WF. 2000. The roles of glutathione and antioxidant enzymes in menadioneinduced oxidative stress. *Toxicology*; 154 (1): 75-78.
- Fallah Huseini H, Zareei Mahmoudabady A, Ziai SA, Mehrazma M, et al. 2010. The Effects of *Taraxacum officinale* L. and *Berberis vulgaris* L. Root Extracts on Carbon Tetrachloride Induced Liver Toxicity in Rats. *Journal of Medicinal Plants*, 9: 45-52.
- Farhadi A Gavadifar K. 2008. [Effects of *Berberise vulgaris* fruit extract on blood cholesterol and triglyceride in hyperlipidemic patients.] In Persian. *Koomesh*, 9: 211-216.
- Gainsford T, Willson TA, Metcalf D, Handman E, and et al. 1996. Leptin can induce proliferation, differentiation, and functional activation of hemopoietic cells. *Proc Natl Acad Sci USA*, 93: 14564-14568.
- Gatreh-Samani K, Farrokhi E, Khalili B, Rafieian M, et al. 2011. [Purslane (*Portulaca oleracea*) effects on serum paraoxanase-1 activity.] Persian. *Journal of Shahrekord university of medical sciences*, 13: 9-15.
- Golfaraz M, Ahmad A. 2008. Hypoglysemic effect of methanolic extract of *Berberis*. *Phytother Res*, 22:1208-1212.
- Gosain S, Ircchiaya R, Sharma PC, Thareja S, Kalra A, et al. 2010. Hypolipidemic effect of ethanolic extract from the leaves of *Hibiscus sabdariffa* L. in hyperlipidemic rats. *Acta Pol Pharm*, 67:179-184.
- Heidarian E, Hajihossaini R, Jafari Dehkordy E, Omidi H. 2008. [Effects of Pistachio Nut Powder on Rat Liver Phosphatidate Phosphohydrolase and Serum Lipids and Lipoproteins Profile.] In Persian. *Journal of Ilam university of medical sciences*, 16: 47-55.
- Hersoug LG, Linneberg A. 2007. [The link between the epidemics of obesity and allergic diseases: does obesity induce decreased immune tolerance? Allergy] In Persian, 62: 1205-1213.
- Hu LF, Xu XY, Wang BQ. 2003. Research and utilization situation of *Portulaca oleracea* L. in China. *Practical Journal of Medicine and Pharmacy*, 20: 315-316.
- Hwong MJ. 2006. Berberin induced apoptosis through a mitochondria /caspases pathway in human hepatoma cells. *Archtoxicol*, 80: 62-73.
- Jellin JM, Batz F and Hitchens K. 2000. *Natural Medicines Comprehensive Database*. 3<sup>rd</sup> Ed. Stockton, California: Therapeutic Research Faculty.
- Jelodar GA, Nazifi habibabadi S. 1997. [Effect of celery, sour apple and carrot on some of serum biochemical parameters of diabetic rats] In Persian. *Journal of Kerman university of medical sciences*, 4: 114-119.
- Kaneda Y, Torii M, Tanaka T, Aikawa M. 1991. In vitro effects of berberine sulphate on the

- growth and structure of *Entamoeba histolytica*, *Giardia lamblia* and *Trichomonas vaginalis*. *Ann. Trop. Med. Parasitol.*, 85: 417-425.
- Kelly GE Husband AJ. 2003. Flavonoid compounds in the prevention of prostate cancer. *Methods Mol Med*, 81: 377-394.
- Lehrke M, Broedl UC, Biller-Friedmann IM, Vogeser M, and et al. 2008. Serum concentrations of cortisol, interleukin 6, leptin and adiponectin predict stress induced insulin resistance in acute inflammatory reactions. *crit care*, 12: R157.
- Meyer DJ, Coles EH and Rich LJ. 1992. *Veterinary Laboratory Medicin Interpretation and diagnosis*. Is ted WB. Saunders Co, 84-86.
- Mohammadi M, Bagheri A, Marashi H, Moshtaghi N, et al. 2011. Investigation into Seasonal Effect and Browning Inhibitor on Callus Regeneration of Seedless Barberry (*Berberis vulgaris* var. *asperma*), *Plant Tissue Cult & Biotech*, 21: 161-168.
- Motalleb G, Hanachi P. 2005. Evaluation of Phenolic content and total antioxidant activity in *Berberis vulgaris* fruit extracts. *J Biol*, 5: 648-653.
- Nazari A, Delfan B, Shahsavari G, 2005. [The effect of *Satureja khuzestanica* on triglyceride, glucose, creatinine and alkaline phosphatase activity in rat] Persian. *Journal of Shahrekord Unniversity of Medical Sciences*, 7: 1-8.
- Noori S, Naderi GA, Hassan ZM, Habib Z. 2004. Immunosuppressive activity of molecule isolated from *Artemisia annua* on DTH responses compared with cyclosporine A. *Immonopharmacol*, 4:1301-1306.
- Pourghassem-Gargari B, Ebrahimzadeh-Attary V, Rafraf M, Gorbani A. 2009. Effect of dietary supplementation with *Nigella sativa* L. on serum lipid profile, lipid peroxidation and antioxidant defense system in hyperlipidemic rabbits. *Journal of Medicinal Plants Research*, 3: 815-821.
- Rafati A, Moradi S, Ismaeili Dahadje M, et al. 2006. [Comparison of the efficiency of Lovostatin and seed dill extract in reduction of serum lipid levels in rats]. In Persian. *Journal of Shahid Sadoughi University of Medical Sciences and Health Services*, 13: 41-49.
- Saeb M, Nazifi S, Sabet M, Nazem H, and et al. 2010. [Effect of dietary wild pistachio oil on serum thyroid hormones, lipids and leptin concentration in experimental hyperthyroidism in male Rat] In Persian. *Journal of Gorgan University of Medical Sciences*, 11:8-15.
- Sharma A, Vija yacumar M, Rao ch V. Unnhkshnan M, K Reddy GAD. 2009. Action of portulaca oleracea against streptozotocin-induced oxidative stress in experimental diabetic rats. *Journal of complementary and Integrative Medicine*, 6: 12-18.
- Shahid M. Rahim T. Shahzad A, Tajuddin. Latif A, 2009. Ethnobotanical studies on *Berberis aristata* DC. Root extracts. *African Journal of Biotechnology*, 8: 556-563.
- Shariati M, Zarei A. 2006. [The study of physalis alkekengi extract on liver function.] In Persian. Azad university of Kazeran, MS.c Thesis.
- Thirupurasundari CJ, Padmini R, Devaraj SN. 2009. Effect of berberine on the antioxidant status, ultrastructural modifications and protein bound carbohydrates in azoxymethane-induced colon cancer in Rat. *Chemicobiological interactions*, 177: 190-195.
- Tohidi M, Harati H, Hadaegh F, Mehrabi Y and et al. 2007. Association of liver enzymes with incident type 2 diabetes: Teheran lipid and glucose study. *Iranian Journal of Diabetes and Lipid disorders*, 7: 167-176.
- Tomosaka H, Chin YW, Salim AA, Keller WJ, et al. 2008. Antioxidant and cytoprotective compounds from *Berberis vulgaris* (barberry). *Phytother Res*. 22: 979-981.
- Vrzal R, Zdarilova A, Ulrichova J. 2005. Activation of the aryl hydrocarbon receptor by berberin in H4ILE and HepG2 cells. *Scince direct*, 70: 925-936.
- Yu AS, Keeffe EB. 2003. Elevated AST or ALT to nonalcoholic fatty disease: accurate predictor of disease prevalence? *The American Journal of Gastroenterology*, 98: 955-956.
- Zarei A, Ashtiyani SC, Rasekh F, Mohammadi A, Jabary A. 2011. [The effect of physalis alkekengi extracts on lipids concentrations in



## Evaluation of the effects of hydroalcoholic extract of *Berberis vulgaris* root

- Rat] In Persian. Arak University of Medical Journal, 14: 48-55.
- Ziai SA, Salar F, Nasri S, Roghani M, Kamalinejad M. 2010. Neuroprotective Effect of Aqueous Extract of *Berberis vulgaris* L. in a Model of Parkinson's Disease in Rat. Journal of medicinal plants, 9: 24-33.
- Zovko Konci M, Kremer D, Karlovi K, Kosalec I. 2010. Evaluation of antioxidant activities and phenolic content of *Berberis vulgaris* L. and *Berberis croatica* Horvat. Food Chem Toxicol. 48: 2176-2180.

Archive of SID