

Review Article

A review on the inhibitory potential of *Nigella sativa* against pathogenic and toxigenic fungi

Hojjatollah Shokri¹

¹Faculty of Veterinary Medicine, Amol University of Special Modern Technologies, Amol, Iran

Article history:

Received: Feb 27, 2015

Received in revised form:

Jul 26, 2015

Accepted: Aug 10, 2015

Vol. 6, No. 1, Jan-Feb 2016,
21-33.

*** Corresponding Author:**

Tel: 01144153452

Fax: 01144153450

hshokri@ausmt.ac.ir

Keywords:

Nigella sativa

Thymoquinone

Anti-fungal and anti-
aflatoxigenic activity

Yeast

Dermatophyte

Aspergillus

Abstract

Nigella sativa (*N. sativa*) grows in various parts of the world, particularly in Iran. It has been traditionally used as a folk remedy to treat a number of diseases. The seeds of this plant contain moisture, proteins, carbohydrates, crude fiber, alkaloids, saponins, ash, fixed oils and essential oil. The major components of the essential oil are thymoquinone, p-cymene, trans-anethole, 2-methyl-5(1-methyl ethyl)-Bicyclo[3.1.0]hex-2-en and γ -terpinene. So far, several pharmacological effects such as anti-oxidant, anti-inflammatory, anti-cancer and anti-microbial have been reported for *N. sativa* or its active compounds. Thymoquinone, thymohydroquinone and thymol are the most active constituents which have different beneficial properties. The oil, extracts and some of *N. sativa* active components possessed moderate *in vitro* and *in vivo* inhibitory activity against pathogenic yeasts, dermatophytes, non-dermatophytic filamentous fungi and aflatoxin-producing fungi. The main morphological changes of pathogenic and toxigenic fungi treated with *N. sativa* oil were observed in the cell wall, plasma membrane and membranous organelles, particularly in the nuclei and mitochondria. Although this review represents first step in the search for a new anti-fungal drug, the full potential of *N. sativa* as a fungitoxic agent has not been exploited and necessitates further investigations.

Please cite this paper as:

Shokri H. A review on the inhibitory potential of *Nigella sativa* against pathogenic and toxigenic fungi. Avicenna J Phytomed, 2016; 6 (1): 21-33.

Introduction

The incidence of both community-acquired and nosocomial fungal infections has significantly increased over the past few decades, accompanying the growing number of high-risk patients, particularly those with impaired immunity (Kauffman, 2006). The majority of the clinically used antifungals has various drawbacks in terms of toxicity, drug-drug interactions, lack of fungicidal efficacy, cost and emergence of resistant strains caused by the frequent use of some of them (Rapp, 2004). In spite of the introduction of new anti-fungal drugs,

they are still limited in number. Hence, there is a great demand for novel anti-fungal agents, justifying the intense search for new drugs that are more effective and less toxic than those already in use.

Medicinal plants have been used for curing diseases for many countries in different indigenous systems of medicine as well as folk medicine. Essential oils are naturally occurring terpenic mixtures isolated from various parts of plants by steam distillation or other methods (Daferera et al., 2000). Since these oils are 'eco-friendly' and harmless to humans, at

present there is an increasing attention, both in industry and academic research, toward medicinal plants and their inhibitory properties against pathogenic and food spoilage fungi (Fan and Chen, 1999). Essential oils have been empirically used as anti-fungal agents, but the spectrum of activity and mechanisms of action remain unknown for most of them. Although only limited consistent information exists about their activity against human and animals fungal pathogens, some oils have shown important anti-fungal activity against yeasts, dermatophytes and non-dermatophytic filamentous fungi, suggesting their potential therapeutic effect on diseases involving mucosa, the skin and the respiratory tract (Pina-Vaz et al., 2004; Cavaleiro et al., 2006; Pinto et al., 2003). Although oils constitute complementary or alternative therapeutic options that are increasing in popularity, they still have scant scientific credibility. Among various medicinal plants, *Nigella sativa* (*N. sativa*) (family Ranunculaceae) is emerging as a miraculous herb with a rich historical and religious background as many investigators revealed a wide spectrum of pharmacological potential for it (Ali and Blunden, 2003). *N. sativa* is generally known as black seed and commonly grows in the Middle East, Eastern Europe and Western and Middle Asia. In French it is called nigelle and cumin noir, in German as schwarzkummel, in Italian as nigella, in Spanish as neguilla and pasionara, in Turkish as kolonji, in Hindi as kala zira, in Arabic as Habat-ulSauda and in English as black cumin. Among muslims, *N. sativa* is considered as one of the greatest source of healing medicine available because the black seed is the remedy for all diseases except death, according to prophet Muhammad. It is also recommended for use on regular basis in Tibb-e-Nabawi and identified as the curative black cumin in the Holy Bible, as the Melanthion of Hippocrates and Discroides and as the Gith

of Pliny (Junemann, 1998). Crude extracts and essential oil of *N. sativa* seeds have been reported to possess a number of pharmacological properties such as anti-oxidant (Burits and Bucar, 2000), anti-tumor (Ivankovic et al., 2006; Amara et al., 2008), anti-parasitic (EL Wakil, 2007), anti-inflammatory (Salem, 2005; Boskabady et al., 2010), anti-diabetic (Rchid et al., 2004), anti-bacterial (Ozmen et al., 2007; Mariam and Al-Basal, 2009), anti-fungal effects (Goreja, 2003; Randhawa and Al-Ghamdi, 2002; Shigeharu et al., 2006), protective activity against nephrotoxicity (Uz et al., 2008) Neurotoxicity (Khazdair, 2015) and hepatotoxicity (Mahmoud et al., 2002). This article aims to provide a review of the inhibitory effect of *N. sativa* against pathogenic and aflatoxin-producing fungi as well as describing ultrastructural changes in fungi treated with *N. sativa* oil.

Chemical compositions of *N. sativa*

Various varieties of *N. sativa* have a range of phytochemical components, of which only some molecules were characterized; so, complementary investigations are needed to identify new compounds in this species. Up to now, many investigations have been done on the seeds of *N. sativa* (Table 1). The main compounds were proteins, carbohydrates, fixed oils, essential oil, crude fiber, alkaloids, minerals, vitamins, ash and moisture. Other components were tannins, resin, saponin, carotene, glucosides and sterols (Randhawa and Al-ghamdi, 2002). α -sitosterol was a major sterol accounting for 44% and 54% of the total sterols in Tunisian and Iranian varieties of black seed oils, respectively (Cheikh-Rouhou et al., 2008). Selenium, DL- α -tocopherol, DL- γ -tocopherol, all trans retinol were among important anti-oxidants present in *N. sativa* seeds (Nasir et al., 2005; AL Saleh et al., 2006). Root and shoot were reported to contain vanillic acid (Al-Jassir, 1992).

Inhibition of pathogenic and toxigenic fungi by *Nigella sativa*

Table 1. Nutrient contents of *N. sativa*

Nutrients	%	Compositions	References
Moisture	4	-	Khan et al., 2003; El-Tahir and Bakheet, 2007
Total protein	20-22	arginine, glutamic acid, leucine, lysine, methionine, tyrosine, proline, threonine, etc.	Khan et al., 2003; El-Tahir and Bakheet, 2007
Total carbohydrate	17-34	-	Takruri and Dameh, 1998; Atta, 2003
Minerals	1.79-3.74	calcium, copper, iron, zinc, potassium, magnesium, phosphorus, sodium, manganese	Sultan et al., 2009; Youssef et al., 2013
Alkaloids	2.06	nigellidine, nigellimine	Atta-Ur-Rahman, 1995
Fixed oil	31-36	linoleic acid, oleic acid, palmitic acid, palmitoleic acid, myristic acid, stearic acid, linolenic acid, arachidic acid, lauric acid, eicosadienoic acid	Staphylakis and Gegiou, 1986; Asdadi et al., 2014; Nickavar et al., 2003
Essential oil	0.5-1.6	-	Zaoui et al., 2002; Morikawa et al., 2004; Morikawa et al., 2004; Ali et al., 2008; Mehta et al., 2009
Crude fiber	8-8.4	-	Khan et al., 2003; El-Tahir and Bakheet, 2007
Ash	4.5-4.8	-	Khan et al., 2003; El-Tahir and Bakheet, 2007
Vitamins	0.3	B ₁ , B ₂ , B ₆ , niacin, folic acid	Nergiz and Otles, 1993

The major components of the essential oil are thymoquinone (Venkatachallam et al., 2010; Mahmoudvand et al., 2014), p-cymene (Sultan et al., 2009; Toma et al., 2010; Sunita and Meenakshi, 2013), trans-anethole (Gerige et al., 2009; Shokri et al., 2012), 2-methyl-5-(1-methyl ethyl)-Bicyclo[3.1.0]hex-2-en (Adamu et al., 2010) and γ -terpinene (Golparvar et al., 2013). Some previous studies reported other chemical components including limonene, α -thujene, α -pinene, β -pinene, thymol, carvacrol, nigellone, anisaldehyde, n-nonane, miristicine, camphene, β -myrcene, 1,8-cineole, longipinene, camphor, linalool, estragole, junipene, germacrene, aromadendrene, borneol, bornylacetat, 2-tridecanone, 4-terpineol, sabinene, phencen, apiol, carvene, carvone, caryophyllene, avenasterol-5-ene, avenasterol-7-ene, campesterol, cholesterol, citrostadienol, cycloeucaenol, gramisterol, lophenol, obtusifoliol, longifolene, stigmastanol, stigmasterol-7-ene, β -amyrin, butyrospermol, cycloartenol, 24-methylene-cycloartanol, taraxerol, tirucallol, 3-O- $[\beta$ -D-xylopyranosyl (1 \rightarrow 3)- α -L-hamnopyranosyl (1 \rightarrow 2)- α -L-arabino-pyranosyl]-28-O- $[\alpha$ -L-rhamnopyranosyl (1 \rightarrow 4)- β -D-glucopyranosyl (1 \rightarrow 6)- β -D-glucopyranosyl] hederagenin, aliphatic alcohol, β -unsaturated hydroxy ketone,

melanthin, melanthigenin, 3-O- $[\beta$ -D-xylopyranosyl-(1 \rightarrow 2)- α -L-rhamnopyranosyl-(1 \rightarrow 2)- β -D-glucopyranosyl]-11-ethoxy-6, 23-dihydroxy-28-methyl-olean-12-enoate, stigma-5, 22-dien-3- β -D-glucopyranoside and cycloart-triene-23-ethyl-7 (Zaoui et al., 2000; Morikawa et al., 2004; Ali et al., 2008; Mehta et al., 2009).

Inhibitory effect of *N. sativa* against pathogenic yeasts

Candidiasis caused by *Candida* species has increased dramatically in recent years. *Candida* is the third- or fourth-most-common isolate in nosocomial bloodstream infections in the world. Among various species, *Candida albicans* (*C. albicans*) is the most causative agent associated with serious fungal infection, accounting for more than 90% of cases (Douglas, 2003). The difficulties associated with the management of *Candida* infections necessitate the discovery of new anti-fungal agents in order to increase the spectrum of activity against *Candida* and combat strains showing resistance to the available anti-fungal drugs. According to the literature, the investigation of natural products activity against *Candida* species increased significantly in the last 10 years, focusing on investigation of approximately

258 plant species from 94 families (Feldmesser, 2003; Duarte and Figueira, 2008). Table 2 summarizes the effect of *N. sativa* against different pathogenic yeasts as assessed by standard susceptibility methods. In general, a moderate efficacy of the oil from *N. sativa* against different *Candida* species were demonstrated

(Naeini et al., 2009; Shokri et al., 2012; Asdadi et al., 2014). In addition, several studies demonstrated that methanolic extract of *N. sativa* has the strongest anti-fungal effect against different strains of pathogenic yeasts, followed by ethanolic and chloroform extracts (Raval et al., 2010; Ahmad et al., 2013).

Table 2. Antifungal activity of *N. sativa* against different pathogenic yeasts

Fungus	Type	Susceptibility method	Growth Inhibition	References
<i>Candida albicans</i>	Essential oil	Disc diffusion	IZ: 35 mm**	Naeini et al., 2009
<i>C. albicans</i>	Essential oil	Broth macrodilution	MIC: 2300 µg/ml	Naeini et al., 2009
<i>C. zeylanoides</i>	Essential oil	Disc diffusion	IZ: 40.8 mm	Shokri et al., 2012
<i>C. albicans</i>	Essential oil	Broth macrodilution	MIC ₉₀ : 4.916 mg/ml MFC: 6.360 mg/ml	Asdadi et al., 2014
<i>C. dubliniensis</i>	Essential oil	Broth macrodilution	MIC ₉₀ : 5.183 mg/ml MFC: 6.360 mg/ml	Asdadi et al., 2014
<i>C. glabrata</i>	Essential oil	Broth macrodilution	MIC ₉₀ : 5.992 mg/ml MFC: 6.360 mg/ml	Asdadi et al., 2014
<i>C. krusei</i>	Essential oil	Broth macrodilution	MIC ₉₀ : 4.939 mg/ml MFC: 6.360 mg/ml	Asdadi et al., 2014
<i>C. parapsilosis</i>	Methanolic extract	Broth microdilution	MIC: 4.846 µg/ml	Raval et al., 2010
<i>C. albicans</i>	Methanolic extract	Broth microdilution	MIC: 6.484 µg/ml	Raval et al., 2010
<i>Issatchenkia orientalis</i>	Methanolic extract	Broth microdilution	MIC: 6.795 µg/ml	Raval et al., 2010
<i>I. orientalis</i>	Ethanolic extract	Broth microdilution	MIC: 5.805 µg/ml	Raval et al., 2010
<i>C. parapsilosis</i>	Ethanolic extract	Broth microdilution	MIC: 7.093 µg/ml	Raval et al., 2010

* IZ: Inhibition zone, ** mm: millimeter

In an experimental study by Khan et al. (2003), the aqueous extract of *N. sativa* seed exhibited inhibitory effect against candidiasis in mice. A 5-fold decrease in *Candida* organisms in kidneys, 8-fold in liver and 11-fold in spleen was observed in the groups of animals post-treated with the plant extract. It has been shown that the candidacidal pathway in mice neutrophils is nitric oxide (NO)-dependent (Fierro and Fidalgo, 1996). It is possible that the plant extract contains active ingredient(s), which may directly stimulate the granulocytes and monocytes to generate NO leading to an excellent anti-fungal activity, which in turn kills *C. albicans*. One of the anti-fungal actions of *Nigella* seed's oil may be attributed to the presence of β -sitosterol and oleic acid as the main components in the oil of *N. sativa* (Asdadi et al., 2014). Several previous studies have shown that long-chain fatty acid has a fungistatic effect against a few strains of *Candida* (Ouraini et al., 2007). In addition, Gupta et al. (2012) exhibited that different components of *N. sativa* oil such as β -sitosterol and stigmasterol have anti-fungal

activity against pathogenic yeasts such as *Candida tropicalis*, *C. albicans* and *Geotrichum candidum*. In a study by Sitheequ et al. (2009), all the polyphenols of black seed had anti-*Candida* activity against all tested *Candida* species such as *C. albicans* and *C. glabrata*, followed by *C. parapsilosis*, *C. krusei* and *C. tropicalis*.

The anti-yeast activity of quinones such as dithymoquinone and thymohydroquinone from *N. sativa* were evaluated *in vitro* using a broth microdilution method against six dairy spoilage yeast species. It is found that anti-fungal effects of quinones were comparable with those of preservatives commonly used in milk products (calcium propionate, natamycin and potassium sorbate), while thymohydroquinone possessed significant anti-yeast activity (Halamova et al., 2010). As reported by Taha et al. (2010), the minimum inhibitory concentration (MIC) of thymol was 0.5 mg/ml for *C. albicans*, *Rhodoturula rubra* (*R. rubra*) and *Trichosporon* species. *R. rubra* was the most sensitive yeast to

Inhibition of pathogenic and toxigenic fungi by *Nigella sativa*

thymoquinone (0.1 mg/ml), followed by *Trichosporum* species (0.25 mg/ml) and *C. albicans* (the resistant yeast). The MIC of thymohydroquinone was 0.5 mg/ml for all yeasts except *R. rubra*, which was inhibited at 0.25 mg/ml. In general, thymoquinone was the most potent active compound of *N. sativa* oil against the yeasts, followed by thymohydroquinone and thymol, respectively. In a comparative study on anti-fungal activity of tannins, saponins, alkaloids, essential oil, brown and yellow crystals separated from essential oil, all tested *N. sativa* constituents except for tannins showed inhibitory activity against *Saccharomyces cerevisiae* (Shohayeb and Halawani, 2012).

Inhibitory effect of *N. sativa* against dermatophytic fungi

One of the most important groups of fungi, which causes worldwide human and animal infections is the dermatophytes. They have the ability to invade keratinized tissues such as hair, skin and nails to

produce an infection, commonly referred to as dermatophytosis (Sidat et al., 2006). At present, there are many anti-dermatophytic drugs including imidazoles and terbinafine for topical, and triazoles and griseofulvin for systemic treatment of dermatophytosis, which exhibited some problems such as prolonged systemic therapy, fungal resistance, high toxicity and high cost (Martinez-Rossi et al., 2008). For these reasons, development of new drugs or combination therapy for treatment of dermatophytosis is urgent. Various studies have shown that plant extracts and plant-derived components are effective against dermatophytosis. Table 3 exhibited the efficacy of *N. sativa* against different dermatophytes by using standard susceptibility methods. Several studies showed a weak to moderate inhibitory effect against a considerable number of clinical isolates of various dermatophytes (Aljabre et al., 2005; Gerige et al., 2009; Khosravi et al., 2013; Sunita and Meenakshi, 2013; Mahmoudvand et al., 2014).

Table 3. Antifungal activity of *N. sativa* against different dermatophytes

Fungus	Type	Susceptibility method	Growth inhibition	References
<i>Trichophyton mentagrophytes</i>	Essential oil	Broth microdilution	MIC: 2±0.6 mg/ml MFC: 4±1.1 mg/ml	Khosravi et al., 2013
<i>Trichophyton rubrum</i>	Essential oil	Broth microdilution	MIC: 4±1.1 mg/ml MFC: 4±1.1 mg/ml	Khosravi et al., 2013
<i>Epidermophyton floccosum</i>	Essential oil	Broth microdilution	MIC: 2±0.6 mg/ml MFC: 4±1.1 mg/ml	Khosravi et al., 2013
<i>Microsporium gypseum</i>	Essential oil	Broth microdilution	MIC: 2±0.6 mg/ml MFC: 4±1.1 mg/ml	Khosravi et al., 2013
<i>Microsporium canis</i>	Essential oil	Broth microdilution	MIC: 4±1.1 mg/ml MFC: 8±2.8 mg/ml	Khosravi et al., 2013
<i>Trichophyton mentagrophytes</i>	Essential oil	Disc diffusion	IZ*: 0.0 mm** (resistant)	Gerige et al., 2009
<i>Trichophyton mentagrophytes</i>	Ether extract	Agar diffusion	MIC: 40 mg/ml	Aljabre et al., 2005
<i>Trichophyton interdigitale</i>	Ether extract	Agar diffusion	MIC: 40 mg/ml	Aljabre et al., 2005
<i>Trichophyton rubrum</i>	Ether extract	Agar diffusion	MIC: 40 mg/ml	Aljabre et al., 2005
<i>Microsporium canis</i>	Ether extract	Agar diffusion	MIC: 10 mg/ml	Aljabre et al., 2005
<i>Epidermophyton floccosum</i>	Ether extract	Agar diffusion	MIC: 40 mg/ml	Aljabre et al., 2005
<i>Trichophyton mentagrophytes</i>	Essential oil	Broth macrodilution	MIC: 4 mg/ml	Mahmoudvand et al., 2014
	Methanolic extract		MIC: 8 mg/ml	
	Aqueous extract		MIC: 16 mg/ml	
<i>Microsporium canis</i>	Essential oil	Broth macrodilution	MIC: 4 mg/ml	Mahmoudvand et al., 2014
	Methanolic extract		MIC: 4 mg/ml	
	Aqueous extract		MIC: 8 mg/ml	
<i>Microsporium gypseum</i>	Essential oil	Broth macrodilution	MIC: 4 mg/ml	Mahmoudvand et al., 2014
	Methanolic extract		MIC: 8 mg/ml	
	Aqueous extract		MIC: 16 mg/ml	
<i>Microsporium gypseum</i>	Essential oil	Disc diffusion	IZ: 38 mm	Sunita and Meenakshi, 2013
<i>Trichophyton rubrum</i>	Essential oil	Disc diffusion	IZ: 20 mm	Sunita and Meenakshi, 2013
<i>Trichophyton simii</i>	Essential oil	Disc diffusion	IZ: 35 mm	Sunita and Meenakshi, 2013

* IZ: Inhibition zone, ** mm: millimeter

As reported by Taha et al. (2010), thymol was found to be effective against *Trichophyton violaceum* (*T. violaceum*) (MIC 0.1 mg/ml), *Microsporum canis* (*M. canis*) (MIC 0.1 mg/ml) and *T. mentagrophytes* (MIC 0.05 mg/ml). The MIC of thymoquinone was found to be 0.05 mg/ml for tested dermatophytes. The MIC of thymohydroquinone was found to be 0.1 mg/ml for *M. canis*, while it showed various values (from 0.025 to 0.25 mg/ml) for *T. mentagrophytes* isolates. In general, thymoquinone was more efficient against the dermatophytes, followed by thymohydroquinone and thymol, respectively. Similarly, Ali and Blunden (2003) indicated that most of biological activity of the *N. sativa* seeds was due to thymoquinone, the major component of the essential oil obtained using soxhlet extraction. Thus, high anti-dermatophytic activity of *N. sativa* oil could be due to higher content of thymoquinone in comparison with methanolic and aqueous extracts, as previously described (Aljabre et al., 2005). Geweely and Alakilli (2012) isolated a variety of anti-fungal proteins from *N. sativa*. The crude proteins of *N. sativa* had highly significant anti-

dermatophytic activity on four zoophilic dermatophytes (*M. canis*, *M. equinum*, *T. mentagrophytes* and *T. verrucosum*). *N. sativa* proteins had considerable effects on the fungal cell permeability of all zoophilic dermatophytes. Two purified proteins (Pr1 and Pr2) from *N. sativa* showed higher anti-dermatophytic activities, representing 1.43-fold of the crude protein. The above-mentioned results denote the potential efficacy of *N. sativa* as a source of anti-dermatophytic drugs and support its use in folk medicine for the treatment of fungal infections.

Inhibitory effect of *N. sativa* against non-dermatophytic filamentous fungi

Literature survey showed that there is not enough research about anti-fungal activity of *N. sativa* against non-dermatophytic fungi. Table 4 exhibited the effect of *N. sativa* against different non-dermatophytic filamentous fungi by standard susceptibility methods. A relatively moderate activity of the oil and extracts of *N. sativa* against filamentous fungi was demonstrated (Sitara et al., 2008; Khosravi et al., 2011; Singh et al., 2015).

Table 4. Antifungal activity of *N. sativa* against non-dermatophytic filamentous fungi

Fungus	Type	Susceptibility method	Growth inhibition	References
<i>Aspergillus fumigatus</i>	Essential oil	Broth macrodilution	MIC ₉₀ : 1.5 mg/ml	Khosravi et al., 2011
<i>A. flavus</i>	Essential oil	Broth microdilution	MIC ₉₀ : 1.5 mg/ml	Khosravi et al., 2011
		Broth macrodilution	MIC ₉₀ : 1.5 mg/ml	
<i>A. niger</i>	Essential oil	Broth microdilution	MIC ₉₀ : 2 mg/ml	Sitara et al., 2008
		Agar diffusion	IZ*: 0.33 mm**	
<i>A. flavus</i>	Essential oil	Agar diffusion	IZ: 0.16 mm	Sitara et al., 2008
<i>Fusarium moniliforme</i>	Essential oil	Agar diffusion	IZ: 0.0 mm	Sitara et al., 2008
<i>F. oxysporum</i>	Essential oil	Agar diffusion	IZ: 0.06 mm	Sitara et al., 2008
<i>F. nivale</i>	Essential oil	Agar diffusion	IZ: 0.260 mm	Sitara et al., 2008
<i>F. semitectum</i>	Essential oil	Agar diffusion	IZ: 0.0 mm	Sitara et al., 2008
<i>Drechslera hawiensis</i>	Essential oil	Agar diffusion	IZ: 0.0 mm	Sitara et al., 2008
<i>Alternaria alternata</i>	Essential oil	Agar diffusion	IZ: 0.0 mm	Sitara et al., 2008
<i>A. flavus</i>	Essential oil	Inverted petri plate	IZ: 45.7 mm	Singh et al., 2015
	Ethanol extract		IZ: 8.9 mm	
<i>A. niger</i>	Essential oil	Inverted petri plate	IZ: 43.6 mm	Singh et al., 2015
	Ethanol extract		IZ: 5.7 mm	
<i>F. graminearum</i>	Essential oil	Inverted petri plate	IZ: 39.7 mm	Singh et al., 2015
	Ethanol extract		IZ: 10 mm	
<i>F. moniliforme</i>	Essential oil	Inverted petri plate	IZ: 71.2 mm	Singh et al., 2015
	Ethanol extract		IZ: 17.8 mm	
<i>Penicillium viridicatum</i>	Essential oil	Inverted petri plate	IZ: 34.7 mm	Singh et al., 2015
	Ethanol extract		IZ: 9.9 mm	

* IZ: Inhibition zone, ** mm: millimeter

Inhibition of pathogenic and toxigenic fungi by *Nigella sativa*

In a study conducted by Taha et al. (2010), the various MICs of thymol were shown. *Fusarium oxysporum* (*F. oxysporum*) and *Cladosporium* species were the most sensitive isolates, which were inhibited at 0.1 mg/ml, whereas *Aspergillus niger* (*A. niger*) and *Penicillium chrysogenum* (*P. chrysogenum*) were the most resistant, which were inhibited at 0.5 mg/ml. The MIC of thymoquinone showed similar profile as that obtained by thymol. *F. oxysporum* (MIC 0.1 mg/ml) and *A. niger* (MIC 0.1 mg/ml) together with *P. chrysogenum* (MIC 0.75 mg/ml) were the most sensitive and resistant isolates, respectively. *Cladosporium* species were inhibited at 0.25 mg/ml. The same profile that was obtained by thymol was also obtained by thymohydroquinone. *F. oxysporum* and *Cladosporium* species were the most sensitive organisms to thymohydroquinone. Both of them were inhibited at 0.25 mg/ml. *A. niger* and *P. chrysogenum* were the most resistant organisms (MIC 0.75 mg/ml). In general, thymol was the most efficient chemical against the tested filamentous fungi, followed by thymoquinone and thymohydroquinone, respectively. In another study, two novel anti-fungal defensins, namely Ns-D1 and Ns-D2, were isolated from seeds of *N. sativa*. Both defensins displayed strong divergent anti-fungal activity towards a number of phytopathogenic fungi (Rogozhin et al., 2011).

Inhibition of aflatoxin production by *N. sativa*

Mycotoxin-producing fungi are important contaminants and destroyers of food and feedstuffs during the storage, rendering them inappropriate for human consumption by reducing their nutritive value and sometimes by producing mycotoxins (Kumar et al., 2007). Mycotoxins are toxic to human and animals, and cause significant reductions in crop yield resulting in economic loss

(Iheshiulor et al., 2011). The food and agriculture organization of the United Nations (FAO) estimated that at least 25% of the cereal grains in the world are contaminated by mycotoxins like aflatoxins (Bathnagar and Garcia, 2001). Beside synthetic agents, some plants and their active metabolites have been introduced as inhibitors of aflatoxin biosynthesis (Sakuda et al., 2000; Rasooli and Razzaghi-Abyaneh, 2004; Yoshinari et al., 2007; Roze et al., 2011). To our knowledge, little data have been reported on the effect of *N. sativa* against aflatoxin-producing fungi. In a study by Khosravi et al. (2011), the effect of the essential oil of *N. sativa* on growth and aflatoxin production of *A. parasiticus* was evaluated. The results indicated that *N. sativa* oil (MIC₉₀ 2.75 mg/ml and MFC 6.25 mg/ml) had a moderate activity against *A. parasiticus*. Essential oil of *N. sativa* (1.5 mg/ml) exhibited a growth inhibition percent of mycelia production by *A. parasiticus* in value of 67.4%. In addition, aflatoxin production was inhibited by 1.5 mg/ml of *N. sativa* oil, representing significant reductions in values of 91.4% for AFB₁, 100% for AFB₂, 98.5% for AFG₁, 87.3% for AFG₂ and 96.2% for total aflatoxin. The oil had a more marked effect on aflatoxin B₂ as compared to AFB₁. This study showed that the extent of inhibition in mycelial growth was associated with decreased levels of aflatoxin production. Interestingly, aflatoxin production was significantly inhibited at concentrations lower than fungistatic concentrations of the oil. There is a 10-kb gene cluster which controls the activity of aflatoxins biosynthesis pathway. Regarding different aflatoxins, it seems that some of the genes are more active than the others. The correlations between some gene expressions and production of different kinds of aflatoxins have been previously noted. It is speculated that the essential oil of *N. sativa* may affect the expression of some special aflatoxin gene (Yahyaraeyat et al., 2013). Further studies are needed to evaluate this

issue. In a study by Maraqa et al. (2007), the crude extract of *N. sativa* inhibited the production of three types of aflatoxins (B_1 , B_2 and G_1) at 5% (w/v) concentration, whereas *N. sativa* oil at a concentration of 3% completely inhibited aflatoxins B_1 , B_2 , G_1 and G_2 . The authors recommended that the anti-aflatoxin property of the oil is mainly related to its high phenolic content as demonstrated previously (Cosentino et al., 1999). In a similar study conducted by El-Nagerabi et al. (2012), *N. sativa* oil at a concentration of 3% significantly inhibited aflatoxin B_1 production by *A. flavus* and *A. parasiticus* by 47.9-58.3% and 32-48%, respectively. On the other hand, the mycelial fresh weights of the two fungal species were similar at the tested concentrations. *N. sativa* oil 3% against pure aqueous aflatoxin B_1 led to decrease of the concentration to 670 ppb in comparison with the control (690 ppb), representing no detoxification ability on aflatoxin B_1 . Moreover, Al-Ghasham et al. (2008) demonstrated that *N. sativa* reduces the toxic effect of AFB₁ in liver and kidneys of rats with aflatoxicosis, which may be related to their cytoprotective and anti-oxidant properties. Recently, Abdel-Wahhab and Aly (2005) found that *N. sativa* oil administration to rats fed with aflatoxin-contaminated diet resulted in significant protection against aflatoxicosis. Similarly, Hussein et al. (2000) found that dietary addition of *N. sativa* significantly ameliorated the adverse effects of dietary AFB₁ on Nile tilapia fish. To the best of our knowledge, there were no previous studies exploring the effect of *N. sativa* on other toxins produced by toxigenic fungi.

Ultrastructural changes of pathogenic and toxigenic fungi due to *N. sativa*

As mentioned above, various studies have been carried out to investigate the anti-fungal activity of *N. sativa*, but its exact mechanism of action was not well established. Furthermore, very few data have been documented on the morphological changes of pathogenic and/or toxigenic fungi grown in the

presence of *N. sativa* oil and its components. In a study conducted by Khosravi et al. (2011), *A. flavus* (toxigen) and *A. fumigates* (pathogen) exposed to 0.25, 0.5, 1, 1.5 and 2 mg/ml of *N. sativa* essential oil were processed for transmission electron microscopy (TEM) (Figures 1a and 1b; normal shapes). The early changes in fungal compartments in the presence of the lowest concentrations of oil (0.25 and 0.5 mg/ml) were noticed in both hyphae and conidia, showing abnormal shaped and swelled hyphae, high vacuolation of the cytoplasm accompanied by vacuole fusion (Figures 1c and 1d). Subsequent events were loss of normal conidia and hyphae shape, detachment of fibrillar layer of the cell wall (Figures 1e, 1f and 1g), destruction of hypha membranous organelles including nuclei and mitochondria, and finally disorganization of cytoplasmic contents accompanied by intensive degradation and lysis of the nucleus and mitochondria (Figure 1h). The most remarkable changes in fungal compartments were observed in fungi treated with the highest fungistatic concentrations of the oils (1.5-2 mg/ml). Destruction and breaking down of plasma membrane at different sites (Figure 1i), disorganization of conidial and hyphal cytoplasm and complete lysis of membranous organelles seemed to be resulting in cells death (Figures 1j and 1k).




Figure 1. Transmission electron micrographs of *Aspergillus* species (CW: Cell wall; PM: Plasma membrane; S: Septum; W: Woronin body).

Inhibition of pathogenic and toxigenic fungi by *Nigella sativa*

All together, the main changes were in the cell wall, plasma membrane and membranous organelles; in particular, in the nuclei and mitochondria. Interestingly, Badary et al. (2000) indicated that adding of thymoquinone as the main component of *N. sativa* seed to the drinking water of mice at a concentration of 0.03% for 3 months led to no sign of toxicity, except for a significant decrease in fasting plasma glucose concentration. Therefore, the *N. sativa* seeds and their derivatives appear to have a low level of toxicity and could be considered safe at low concentrations in the host cells.

The use of herbal drugs as complementary medicine is prevalent and has gained worldwide popularity. Many drugs are derived directly from plants; while others are chemically modified natural products. The original research articles published so far have confirmed the anti-fungal potential of *N. sativa* seeds. The oil, extracts and some of its active components, particularly thymoquinone, thymohydroquinone and thymol, possess moderate *in vitro* and *in vivo* inhibitory effect against pathogenic yeasts, dermatophytes, non-dermatophytic filamentous fungi and aflatoxin-producing fungi.

Acknowledgement

This study was funded by Research Council of Faculty of Veterinary Medicine, Amol University of Special Modern Technologies, Amol, Iran

Conflict of interest

The authors declare that they have no conflicts of interest concerning this article.

References

- Abdel-Wahhab MA, Aly SE. 2005. Antioxidant property of *Nigella sativa* (black cumin) and *Syzygium aromaticum* (clove) in rats during aflatoxicosis. *J Appl Toxicol*, 25: 218-223.
- Adamu HM, Ekanem EO, Bulama S. 2010. Identification of essential oil components from *Nigella sativa* seed by gas chromatography-mass spectroscopy. *Pak J Nutr*, 9: 966-967.
- Ahmad A, Husain A, Mujeeb M, Khan SA, Najmi AK, Siddique NA, Damanhoury ZA, Anwar F. 2013. A review on therapeutic potential of *Nigella sativa*: A miracle herb. *Asian Pac J Trop Biomed*, 3: 337-352.
- Al-Ghasham A, Ata HS, El-Deep S, Meki A, Shehada S. 2008. Study of protective effect of date and *Nigella sativa* on aflatoxin B1 toxicity. *Int J Health Sci*, 2: 26-44.
- Ali B, Blunden G. 2003. Pharmacological and toxicological properties of *Nigella sativa*. *Phytother Res*, 17: 299-305.
- Ali Z, Ferreira D, Carvalho P, Avery MA, Khan IA. 2008. Nigellidine-4-O-sulfite, the first sulfated indazole-type alkaloid from the seeds of *Nigella sativa*. *J Nat Prod*, 71: 1111-1112.
- Aljabre SH, Randhawa MA, Akhtar N, Alaklomy OM, Alqurashi AM, Aldossary A. 2005. Antidermatophyte activity of ether extract of *Nigella sativa* and its active principle, thymoquinone. *J Ethnopharmacol*, 101: 116-119.
- Al-Jassir MS. 1992. Chemical composition and microflora of black cumin (*Nigella sativa* L.) seeds growing in Saudi Arabia. *Food Chem*, 45: 239-242.
- Al Saleh IA, Billedo G, Inam IE. 2006. Level of selenium, DL- α -tocopherol, DL- γ -tocopherol, all trans retinol, thymoquinone and thymol in different brands of *Nigella sativa* seeds. *J Food Comp Anal*, 19: 167-175.
- Amara AA, El-Masry MH, Bogdady HH. 2008. Plant crude extracts could be the solution: extracts showing *in vivo* antitumorigenic activity. *Pak J Pharm Sci*, 21: 159-171.
- Asdadi A, Harhar H, Gharby S, Bouzoubaâ Z, Yadini AE, Moutaj R, Hadek ME, Chebli B, Hassani LMI. 2014. Chemical composition and antifungal activity of *Nigella Sativa* L. oil seed cultivated in Morocco. *Int J Pharma Sci Invent*, 3: 9-15.
- Atta MB. 2003. Some characteristics of *Nigella* (*Nigella sativa* L.) seed cultivated in Egypt and its lipid profile. *Food Chem*, 83: 63-68.

- Atta-Ur-Rahman. 1995. Nigellidine-a new indazole alkaloid from the seed of *Nigella sativa*. *Tetrahedron Lett*, 36: 1993-1994.
- Badary OA, Abdel-Naim AB, Abdel-Wahab MH, Hamada FM. 2000. The influence of thymoquinone on doxorubicin-induced hyperlipidemic nephropathy in rats. *Toxicol*, 143: 219-226.
- Boskabady MH, Mohsenpoor N, Takaloo L. 2010. Antiasthmatic effect of *Nigella sativa* in airways of asthmatic patients. *Phytomed*, 17: 707-713.
- Bathnagar D, Garcia S. 2001. *Aspergillus*. In Labbe RG, Garcia S. (Eds.), *Guide to Foodborne Pathogens*. New York: John Wiley and Sons, pp. 35-49.
- Burits M, Bucar F. 2000. Anti-oxidant activity of *Nigella sativa* oil. *Phytother Res*, 14: 323-328.
- Cavaleiro C, Pinto E, Goncalves MJ, Salgueiro L. 2006. Antifungal activity of *Juniperus* essential oils against dermatophyte, *Aspergillus* and *Candida* strains. *J Appl Microbiol*, 100: 1333-1338.
- Cheikh-Rouhou S, Besbes S, Lognay G, Blecker C, Deroanne C, Attia H. 2008. Sterol composition of black cumin (*Nigella sativa* L.) and Aleppo pine (*Pinus halpensis* Mill.) seed oils. *J Food Comp Anal*, 21: 162-168.
- Cosentino S, Tuberoso CIG, Pisano B. 1999. In-vitro antimicrobial activity and chemical composition of Sardinian *Thymus* essential oils. *Lett Appl Microbiol*, 29: 130-136.
- Daferera DJ, Zircas BN, Polission MG. 2000. GC-MS analysis of essential oil from some Greek aromatic plants and their fungitoxicity on *Penicillium digitatum*. *J Agric Food Chem*, 48: 2576-2581.
- Douglas LJ. 2003. *Candida* biofilms and their role in infection. *Trends Microbiol*, 11: 30-36.
- Duarte MCT, Figueira GM. 2008. Anti-*Candida* activity of essential oils and extracts from native and exotic medicinal plants used in Brazil. In: Rai MK, Carpinella C. (Eds.), *Naturally Occurring Bioactive Compounds: A Newer and Safer Alternative for Control of Pest and Diseases*. The Haworth Press Inc.
- El-Nagerabi SAF, Al-Bahry SN, Elshafie AE, Al-Hilali S. 2012. Effect of *Hibiscus sabdariffa* extract and *Nigella sativa* oil on the growth and aflatoxin B1 production of *Aspergillus flavus* and *Aspergillus parasiticus* strains. *Food Control*, 25: 59-63.
- El-Tahir KE, Bakheet DM. 2007. The black seed *Nigella sativa* linnaeus-a mine for multi-cures: A plea for urgent clinical evaluation of its volatile oil. *J T U Med Sci*, 1: 1-19.
- El-Wakil SS. 2007. Evaluation of the in vitro effect of *Nigella sativa* aqueous extract on *Blastocystis hominis* isolates. *J Egypt Soc Parasitol*, 7: 801-813.
- Fan JJ, Chen JH. 1999. Inhibition of aflatoxin-producing by welsh onion extracts. *J Food Prot*, 62: 414-417.
- Feldmesser M. 2003. New and emerging antifungal agents: Impact on respiratory infections. *Am J Respir Med*, 2: 371-383.
- Fierro M, Fidalgo CB. 1996. The involvement of nitrous oxide in the anti-*Candida albicans* activity of rat neutrophils. *Immunol*, 89: 295-300.
- Iheshiulor OOM, Esonu BO, Chuwuka OK, Omede AA, Okoli IC, Ogbuewu IP. 2011. Effects of mycotoxins in animal nutrition. *Asian J Anim Sci*, 5: 19-33.
- Junemann M. 1998. *Three great healing herbs*, Lotus Light Publications, Twin Laked, WI, pp. 45.
- Gerige SJ, Gerige MKY, Rao M. 2009. GC-MS Analysis of *Nigella sativa* seeds and antimicrobial activity of its volatile oil. *Braz Arch Biol Technol*, 52: 1189-1192.
- Geweely NS, Alakilli SYM. 2012. Effect of the purification of antidermatophytic proteins from *Nigella sativa* on four zoophilic species. *Afr J Biotechnol*, 11: 9422-9434.
- Golparvar AR, Ghaisari MM, Hadipanah A, Armin A. 2013. Chemical analysis and identification of the components of black seed and Thyme cultivated in Iran. *Sci Agri*, 4: 55-57.
- Goreja WG. 2003. *Black seed: Nature's Miracle Remedy*, Amazing Herbs Press, New York, NY.
- Gupta S, Satishkumar MN, Duraiswamy B, Das S, Chhajed M. 2012. Potential herbs and its phytoconstituents against fungal infection: A systematic review. *World J Pharma Res*, 1: 1-20.
- Halamova K, Kokoska L, Flesar J, Sklenickova O, Svobodova B, Marsik P. 2010. In vitro antifungal effect of black cumin seed quinones against dairy spoilage

Inhibition of pathogenic and toxigenic fungi by *Nigella sativa*

- yeasts at different acidity levels. *J Food Prot*, 73: 2291-2295.
- Hussein SY, Mekkawy IAA, Moktar ZZ Mubarak M. 2000. Protective effect of *Nigella sativa* seed against aflatoxicosis in *Oreochromis niloticus*. *Proc. Conf. Mycotoxins and Dioxins and the Environment*, Bydgoszcz, 25-27 Sept., pp: 109-130.
- Ivankovic SR, Stojkovic M, Jukic M, Milos M. 2006. The antitumor activity of thymoquinone and thymohydroquinone in vitro and in vivo. *Exp Oncol*, 28: 220-224.
- Kauffman CA. 2006. Fungal infections. *Proc Am Thorac Soc*, 3: 35-40.
- Khan MA, Ashfaq MK, Zuberi HS, Zuberi AH. 2003. The in vivo anti-fungal activity of the aqueous extract from *Nigella sativa* seed. *Phytother Res*, 17: 183-186.
- Khazdair MR. 2015. The Protective Effects of *Nigella sativa* and Its Constituents on Induced Neurotoxicity. *J Toxicol*, 2015, 1-7.
- Khosravi AR, Minooeianhaghighi MH, Shokri H, Emami SA, Alavi SM, Asili J. 2011. The potential inhibitory effect of *Cuminum cyminum*, *Ziziphora clinopodioides* and *Nigella sativa* essential oils on the growth of *Aspergillus fumigatus* and *Aspergillus flavus*. *Braz J Microbiol*, 42: 216-224.
- Khosravi AR, Shokri H, Farahnejat Z, Chalangari R, Katalin M. 2013. Antimycotic efficacy of Iranian medicinal plants towards dermatophytes obtained from patients with dermatophytosis. *Chinese J Nat Med*, 11: 43-48.
- Khosravi AR, Shokri H, Minooeianhaghighi M. 2011. Inhibition of aflatoxin production and growth of *Aspergillus parasiticus* by *Cuminum cyminum*, *Ziziphora clinopodioides*, and *Nigella sativa* essential oils. *Foodborne Pathog Dis*, 8: 1275-1280.
- Kumar R, Mishra AK, Dubey NK, Tripathi YB. 2007. Evaluation of *Chenopodium ambrosioides* as a potential source of antifungal, antiaflatoxigenic and antioxidant activity. *Int J Food Microbiol*, 115: 159-164.
- Mahmoudvand H, Sharifi I, Fasihi Harandi M, Shokohi M, Shakibaie M, Rezaei Riabi T. 2014. Anti-leishmania effects of methotrexate (MTX) alone and in combination with meglumine antimoniate (MA) against Iranian isolate of sensitive and MA-resistant *Leishmania tropica*: an in-vitro assay. *Asian Pac J Trop Med*, 4: 412-420.
- Mahmoud MR, El-Abhar HS, Saleh S. 2002. The effects of *Nigella sativa* oil against the liver damage induced by *Schistosoma mansoni* in mice. *J Ethnopharmacol*, 79: 1-11.
- Maraqa A, Al-Sharoha NF, Farah H, Elbjeirami WM, Shakya AK, Sallal AJ. 2007. Effect of *Nigella sativa* extract and oil on aflatoxin production by *Aspergillus flavus*. *Turk J Biol*, 31: 155-159.
- Mariam A, Al-Basal A. 2009. In vitro and in vivo anti-microbial effects of *Nigella sativa* Linn. seed extracts against clinical isolates from skin wound infections. *Am J Appl Sci*, 6: 1440-1447.
- Martinez-Rossi NM, Peres NTA, Rossi A. 2008. Antifungal resistance mechanisms in dermatophytes. *Mycopathologia*, 166: 369-383.
- Mehta BK, Pandit V, Gupta M. 2009. New principles from seeds of *Nigella sativa*. *Nat Prod Res*, 23: 138-148.
- Morikawa T, Xu F, Ninomiya K, Matsuda H, Yoshikawa M. 2004. N. mines A3, A4, A5 and C, new dolabellane-type diterpene alkaloids with lipid metabolism-promoting activities from the Egyptian medicinal food black cumin. *Chem Pharm Bull*, 52: 494-497.
- Naeini A, Khosravi AR, Chitsaz M, Shokri H, Kamlnjad M. 2009. Anti-*Candida albicans* activity of some Iranian plants used in traditional medicine. *J Mycol Méd*, 19: 168-172.
- Nasir Z, Abid AR, Hayat Z, Shakoor HI. 2005. Effect of kalongi (*Nigella sativa*) seeds on egg production and quality in white Leghorn layers. *J Anim Plant Sci*, 15: 22-24.
- Nergiz C, Otles S. 1993. Chemical composition of *Nigella sativa* L. seeds. *Food Chem*, 48: 259-261.
- Nickavar B, Mojab F, Javidnia K, Amoli MA. 2003. Chemical composition of the fixed and volatile oils of *Nigella sativa* L. from Iran. *Z Naturforsch C*, 58: 629-631.
- Ouraïni D, Agoumi A, Ismaili-Alaoui M, Alaoui K, Cherrah Y, Alaoui MA, Belabbas MA. 2007. Activité antifongique de l'acide oléique et des huiles essentielles de *Thymus saturejoides* L. et de *Mentha pulegium* L., comparée aux antifongiques dans les dermatoses mycosiques. *Phytother*, 5: 6-14.

- Ozmen A, Gamze B, Tugba A. 2007. Antimitotic and antibacterial effects of the *Nigella sativa* L Seed. *Cayological*, 60: 270-272.
- Pina-Vaz C, Rodrigues AG, Pinto E, Costa-de-Oliveira S, Tavares C, Salgueiro LR, Cavaleiro C, Goncalves MJ, Martinez-de-Oliveira J. 2004. Antifungal activity of Thymus oils and their major compounds. *J Eur Acad Dermatol*, 18: 73-78.
- Pinto E, Palmeira A, Salgueiro L, Cavaleiro C, Goncalves MJ, Pina-Vaz C, Rodrigues A, Oliveira S. 2003. Antifungal activity of oregano oils (*Lippia graveolens* and *Origanum virens*) on dermatophyte species. *Clin Microbiol Infect*, 9: 222-230.
- Rapp RP. 2004. Changing strategies for the management of invasive fungal infections. *Pharmacother*, 24: 4S-28S.
- Rasooli I, Razzaghi-Abyaneh M. 2004. Inhibitory effects of thyme oils on growth and aflatoxin production by *Aspergillus parasiticus*. *Food Control*, 1: 479-483.
- Raval BP, Shah TG, Suthar MP, Ganure AL. 2010. Screening of *Nigella Sativa* Seeds for antifungal activity. *Ann Biolog Res*, 1: 164-171.
- Rchid H, Chevassus H, Nmila R, Guiral C, Petit P, Chokairi M. 2004. *Nigella sativa* seed extracts enhance glucose-induced insulin release from rat-isolated langerhans islets. *Fundam Clin Pharmacol*, 18: 525-529.
- Randhawa MA, Al-Ghamdi MS. 2002. A review of the pharmacotherapeutic effects of *Nigella sativa*. *Pak J Med Res*, 41: 77-83.
- Rogozhin EA, Oshchepkova YI, Odintsova TI, Khadeeva NV, Veshkurova ON, Egorov TA. 2011. Novel antifungal defensins from *Nigella sativa* L. seeds. *Plant Physiol Biochem*, 49: 131-137.
- Roze LV, Koptina AV, Laivenieks M, Beaudry RM, Jones DA, Kanarsky AV, Linz JE. 2011. Willow volatiles influence growth, development, and secondary metabolism in *Aspergillus parasiticus*. *Appl Microbiol Biotechnol*, 92: 359-370.
- Sakuda S, Ono M, Ikeda H, Nakamura T, Inagaki Y, Kawachi R, Nakayama J, Suzuki A, Isogai A, Nagasawa H. 2000. Blastidin A as an inhibitor of aflatoxin production by *Aspergillus parasiticus*. *J Antibiotics*, 68: 407-412.
- Salem ML. 2005. Immunomodulatory and therapeutic properties of the *Nigella Sativa* L. Seed. *Int Immunopharmacol*, 5: 1749-1770.
- Shigeharu I, Katsuhisa U, Toshio T, Hideyo Y, Shigeru A. 2006. Evaluation of the effect of terpenoid quinones on *Trichophyton mentagrophytes* by solution and vapor contact. *J Infect Chemother*, 12: 100-104.
- Shohayeb M, Halawani E. 2012. Comparative antimicrobial activity of some active constituents of *N. sativa* L. *World Appl Sci J*, 20: 182-189.
- Shokri H, Sharifzadeh A, Ashrafi Tamai I. 2012. Anti-*Candida zeylanoides* activity of some Iranian plants used in traditional medicine. *J Mycol Méd*, 22: 211-216.
- Sidat MM, Correia D, Buene TP. 2006. *Tinea capitis* among rural school children of the district of Magude, in Maputo province, Mozambique. *Mycoses*, 49: 480-483.
- Singh G, Marimuthu P, de Heluani CS, Catalan C. 2005. Chemical constituents and antimicrobial and antioxidant potentials of essential oil and acetone extract of *Nigella sativa* seeds. *J Sci Food Agric*, 85: 2297-2306.
- Singh SS, Singh DG, Schuff C, de Lampasona MP, Catalán CAN. 2015. Composition, in vitro antioxidant and antimicrobial activities of essential oil and oleoresins obtained from black cumin seeds (*Nigella sativa* L.). *BioMed Res Int*, In Press.
- Sitara U, Niaz I, Naseem J, Sultana N. 2008. Antifungal effect of essential oils on in vitro growth of pathogenic fungi. *Pak J Bot*, 40: 409-414.
- Sitheequ MAM, Panagoda GJ, Yau J, Amarakoon AMT, Udagama URN, Samaranayake LP. 2009. Antifungal activity of black tea polyphenols (catechins and thea flavins) against *Candida* Species. *Chemother*, 55: 189-196.
- Staphylakis PK, Gegiou D. 1986. The sterols of *Nigella sativa* seed oil. *Phytochem*, 25: 761-763.
- Sultan MT, Butt MS, Anjum FM, Jamil A, Akhtar S, Nasir M. 2009. Nutritional profile of indigenous cultivar of black cumin seeds and antioxidant potential of its fixed and essential oil. *Pak J Bot*, 41: 1321-1330.
- Sunita M, Meenakshi S. 2013. Chemical composition and antidermatophytic activity

Inhibition of pathogenic and toxigenic fungi by *Nigella sativa*

- of *Nigella sativa* essential oil. *Afr J Pharma Pharmacol*, 7: 1286-1292.
- Taha M, Abdel Azeiz AZ, Saudi W. 2010. Antifungal effect of thymol, thymoquinone and thymohydroquinone against yeasts, dermatophytes and non-dermatophyte molds isolated from skin and nails fungal infections. *Egypt J Biochem Mol Biol*, 28: 109-126.
- Takruri HRH, Dameh MAF. 1998. Study of nutritional value of black cumin seeds (*Nigella sativa* L.). *J Sci Food Agric*, 76: 404-410.
- Toma CC, Simu GM, Hanganu M, Olah N, Vata FMG, Hammami C, Hammami M. 2010. Chemical composition of the Tunisian *Nigella sativa*. Note I. Profile on essential oil. *Farmacia*, 58: 458-464.
- Uz E, Bayrak O, Uz E, Kaya A, Baayrak R. 2008. *Nigella sativa* oil for prevention of chronic cyclosporine nephrotoxicity: An experimental model. *Am J Nephrol*, 28: 517-522.
- Venkatachallam UKT, Pattekhani H, Divakar S, Kadimi US. 2010. Chemical composition of *Nigella sativa* L. seed extracts obtained by supercritical carbon dioxide. *J Food Sci Technol*, 47: 598-605.
- Yahyaraeyat R, Khosravi AR, Shahbazzadeh D, Khalaj V. 2013. The potential effects of *Zataria multiflora* Boiss essential oil on growth, aflatoxin production and transcription of aflatoxin biosynthesis pathway genes of toxigenic *Aspergillus parasiticus*. *Braz J Microbiol*, 44: 649-655.
- Yoshinari T, Akiyama T, Nakamura K, Kondo T, Takahashi Y, Muraoka Y, Nonomura Y, Nagasawa H, Sakuda S. 2007. Diocatin A is a strong inhibitor of aflatoxin production by *Aspergillus parasiticus*. *Microbiol*, 153: 2774-2780.
- Youssef MKE, Eshak NS, Hana RS. 2013. Physicochemical characteristics, nutrient content and fatty acid composition of *Nigella sativa* oil and sesame oil. *Food Pub Health*, 3: 309-314.
- Zaoui A, Cherrah Y, Lacaille-Dubois MA, Settaf A, Amarouch H, Hassar M. 2000. Diuretic and hypotensive effects of *Nigella sativa* in the spontaneously hypertensive rat. *Therapie*, 55: 379-382.