

Original Research Article

The neuroprotective mechanism of cinnamaldehyde against amyloid- β in neuronal SHSY5Y cell line: The role of N-methyl-D-aspartate, ryanodine, and adenosine receptors and glycogen synthase kinase-3 β

Masoumeh Emamghoreishi^{1,2}, Majid Reza Farrokhi³, Atena Amiri³, Mojtaba Keshavarz^{3,*}

¹ Department of Pharmacology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran.

² Department of Neuroscience, School of Advanced Medical Sciences and Technologies, Shiraz University of Medical Sciences, Shiraz, Iran.

³ Shiraz Neuroscience Research Center, Shiraz University of Medical Sciences, Shiraz, Iran.

Article history:

Received: Jul 02, 2018

Received in revised form:

Jan 16, 2019

Accepted: Jan 17, 2019

Vol. 9, No. 3, May-Jun 2019,
271-280.

* Corresponding Author:

Tel: +9871 3623 4508

Fax: +9871 3623 4508

mkeshavar@sums.ac.ir

moj.ph60@yahoo.com

Keywords:

Adenosine

Cinnamaldehyde

Dantrolene

Glycogen synthase kinase

Neuroprotection

N-methyl-D-aspartate

Abstract

Objective: Cinnamaldehyde may be responsible for some health benefits of cinnamon such as its neuroprotective effects. We aimed to investigate the cinnamaldehyde neuroprotective effects against amyloid beta ($A\beta$) in neuronal SHSY5Y cells and evaluate the contribution of N-methyl-D-aspartate (NMDA), ryanodine, and adenosine receptors and glycogen synthase kinase (GSK)-3 β , to its neuroprotective effects.

Materials and Methods: After seeding the cells in 96-well plates, adenosine (20, 40, 80, and 120 μ M), NMDA (20, 40, 80, and 120 μ M), and dantrolene (as a ryanodine receptor antagonist; 2, 4, 6, 8, and 16 μ M) were added to the medium containing $A\beta$ 25-35 and/or cinnamaldehyde. The 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyl tetrazolium bromide method was used to assess neurotoxicity and western blot to measure the GSK-3 β protein level.

Results: Cinnamaldehyde (15, 20, 23, and 25 μ M) significantly reversed $A\beta$ -induced toxicity in SHSY5Y neuronal cells. Adenosine (20, 40, 80 and 120 μ M) inhibited the neuroprotective effects of cinnamaldehyde (15 μ M). NMDA (20, 40, 80, and 120 μ M) reduced cinnamaldehyde (15 and 23 μ M) neuroprotective effects against $A\beta$ neurotoxicity. Dantrolene (2, 4, 8, and 16 μ M) significantly reduced cinnamaldehyde (15 μ M) neuroprotective effects. Cinnamaldehyde (15 and 23 μ M) suppressed the $A\beta$ -induced increment of GSK-3 β protein level.

Conclusion: NMDA and adenosine receptors suppression together with ryanodine receptors stimulation may be relevant to cinnamaldehyde neuroprotective effects against $A\beta$ neurotoxicity. Moreover, the inhibition of GSK-3 β may contribute to the cinnamaldehyde neuroprotection.

Please cite this paper as:

Emamghoreishi M, Farrokhi MR, Amiri A, Keshavarz M. The neuroprotective mechanism of cinnamaldehyde against amyloid- β in neuronal SHSY5Y cell line: The role of N-methyl-D-aspartate, ryanodine, and adenosine receptors and glycogen synthase kinase-3 β . Avicenna J Phytomed, 2019; 9(3): 271-280.

Introduction

Alzheimer's disease (AD) is the most popular neurodegenerative disorder which is characterized by cognitive impairment especially in older persons (Lindeboom and Weinstein, 2004). The extracellular deposition of β -amyloid ($A\beta$) plaques and aggregation of hyperphosphorylated form of tau protein (neurofibrillary tangles) are classic signs of AD (Checler, 1995; Delacourte and Buée, 2000). Recent evidence has shown that glycogen synthase kinase-3 β (GSK-3 β) may enhance the $A\beta$ -induced tau phosphorylation and progression of AD pathophysiology (Hernandez et al., 2013). Furthermore, deregulated calcium homeostasis may activate apoptotic pathways and cause neural cell death in AD (LaFerla, 2002). Abnormal intracellular calcium homeostasis may precede pathophysiological changes in AD (Chui et al., 1999).

Calcium as an essential second messenger (Berridge et al., 1998) regulates neural cell function and viability (Berridge et al., 1998). N-methyl-D-aspartate (NMDA), adenosine, and ryanodine receptors are important systems that regulate the intracellular calcium concentration (Gomes et al., 2011; Lüscher and Malenka, 2012; McPherson et al., 1991). These receptors may be connected to the pathophysiology and treatment of AD (Bruno et al., 2012; Danysz and Parsons, 2012).

Cinnamomum verum extract showed a neuroprotective effect in AD models (Frydman-Marom et al., 2011; Peterson et al., 2009). Moreover, the bioactive compounds of cinnamon were shown to reduce GSK-3 β activity in the peripheral tissues (Imparl-Radosevich et al., 1998). The therapeutic potential of cinnamon is mainly related to its phytochemicals like cinnamaldehyde, cinnamyl, and eugenol (Stavinoha and Vattam, 2015). However, the active compound responsible for the neuroprotective effects of cinnamon is unknown.

Cinnamaldehyde is one of the main components of cinnamon (Chen et al., 2016) with beneficial effects on inflammatory and oxidative stress, blood glucose and malignancy (Anderson et al., 2004; Ataie et al., 2019; Kwon et al., 1997; Zhao et al., 2015). Cinnamaldehyde may be also responsible for some health benefits of cinnamon such as its neuroprotective effects (Stavinoha et al., 2015). In this regard, cinnamaldehyde was shown to reduce neuroinflammation, microglia activation and tau aggregation (Ho et al., 2013; Peterson et al., 2009). However, the potential effects and mechanism of action of cinnamaldehyde against $A\beta$ neurotoxicity are relatively unknown. Therefore, we aimed to assess the cinnamaldehyde neuroprotective effects in SHSY5Y cells exposed to $A\beta$ and the involvement of NMDA, ryanodine and adenosine receptors in its neuroprotective effects. Moreover, we tried to explore the relationship between GSK-3 β inhibition and neuroprotective effects of cinnamaldehyde.

Materials and Methods

Human SHSY5Y neuroblastoma cell line were obtained from Pasteur Institute (Iran). The cell culture reagents including DMEM/F12, fetal bovine serum (FBS), and penicillin-streptomycin were purchased from Gibco® life technologies™ (USA). We purchased $A\beta$ 25-35, NMDA, cinnamaldehyde (W228613), dantrolene sodium salt, and adenosine from Sigma-Aldrich (USA). The anti-actin, anti-GSK-3 β , and anti-phosphorylated (p)-GSK-3 β antibodies were obtained from Cell Signaling Technology® (USA). The phosphate buffered solutions (PBS) was used as the solvent of cinnamaldehyde, dantrolene, adenosine, and NMDA.

Neuronal Cell Culture

The seeded cells (1×10^5 cells/well) were kept in 96-well plates and in a Dulbecco's modified Eagle's medium (DMEM) plus Ham's nutrient mixture F-12 (1:1). Other reagents including 10% FBS, 100 U/ml

penicillin, and 100 µg/ml streptomycin were added to the plates. Then, the plates were maintained in the condition of 95% air/5% CO₂ at 37°C.

Treatment

We incubated Aβ_{25–35} for four days at 37°C to induce the aggregation process. The best concentrations of Aβ_{25–35} and cinnamaldehyde were determined in a pilot study. In the pilot study, we used different concentrations of cinnamaldehyde (1–50 µM) and chose the concentrations of 15 and 23 µM. We also chose 20 µM as the best concentration of Aβ_{23–35}. After replacement of serum-free media, Aβ_{23–35} (20 µM), cinnamaldehyde (15 and 23 µM) or both was added to the plates. Adenosine (20, 40, 80, and 120 µM), NMDA (20, 40, 80, and 120 µM), and dantrolene (2, 4, 6, 8, and 16 µM) were added to the culture media containing Aβ_{23–35}, cinnamaldehyde or both in independent experiments (4 experiment).

Cell viability assay

Cell viability was measured using 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyl tetrazolium bromide (MTT) reagent (5 mg/ml) 24h after the treatment. The cells were incubated with MTT for 4h, and then, the precipitate was dissolved in 100 µl of dimethyl sulfoxide (DMSO). The absorbance was determined at a 570 nm by a microplate reader (Synergy HT, Biotek®).

Total protein determination

The SHSY5Y cells were transported to the 6-well plates (10⁶ cells/ml) for protein analysis. After incubation with cinnamaldehyde and Aβ, cells were removed by centrifuging at 14000 g for 5 min. The radio-immunoprecipitation assay lysis (RIPA) buffer containing protease and phosphatase inhibitor cocktail was used to lyse the cells. The lysate cells were centrifuged at 19000 g for 25 min at 4°C to eliminate cell debris. The soluble part was used to determine total protein level. The

Lowry method was used to determine the total protein level (Lowry et al., 1951).

Western blot analysis

SDS-PAGE separated the proteins and equal amounts of proteins were transblotted onto polyvinylidene fluoride (PVDF) membrane. For blocking, the membrane was incubated with 5% bovine serum albumin (BSA) for 1 hr at room temperature. Then, the membrane incubated with the GSK-3β rabbit monoclonal antibody, p-GSK-3β (Ser9) antibody, and actin overnight at 4°C. The membrane was washed with TBST (tris-buffered saline and tween 20) buffer. After that, the anti-rabbit IgG, HRP-linked antibody (7074s, Cell Signaling) (1:25000) was poured on the membrane and kept for 1 hr at 37°C. Then, we scanned the membrane using an enhanced chemiluminescence kit (GE Healthcare) and ChemiDoc™ XRS+ imaging System. The bands were analyzed by image-J software.

Statistical analysis

The Shapiro-Wilk test was used for the evaluation of the normal distribution. The data were analyzed with the one-way analysis of variance (ANOVA) followed by the Bonferroni test. The significance level was set at 0.05. All the analyses were executed using SPSS software version 23.

Results

Cinnamaldehyde neuroprotection against Aβ neurotoxicity

Aβ (20 µM) reduced neural cell survival by 57%. Cinnamaldehyde (15, 20, 23, and 25 µM) significantly reversed Aβ neurotoxicity (F(5)=14.66, p=0.000) (Figure 1). However, cell viability was not different between cinnamaldehyde 15 µM and cinnamaldehyde 23 µM groups (p>0.05).




Figure 1. Effect of cinnamaldehyde (Cinn) on the viability of SHSY-5Y cells in the presence of beta amyloid (Aβ). The percent of viable cells was determined using MTT assay and normalized against untreated cells. Data are presented as the mean ± standard error of the mean of four experiments. TTT: p<0.001 compared to the vehicle-treated cells (white column); ***: p<0.001 compared to the Aβ-treated cells (black column).

Adenosine interference with cinnamaldehyde neuroprotection against Aβ

Adenosine (without cinnamaldehyde) only at the concentration of 80 μM

produced a neuroprotective effect against Aβ neurotoxicity (p=0.043) (Figure 2). Adenosine (20, 40, 80 and 120 μM) inhibited cinnamaldehyde neuroprotective effects (15 μM) against Aβ (F(9)=10.36, p=0.000) (Figure 2). Adenosine had no effect on cinnamaldehyde (23 μM) neuroprotective effects (Figure 2).

NMDA interference with cinnamaldehyde neuroprotection against Aβ neurotoxicity

The present study showed that NMDA (20, 40, 80, and 120 μM) decreased the cinnamaldehyde (15 and 23 μM) neuroprotective effects (F(9)=26.41, p=0.000, and F(9)=14.35, p=0.000, respectively) (Figure 3). Moreover, NMDA alone (20, 80, and 120 μM) had no effect on Aβ neurotoxicity (F(3)=0.66, p=0.59).




Figure 2. Effect of cinnamaldehyde (Cinn) on the viability of SHSY-5Y cells in the presence of adenosine (Ad) and beta amyloid (Aβ). The percent of viable cells was determined using MTT assay and normalized against untreated cells. Data are presented as the mean ± standard error of the mean of four experiments; ***: p<0.001 compared to the Aβ-treated cells (black column); †††: p<0.001 compared to the Aβ+Cinn (15 μM)-treated cells.




Figure 3. Effect of cinnamaldehyde (Cinn) on the viability of SHSY-5Y cells in the presence of N-Methyl-D-aspartate (NMDA) and beta amyloid (Aβ). The percent of viable cells was determined using MTT assay and normalized against untreated cells. Data are presented as the mean ± standard error of the mean of four experiments. ***: p<0.001 compared to the Aβ-treated cells (white column); †: p<0.05 and †††: p<0.001 compared to the Aβ+Cinn (15 μM)-treated cells; ‡‡‡: p<0.001 compared to the Aβ+Cinn (23 μM)-treated cells

Dantrolene interference with the cinnamaldehyde neuroprotection against Aβ neurotoxicity

Dantrolene (2, 4, 8, and 16 μM) significantly reduced cinnamaldehyde (15 μM) neuroprotective effects (F(9)=13.76, p=0.000) (Figure 4).

Moreover, dantrolene (4 and 16 μM) suppressed the neuroprotective effects of cinnamaldehyde (23 μM) (Figure 4). Furthermore, dantrolene alone (2, 8 and 16 μM) had no effect on the Aβ-induced neurotoxicity (p>0.05).




Figure 4. Effect of cinnamaldehyde (Cinn) on the viability of SHSY-5Y cells in the presence of dantrolene (Dant) and beta amyloid (Aβ). The percent of viable cells was determined using MTT assay and normalized against untreated cells. Data are presented as the mean ± standard error of the mean of four experiments. ***: p<0.001 compared to the Aβ-treated cells (white column); †: p<0.05 and †††: p<0.001 compared to the Aβ+Cinn (15 μM)-treated cells; ‡‡‡: p<0.001 compared to the Aβ+Cinn (23 μM)-treated cells

Cinnamaldehyde effects on the GSK-3 β protein

The expression of GSK-3 β protein was different in the studied groups (F(5)=172.483, p=0.000) (Figure 5). Treatment of neuronal cells with A β increased the total GSK-3 β protein levels compared to the control group (p=0.000) (Figure 5). Cinnamaldehyde 15 and 23 μ M diminished the A β effects on the total GSK-3 β protein levels (p<0.001) (Figure 5). Cinnamaldehyde (23 μ M) decreased the total GSK-3 β protein compared to the control group (p=0.004) (Figure 5).

The p-GSK-3 β protein levels in the studied groups were different (F(5)=12.972, p=0.000) (Figure 5). A β decreased, though with no statistical significance, the p-GSK-3 β protein level (p=0.057) (Figure 5). The p-GSK-3 β protein level in the A β +cinnamaldehyde (15 μ M) group was lower than the A β -treated group (p=0.034) (Figure 5). However, the p-GSK-3 β protein level in the A β +cinnamaldehyde (23 μ M) group was not different from the A β group (p=0.934) (Figure 5). Moreover, the p-GSK-3 β protein level in the cinnamaldehyde (15 and 23 μ M) groups were lower compared to the control group (p=0.001 and p=0.003, respectively) (Figure 5).




Figure 5. Effect of cinnamaldehyde (Cin) on the level of total and phosphorylated (p)-glycogen synthase kinase (GSK)-3 β in SHSY-5Y cells in the presence of beta amyloid (A β). The protein level was determined using Western blot analysis and normalized against untreated control cells. Data are presented as the mean \pm standard error of the mean of four experiments. †: p<0.05 and †††: p<0.001 compared to the A β -treated cells; ††††: p<0.0001 compared to the vehicle-treated cells (control).

Discussion

Our study showed that cinnamaldehyde, the main component of cinnamon, protected neuronal cells against A β neurotoxicity. Cinnamon extract inhibited A β neurotoxicity and improved cognitive function in an animal model of AD (Frydman-Marom et al., 2011). Furthermore, cinnamon decreased A β oligomerization and tau aggregation in cell culture and in an animal model of AD (Frydman-Marom et al., 2011; Peterson et al., 2009). Therefore, cinnamaldehyde may be regarded as an alternative therapy for the treatment of AD.

The present study showed that cinnamaldehyde reduced GSK-3 β protein level and prevented the A β -induced increases in GSK-3 β . Similarly, it was shown that cinnamon extract suppressed the GSK activation and reduced tau phosphorylation in the transfected HEK293 cells (Donley et al., 2016). Moreover, cinnamon decreased the GSK-3 β mRNA and protein levels in the liver and muscle of rats (Couturier et al., 2011). On the other hand, cinnamaldehyde did not affect GSK phosphorylation in primary astrocyte culture (Sartorius et al., 2014). Thus, our study revealed the inhibitory effects of cinnamaldehyde on the GSK-3 β protein in a neuronal cell line and showed the contribution of this effect to the neuroprotection against A β -induced neurotoxicity.

Adenosine is an important neuromodulator with potential roles in AD (Dall'Igna et al., 2007). However, the exact function of this endogenous substance should be clarified (Gomes et al., 2011). Our study showed that the activation of adenosine receptors might induce neuroprotection, while adenosine diminished the neuroprotective effects of cinnamaldehyde. These effects may imply that adenosine and cinnamaldehyde affect similar receptors and compete for binding to these receptors. To the best of our knowledge, there is no other study about possible interactions between

cinnamaldehyde and adenosine receptors. However, other adenosine antagonist like caffeine protected neuronal cells against A β (Dall'Igna et al., 2003; Keshavarz et al., 2017). Moreover, an A2A agonist blocked the caffeine neuroprotective effects against A β (Dall'Igna et al., 2003). Caffeine and an A2A antagonist reversed A β -induced cognitive impairments in animal models (Dall'Igna et al., 2007). Thus, the adenosine receptors may be potential targets for the cinnamaldehyde neuroprotective effects. However, cinnamaldehyde only at a lower concentration diminished the effects of adenosine.

The inhibition of NMDA receptors may be involved the neuroprotective effects of cinnamaldehyde.

Similarly, cinnamaldehyde protected PC12 cells against NMDA-induced neurotoxicity (Lv et al., 2017). Furthermore, cinnamon extract produced neuroprotection against glutamate in primary neuronal culture (Shimada et al., 2000) and protected neurons against glutamate and NMDA neurotoxicity in the primary chick embryo neuronal culture (Gomada et al., 2012). An animal study showed that cinnamaldehyde increased the glutamate release (Klafke et al., 2012). However, cinnamaldehyde produced no effect on the glutamate binding (Klafke et al., 2012). The discrepancies among the studies may arise from methodological differences or the nature of used neurons.

Cinnamon neuroprotective effect may result from the modulation of intracellular calcium. Ryanodine receptors are sarcoplasmic reticulum calcium channels that modulate intracellular calcium in the brain and peripheral tissues (McPherson et al., 1991). Therefore, blockade of ryanodine receptors may diminish the neuroprotective effects of cinnamaldehyde. There are some controversies about the roles of ryanodine receptor in neurotoxicity and neuroprotection. In accordance with our study, dantrolene deteriorated A β -induced hippocampal neuronal damage in an animal model of AD (Zhang et al.,

2010). Furthermore, the inability to up-regulate ryanodine receptor-3 may be the cause of neuronal vulnerability against multiple insults like A β , oxidative stress, and excitotoxicity (Allan Butterfield, 2002). Accordingly, activation of the ryanodine receptor may be a cellular mechanism to protect neurons in the initial stage of AD (Supnet and Bezprozvanny, 2010). On the other hand, dantrolene ameliorated cognitive dysfunction by decreasing A β load in an animal model of AD (Peng et al., 2012).

GSK is a regulatory enzyme involved in several CNS functions like neuronal development and neurodegeneration (Bhat et al., 2004). Accordingly, GSK is an important target in the pathophysiology and treatment of neurodegenerative disorders including AD (P. Chen et al., 2007). The GSK-3 β protein may be a mediator in the neurotoxic effects of A β (Phiel et al., 2003; Takashima, 2006). Our study showed that cinnamaldehyde reversed the A β effects on the GSK-3 β protein levels. Similarly, Imparl-Radosevich and colleagues showed that cinnamon inhibited the GSK-3 β activity in the peripheral tissue (Imparl-Radosevich et al., 1998). Thus, cinnamaldehyde neuroprotective effects may be due to GSK-3 β inhibition. There is limited information regarding the cinnamaldehyde interactions with the GSK-3 β protein. Our study showed that NMDA activation suppressed the neuroprotective effects of cinnamaldehyde. On the other hand, it was shown that NMDA stimulation enhance the activity of GSK-3 β in primary neuronal culture and animal brain (Szatmari et al., 2005). Similarly, GSK-3 β activation enhanced NMDA activity in neurons (Szatmari et al., 2005). The interaction between NMDA receptors and GSK-3 β protein may induce neuronal toxicity. Inhibition of NMDA receptors and GSK-3 protein by cinnamaldehyde may produce neuroprotective effects.

The application of non-selective agonist and antagonists of adenosine and ryanodine receptors may be the main limitation of this

study. We assessed cinnamaldehyde effects on these two receptors. Future studies should employ selective modulators of these receptors. Moreover, the MTT method has some limitations. We suggest reassessing the results of this study by other apoptosis evaluation techniques.

Cinnamaldehyde protected neurons against A β . The exact mechanism of cinnamaldehyde neuroprotective effects should be studied in future. However, the neuroprotective effects of cinnamaldehyde may be related to inhibition of NMDA and adenosine receptors together with stimulation of ryanodine receptors. Inhibition of the GSK-3 β protein may enhance cinnamaldehyde neuroprotective effects.

Acknowledgment

This work was funded by Shiraz University of Medical Sciences [grant No. 13761]. We would like sincerely to thank deputy for research of Shiraz University of Medical Sciences for supporting this investigation.

Conflicts of interest

All the authors confirm that there were no conflicts of interest.

References

- Allan Butterfield D. 2002. Amyloid β -peptide (1-42)-induced oxidative stress and neurotoxicity: implications for neurodegeneration in Alzheimer's disease brain. A review. *Free Radic Res*, 36: 1307-1313.
- Anderson RA, Broadhurst CL, Polansky MM, Schmidt WF, Khan A, Flanagan VP, Schoene NW, Graves DJ. 2004. Isolation and characterization of polyphenol type-A polymers from cinnamon with insulin-like biological activity. *J Agric Food Chem*, 52: 65-70.
- Ataie Z, Mehrani H, Ghasemi A, Farrokhfall K. 2019. Cinnamaldehyde has beneficial effects against oxidative stress and nitric oxide metabolites in the brain of aged rats fed with long-term, high-fat diet. *J Funct Foods*, 52: 545-551.
- Berridge MJ, Bootman MD, Lipp P. 1998. Calcium-a life and death signal. *Nature*, 395: 645-648.
- Bhat RV, Budd Haeberlein SL, Avila J. 2004. Glycogen synthase kinase 3: a drug target for CNS therapies. *J Neurochem*, 89: 1313-1317.
- Bruno AM, Huang JY, Bennett DA, Marr RA, Hastings ML, Stutzmann GE. 2012. Altered ryanodine receptor expression in mild cognitive impairment and Alzheimer's disease. *Neurobiol Aging*, 33: 1001.e1001-e1006.
- Checler F. 1995. Processing of the β -amyloid precursor protein and its regulation in Alzheimer's disease. *J Neurochem*, 65: 1431-1444.
- Chen, Wang YW, Huang WS, Lee MM, Wood WG, Leung YM, Tsai HY. 2016. Trans-cinnamaldehyde, an essential oil in cinnamon powder, ameliorates cerebral ischemia-induced brain injury via inhibition of neuroinflammation through attenuation of iNOS, COX-2 expression and NF κ -B signaling pathway. *Neuromolecular Med*, 18: 322-333.
- Chen P, Gu Z, Liu W, Yan Z. 2007. Glycogen synthase kinase 3 regulates N-methyl-D-aspartate receptor channel trafficking and function in cortical neurons. *Mol Pharmacol*, 72: 40-51.
- Chui D-H, Tanahashi H, Ozawa K, Ikeda S, Checler F, Ueda O, Suzuki H, Araki W, Inoue H, Shirotani K. 1999. Transgenic mice with Alzheimer presenilin 1 mutations show accelerated neurodegeneration without amyloid plaque formation. *Nat Med*, 5: 560-564.
- Couturier K, Qin B, Batandier C, Awada M, Hininger-Favier I, Canini F, Leverve X, Roussel AM, Anderson RA. 2011. Cinnamon increases liver glycogen in an animal model of insulin resistance. *Metabolism*, 60: 1590-1597.
- Dall'Igna OP, Fett P, Gomes MW, Souza DO, Cunha RA, Lara DR. 2007. Caffeine and adenosine A_{2a} receptor antagonists prevent β -amyloid (25-35)-induced cognitive deficits in mice. *Exp Neurol*, 203: 241-245.
- Dall'Igna OP, Porciúncula LO, Souza DO, Cunha RA, Lara DR. 2003. Neuroprotection by caffeine and adenosine

- A2A receptor blockade of β -amyloid neurotoxicity. *Br J Pharmacol*, 138: 1207-1209.
- Danysz W, Parsons CG. 2012. Alzheimer's disease, β -amyloid, glutamate, NMDA receptors and memantine—searching for the connections. *Br J Pharmacol*, 167: 324-352.
- Delacourte A, Buée L. 2000. Tau pathology: a marker of neurodegenerative disorders. *Curr Opin Neurol*, 13: 371-376.
- Donley J, Hurt W, Stockert A. 2016. Effects of cinnamon extract on glycogen synthase kinase 3 phosphorylation of tau. *FASEB J*, 30: 814.815-814.815.
- Frydman-Marom A, Levin A, Farfara D, Benromano T, Scherzer-Attali R, Peled S, Vassar R, Segal D, Gazit E, Frenkel D. 2011. Orally administered cinnamon extract reduces β -amyloid oligomerization and corrects cognitive impairment in Alzheimer's disease animal models. *PLoS One*, 6: e16564.
- Gomada Y, Jamison BY, Maitin V, Vattam DA. 2012. Cinnamon decreases excitation toxicity in primary neuronal cultures from chick embryos. *FASEB J*, 26: 1011-1025.
- Gomes CV, Kaster MP, Tomé AR, Agostinho PM, Cunha RA. 2011. Adenosine receptors and brain diseases: neuroprotection and neurodegeneration. *Biochim Biophys Acta*, 1808: 1380-1399.
- Hernandez F, Lucas JJ, Avila J. 2013. GSK3 and tau: two convergence points in Alzheimer's disease. *J Alzheimers Dis*, 33: S141-S144.
- Ho S-C, Chang K-S, Chang P-W. 2013. Inhibition of neuroinflammation by cinnamon and its main components. *Food Chem*, 138: 2275-2282.
- Imparl-Radosevich J, Deas S, Polansky MM, Baedke DA, Ingebritsen TS, Anderson RA, Graves DJ. 1998. Regulation of PTP-1 and insulin receptor kinase by fractions from cinnamon: implications for cinnamon regulation of insulin signalling. *Horm Res Paediatr*, 50: 177-182.
- Keshavarz M, Farrokhi M, Amiri A. 2017. Caffeine neuroprotective mechanism against β -amyloid neurotoxicity in SHSY5Y cell line: involvement of adenosine, ryanodine, and N-methyl-D-aspartate receptors. *Adv Pharm Bull*, 7: 579-584.
- Klafke J, Da Silva M, Trevisan G, Rossato M, Da Silva C, Guerra G, Villarinho J, Rigo F, Dalmolin G, Gomez M. 2012. Involvement of the glutamatergic system in the nociception induced intrathecally for a TRPA1 agonist in rats. *Neuroscience*, 222: 136-146.
- Kwon B-M, Lee S-H, Cho Y-K, Bok S-H, So S-H, Youn M-R, Chang S-I. 1997. Synthesis and biological activity of cinnamaldehydes as angiogenesis inhibitors. *Bioorg Med Chem Lett*, 7: 2473-2476.
- LaFerla FM. 2002. Calcium dyshomeostasis and intracellular signalling in Alzheimer's disease. *Nat Rev Neurosci*, 3: 862-872.
- Lindeboom J, Weinstein H. 2004. Neuropsychology of cognitive ageing, minimal cognitive impairment, Alzheimer's disease, and vascular cognitive impairment. *Eur J Pharmacol*, 490: 83-86.
- Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. 1951. Protein measurement with the Folin phenol reagent. *J Biol Chem*, 193: 265-275.
- Lüscher C, Malenka RC. 2012. NMDA receptor-dependent long-term potentiation and long-term depression (LTP/LTD). *Cold Spring Harb Perspect Biol*, 4: a005710.
- Lv C, Yuan X, Zeng H-W, Liu R-H, Zhang W-D. 2017. Protective effect of cinnamaldehyde against glutamate-induced oxidative stress and apoptosis in PC12 cells. *Eur J Pharmacol*, 815: 487-494.
- McPherson PS, Kim Y-K, Valdivia H, Knudson CM, Takekura H, Franzini-Armstrong C, Coronadot R, Campbell KP. 1991. The brain ryanodine receptor: a caffeine-sensitive calcium release channel. *Neuron*, 7: 17-25.
- Peng J, Liang G, Inan S, Wu Z, Joseph DJ, Meng Q, Peng Y, Eckenhoff MF, Wei H. 2012. Dantrolene ameliorates cognitive decline and neuropathology in Alzheimer triple transgenic mice. *Neurosci Lett*, 516: 274-279.
- Peterson DW, George RC, Scaramozzino F, LaPointe NE, Anderson RA, Graves DJ, Lew J. 2009. Cinnamon extract inhibits tau aggregation associated with Alzheimer's disease in vitro. *J Alzheimers Dis*, 17: 585-597.
- Phiel CJ, Wilson CA, Lee VM-Y, Klein PS. 2003. GSK-3 α regulates production of

- Alzheimer's disease amyloid- β peptides. *Nature*, 423: 435.
- Sartorius T, Peter A, Schulz N, Drescher A, Bergheim I, Machann J, Schick F, Siegel-Axel D, Schürmann A, Weigert C, Häring H-U, Hennige AM. 2014. Cinnamon extract improves insulin sensitivity in the brain and lowers liver fat in mouse models of obesity. *PLoS One*, 9: e92358.
- Shimada Y, Goto H, Kogure T, Kohta K, Shintani T, Itoh T, Terasawa K. 2000. Extract prepared from the bark of *Cinnamomum cassia* Blume prevents glutamate-induced neuronal death in cultured cerebellar granule cells. *Phytother Res*, 14: 466-468.
- Stavinoha RC, Vattem DA. 2015. Potential neuroprotective effects of cinnamon. *Int J Appl Res Nat Prod* 8: 24-46.
- Supnet C, Bezprozvanny I. 2010. The dysregulation of intracellular calcium in Alzheimer disease. *Cell Calcium*, 47: 183-189.
- Szatmari E, Habas A, Yang P, Zheng J-J, Hagg T, Hetman M. 2005. A positive feedback loop between glycogen synthase kinase 3 β and protein phosphatase 1 after stimulation of NR2B NMDA receptors in forebrain neurons. *J Biol Chem*, 280: 37526-37535.
- Takashima A. 2006. GSK-3 is essential in the pathogenesis of Alzheimer's disease. *J Alzheimers Dis*, 9: 309-317.
- Zhang H, Sun S, Herreman A, De Strooper B, Bezprozvanny I. 2010. Role of presenilins in neuronal calcium homeostasis. *J Neurosci*, 30: 8566-8580.
- Zhao J, Zhang X, Dong L, Wen Y, Zheng X, Zhang C, Chen R, Zhang Y, Li Y, He T. 2015. Cinnamaldehyde inhibits inflammation and brain damage in a mouse model of permanent cerebral ischaemia. *Br J Pharmacol*, 172: 5009-5023.