

Original Research Article

Protective effects of quercetin against hyperglycemia-induced oxidative stress in hepatic HepG2 cell line

Amir Yarahmadi^{1,2}, Mostafa Moradi Sarabi³, Ahmad Sayahi³, Fatemeh Zal^{1,4,*}

¹Department of Biochemistry, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

²Transplant Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

³Department of Biochemistry and Genetics, Lorestan University of Medical Sciences, Faculty of Medicine, Khorramabad, Iran

⁴Research Center for Traditional Medicine and History of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

Article history:

Received: Oct 29, 2019

Received in revised form:

Aug 10, 2020

Accepted: Aug 11, 2020

Vol. 11, No. 3, May-Jun 2021,
269-280.

* Corresponding Author:

Tel: +98 713 234 3848

Fax: +98 713 234 3848

Fatemehzal@Yahoo.Com

Keywords:

Flavonoid

Quercetin

Antioxidant enzymes

Hyperglycemia

Oxidative stress

HepG2 cells

Abstract

Objective: Hyperglycemia is a severe consequence of diabetes mellitus (DM). Through induction of oxidative stress, it plays a major role in the pathogenesis of several complications in DM. Therefore, new strategies and antioxidants should be implemented in the treatment of DM. Quercetin is a flavonoid with strong antioxidant capacity found dominantly in vegetables, fruits, leaves, and grains. The current study aimed to investigate quercetin protective effects under D-glucose-induced oxidative stress by assessing antioxidant defense enzymes in HepG2 cells as an *in vitro* model.

Materials and Methods: HepG2 cells were cultured with different concentrations of D-glucose (5.5, 30 and 50 mM) and/or 25 μ M quercetin for 48 and 72 hr, respectively. The effect of treatments on cellular integrity, antioxidant enzymes superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GPx), glutathione reductase (GR) activity, and cellular levels of glutathione (GSH) and malondialdehyde (MDA) was determined.

Results: D-glucose had various effects on intracellular antioxidant defense at different doses and time-points and quercetin could attenuate oxidative stress and modulate antioxidant defenses.

Conclusion: The results of this study indicated that flavonoid quercetin could be proposed as an agent protecting hepatic HepG2 cells against oxidative stress associated with hyperglycemia.

Please cite this paper as:

Yarahmadi A, Moradi Sarabi M, Sayahi A, Zal F. Protective effects of quercetin against hyperglycemia-induced oxidative stress in hepatic HepG2 cell line. Avicenna J Phytomed, 2021; 11(3): 269-280.

Introduction

Hyperglycemia-induced oxidative damage is one of the important complications in the pathophysiology of type-2 diabetes mellitus

(T2DM) (American Diabetes, 2014; Roden, 2016). Hyperglycemia can play an important role in pathophysiological changes that happen during DM through induction of oxidant formation, advanced glycation end products (AGE) generation,

hyperosmolarity associated with high glucose, abnormality of sorbitol and myoinositol metabolism, and protein kinase C activation (Evcimen and King, 2007; Kalousova et al., 2002). Mechanistically, hyperglycemia generates reactive oxygen species (ROS) that have a pivotal role in D-glucose-induced cellular impairment (Chandrasekaran et al., 2010; Zal et al., 2014a). ROS consist of superoxide anion radical ($O_2^{\cdot-}$), hydrogen peroxide (H_2O_2), hydroxyl radical (OH^{\cdot}), and singlet oxygen (1O_2), which are the most reactive oxygen species that damage cells during oxidative insult (Hamidi Alamdari et al., 2020; Nimse and Pal, 2015). Antioxidants that neutralize ROS to prevent cellular components from oxidative insult, can be divided into enzymatic and non-enzymatic antioxidants. Enzymatic antioxidants, including superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GPx) and glutathione reductase (GR) and non-enzymatic antioxidants (e. g. GSH, vitamin A, C, and E, and bioflavonoids) have been implicated in the protection against ROS (Di Luzio and Hartman, 1967; Kazemi et al., 2015; Yazdi et al., 2020). The enzyme SOD is located in both cytosol and mitochondria and catalyzes conversion of $O_2^{\cdot-}$ into O_2 and H_2O_2 by using metal cofactors copper (Cu), zinc (Zn), and manganese (Mn), respectively. In the peroxisome of cells, the enzyme CAT converts H_2O_2 to H_2O and O_2 . GPx is another enzyme responsible for dissociation of H_2O_2 into H_2O via GSH in both cytoplasm and extracellular fluid of all human tissues. GR functions by reducing oxidized GS-SG into its active form GSH which is necessary for the action of GPx (Nimse and Pal, 2015). Antioxidants tightly regulate the cellular redox balance and can delay or inhibit oxidative damage by converting free radicals to non-toxic products (Amin et al., 2020; Mashhoody et al., 2014; Weydert and Cullen, 2010). Flavonoids are employed as the most attractive agents among the antioxidants

(Lodhi et al., 2016; Saboonchian et al., 2014). They have different actions including antiallergic, anti-inflammatory, antiseptic, and antioxidant, and exert limited toxicity (Choi et al., 2017; Srivastava et al., 2016). However, potent antioxidant activities of flavonoids are due to the high redox potential that allows them to act as reducing agents for reducing and scavenging free radicals (Anand David et al., 2016; Razali et al., 2015; Srivastava et al., 2016). Quercetin, primarily known for its prominent antioxidant capacity, is one of the most abundant flavonoids widely found in various food products, fruits, and vegetables (Ader et al., 2000; Mulholland et al., 2001). Quercetin has been investigated for its beneficial effects such as the prevention of cell death and oxidant damage (Yarahmadi et al., 2017; Zizkova et al., 2017). It is also famous for protecting DNA from oxidative damages of $O_2^{\cdot-}$, H_2O_2 , and OH^{\cdot} on DNA oligonucleotides. It has been suggested that quercetin has a beneficial impact on diabetes by reducing hyperglycemia-induced oxidative stress in hepatic tissue (Chis et al., 2016; Yarahmadi et al., 2018). Moreover, many studies have indicated that the antioxidant capacity of quercetin is stronger than other phenolic compounds (Bai et al., 2013). Given that hyperglycemia injures the cells through oxidative stress in diabetes and considering the role of quercetin as a potent antioxidant, the present study aimed to investigate potential protective effects of quercetin against oxidative stress induced by D-glucose at different doses and time-points in HepG2 cells *in vitro*.

Materials and Methods

Chemicals

Quercetin and D-glucose were purchased from Sigma Chemical Co. (St. Louis, MO, USA); Roswell Park Memorial Institute Medium (RPMI), fetal bovine serum (FBS), penicillin, and streptomycin were obtained from Gibco-BRL (Paisley, UK). BSA total protein assay kit was

purchased from Bio-Rad (Hercules, California, USA). Trypsin was from BDH, England. Potassium chloride, and potassium dihydrogen orthophosphate were from Fluka, England. GR, tert-butyl hydroperoxide (t-BuOOH), bovine serum albumin (BSA), nicotinamide adenine dinucleotide phosphate (NADPH), Triton X-100, ethylenediaminetetraacetic acid (EDTA), and dimethyl sulfoxide (DMSO) were purchased from Sigma Chemical Co. (Poole, Dorset, UK).

Cell culture and treatment

The human HepG2 cell line was acquired from the National Cell Bank of Iran (NCBI, Pasteur Institute, Tehran). Then, cells were cultured in RPMI 1640 medium containing 10% heat-inactivated FBS, supplemented with 100 IU/ml penicillin and 100 µg/ml streptomycin in a humidified atmosphere containing 5% CO₂ at 37°C. All experiments were done using cells at 70% confluences (Zal et al., 2020). Cells were then grown and divided into 6 groups as follows: the first group included media containing 5.5 mM D-glucose (as a normal concentration and served as normal control cells), the second group included D-glucose 30 mM, the third group included D-glucose 50 mM (as the high concentration), and the fourth group was treated with 25 µM quercetin, the fifth group included D-glucose 30 mM + quercetin (25 µM) and finally, the sixth group was treated with D-glucose 50 mM + quercetin (25 µM). The vehicle for the quercetin was ethanol, and a separate group was treated with ethanol (0.3%), and since it did not show significant differences with control, was omitted from analysis. We performed all treatments in quadruplicate for each group. Treated cells were incubated at 37°C with 5% CO₂. The cell lysate was collected after 48 and 72 hr treatment and stored at -20°C until use. Notably, the pH of the culture medium was not affected by addition of D-glucose or quercetin.

Cell viability assay

To assess the viability of cells treated with different concentrations of D-glucose or quercetin, an assay was carried out using 3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyltetrazolium bromide (MTT) as previously described by Mosmann (Mosmann, 1983). First, 12×10³ cells/well in a 96-well plate were incubated with 5.5, 30 and 50 mM D-glucose and/or 25 µM quercetin for 48 and 72 hr, respectively at 37°C. Then, cells were incubated with MTT (0.5 mg/ml) dissolved in serum-free medium. After 3.5 hr incubation, 100 µl DMSO was added to dissolve the formazan crystals, and then, absorbance was determined at 570/650 nm wavelength using an ELISA reader (Bio Rad, USA). Cell viability was determined as the ratio of absorbance of treated cells to that of untreated cells that served as a control.

Measurement of Cu, Zn-SOD activity

We measured the activity of Cu, Zn-SOD in HepG-2 cell lysate based on the procedure that was described by Misra and Fridovich (Misra and Fridovich, 1972). This method relies on assessing inhibition of adrenaline auto-oxidation to adrenochrome by enzyme SOD in alkaline pH 10.2 at 480 nm. First, we drew the standard curve of SOD by plotting the amount of inhibition of adrenaline oxidation against different concentrations of SOD. Each unit of SOD enzyme is the amount of enzyme that inhibits 50% adrenaline oxidation, and SOD enzyme activity is reported as unit/mg protein of the HepG2 cell lysate.

Measurement of CAT activity

For determination of CAT activity, we used a method that was previously described by Aebi (Aebi, 1984). In this method, we spectrophotometrically assayed decomposition of H₂O₂ to H₂O and O₂ by CAT. Enzyme activity is shown as mmol H₂O₂ consumed/min per mg HepG2 cell lysate protein by a molar absorptivity of 43.6 M/cm.

Measurement of GPx activity

GPx activity was measured by the process described by Fecondo and Augustey based on monitoring continuous substitution of reduced GSH from oxidized form (G-S-S-G) in the presence of enzyme glutathione reductase. Besides, we used disodium (Na_2) salt of reduced nicotinamide adenine dinucleotide phosphate for the assessment of GPx activity with minor modifications (Mostafavi-Pour et al., 2008). GPx enzyme activity in the HepG2 cell lysate is shown as μmol of NADPH oxidized/min/mg of cell protein using a molar absorptivity of $6.22 \times 10^6 \text{ M/cm}$ for NADPH. One unit of GPx is shown as U/mg of HepG2 cell protein.

Measurement of GR activity

The enzyme GR activities were measured using the method of Racker and Carlberg (Carlberg and Mannervik, 1985; Racker, 1955). For determination of GR activity, 60 μM buffer, 5 mM EDTA (pH 8.0), 0.033 M GS-SG, 2 mM NADPH, and a sample of HepG2 cell lysate at a final volume of 1000 μl , were used. A reduction in absorbance shows oxidation of NADPH through reduction of GS-SG by the activity of enzyme GR in the HepG2 cell lysate which was monitored spectrophotometrically at 340 nm wavelength for 3 min. Finally, the results are reported through a molar absorptivity of $6.22 \times 10^6 \text{ M/cm}$ for NADPH. One unit of GR is defined as U/mg HepG2 cell protein.

Determination of reduced GSH

Assessment of reduced GSH using 5, 5-dithio-bis (2-nitrobenzoic) acid (DTNB), was performed using Ellman's method with some modifications (Mashhoody et al., 2014; Zal et al., 2014b). We used 1 mM solutions of reduced GSH to draw a standard curve. GSH amount was assayed in the HepG2 cell lysate. For measurement of GSH level, 0.5 ml of DTNB (0.001 M) solution was added to 2.3 ml potassium phosphate buffer (0.2 M, pH 7.6) plus 0.2

ml of HepG2 cell lysate. The absorbance of the solution was read after 5 min at 412 nm.

Determination of MDA

MDA as a marker of lipid peroxidation was estimated according to a colorimetric method. Briefly, 100 μl of the HepG2 cell lysate was added to 400 μl TBA reagent containing 0.375% TBA, 15% trichloroacetic acid, and 0.25 mol/HCl. The mixture was boiled in a water bath at 95°C for 30 min and after fast cooling, it was centrifuged at $8000 \times g$ for 15 min at 4°C. The absorbance of the pink color supernatant was measured at 532 nm. MDA concentration was calculated using tetraethoxypropane (TEP) as standard and expressed as nmol/mg protein.

Measurement of total protein

Protein content was determined using BSA as a standard with BSA total protein assay kit from Bio-Rad according to the manufacturer protocols.

Statistical analysis

GraphPad Prism version 6.01 (GraphPad Software, San Diego, CA, USA) and SPSS18 software (SPSS, Chicago, IL, USA) were used for group comparison. The data are presented as mean \pm SD and analyzed using one-way analysis of variance (one-way ANOVA) followed by Tukey's multiple comparison test. A p -value ≤ 0.05 was used as the level of significance. Mean \pm SD values are the representative of four independent experiments.

Results

Effect of D-glucose and quercetin on the viability of HepG2 cells

Figure 1 (A and B) shows that viability was not affected when HepG2 cells were incubated with 30 and 50 mM D-glucose with or without quercetin for 48 and 72 hr, respectively (85-90% viability).




Figure 1. Effect of D-glucose and/ or quercetin on the viability of HepG2 cells after 48 (A) and 72 (B) hr treatment using the MTT assay. Data are presented as mean±SD (n=4). All data are presented as a percentage with respect to control (100% cell viability). Glu: D-glucose; Q: Quercetin; and MTT: 3-(4, 5-dimethylthiazol-2-yl)-2, 5 diphenyltetrazolium bromide.

Effect of D-glucose and quercetin on SOD activity

The effect of D-glucose on SOD activity was different at different concentrations and time-points. As shown in Figure 2, after both 48 and 72 hr, 30 mM D-glucose significantly increased SOD activity, but 50 mM D-glucose decreased it though it was significant only after 72 hr of treatment with 50 mM D-glucose ($p < 0.05$). The combinations of quercetin and 30 mM D-glucose significantly decreased the level of SOD activity compared to the corresponding values in HepG2 cells treated with 30 mM D-glucose, while

combinations of quercetin and 50 mM D-glucose significantly increased the level of SOD activity compared to the corresponding values at 50 mM D-glucose concentration ($p < 0.05$).




Figure 2. Effect of D-glucose and/ or quercetin on SOD activity in HepG2 cells after 48 (A) and 72 (B) hr treatment. Results are given as mean±SD (n=4). *shows a significant difference compared to the respective control group ($p < 0.05$); Φ shows a significant difference compared to 30 mM D-glucose-treated group ($p < 0.05$); Θ shows a significant difference compared to 50 mM D-glucose-treated group ($p < 0.05$). Glu: D-glucose; Q: Quercetin; and SOD: Superoxide dismutase.

Effect of D-glucose and quercetin on CAT activity

As shown in Figure 3, after 48 hr, a 2.2 fold increase in CAT activity was observed in HepG2 cells treated with 30 mM D-glucose as compared to normal control cells, but after 72 hr, 44.7% decrease in CAT activity in 30 mM D-glucose-treated group occurred ($p < 0.05$). Also, 50 mM D-

glucose significantly decreased CAT activity by about 69% after 72 hr as compared to the normal control cells ($p < 0.05$). At both 48 and 72 hr time-points, quercetin could improve these conditions compared to the corresponding values in HepG2 cells treated with 30 and 50 mM D-glucose, respectively ($p < 0.05$). Quercetin-treated cells showed an increase in CAT activity, which was significant only after 48 hr of treatment as compared to the normal control cells ($p < 0.05$).

Effect of D-glucose and quercetin on GPx activity

Our data showed that treatment of HepG2 cells with 30 and 50 mM D-glucose for 48 hr resulted in significant 33 and 38% decreases in GPx activity, respectively ($p < 0.05$) (Figure 4). However, after 72 hr, this reduction was 40 and 75.7% as compared to the control cells, respectively ($p < 0.05$) (Figure 4). Furthermore, incubation of HepG2 cells with quercetin and quercetin plus 30 and 50 mM D-glucose caused an increase in GPx activity as compared to the control group, but this increase was significant only at 30 mM D-glucose + quercetin after 72 hr.

Effect of D-glucose and quercetin on GR activity

As shown in Figure 5, after 48 hr, we found no significant differences in GR activity in D-glucose-treated HepG2 cells with or without quercetin as compared to the control group. However, after 72 hr, 30 and 50 mM D-glucose significantly decreased GR activity, and quercetin could not overcome this reduction ($p < 0.05$) (Figure 5).

Effect of D-glucose and quercetin on GSH level in HepG2 cells

Figure 6 shows that the level of GSH was significantly decreased by 41% in HepG2 cells treated with 50 mM D-glucose after 48 hr, as compared to the control group ($p < 0.05$).




Figure 3. Effect of D-glucose and/ or quercetin on CAT activity in HepG2 cells after 48 (A) and 72 (B) hr treatment. Results are given as mean±SD (n=4). * shows asignificant difference compared to the respective control group ($p < 0.05$); Φ shows asignificant difference compared to 30 mM D-glucose treated group ($p < 0.05$); Θ shows asignificant difference compared to 50 mM D-glucose treated group ($p < 0.05$). Glu: D-glucose; Q: Quercetin; and CAT: Catalase.

Also, Figure 6 demonstrates that after 72 hr, 30 and 50 mM D-glucose significantly decreased GSH by 34.4 and 55.5%, respectively, in treated HepG2 cells in comparison to the control group ($p < 0.05$).

After 48 hr incubation of HepG2 cells treated with quercetin plus 50 mM D-glucose, a significant increase in GSH level occurred compared to 50 mM D-glucose-treated cells ($p < 0.05$).

Additionally, our data indicated that after 72 hr, incubation of HepG2 cells with quercetin plus 30 and 50 mM D-glucose

caused a significant increase in GSH level compared to the corresponding 30 and 50 mM D-glucose treated HepG2 treated cells, respectively ($p < 0.05$) (Figure 6).




Figure 4. Effect of D-glucose and/ or quercetin on GPx activity in HepG2 cells after 48 (A) and 72 (B) hr treatment. Results are given as mean \pm SD (n=4). *shows asignificant difference compared to the respective control group ($p < 0.05$); Φ shows a significant difference compared to 30 mM D-glucose treated group ($p < 0.05$).Glu: D-glucose; Q: Quercetin; and GPx: Glutathione peroxidase.

Effect of D-glucose and quercetin on MDA level in HepG2 cells

As shown in Figure 7, 48 hr treatment of HepG2 cells with 30 and 50 mM D-glucose resulted in 40 and 56% increases in the level of MDA as a marker of lipid peroxidation compared to the control group ($p < 0.05$).

Moreover, quercetin produced a significant decrease in the level of MDA in cells treated with 30 and 50 mM D-glucose ($p < 0.05$).

Our results also showed that 72hr treatment of HepG2 cells with 50 mM D-glucose significantly increased the level of MDA to about 41%, and quercetin significantly decreased the MDA level in 50 mM D-glucose-treated cells ($p < 0.05$) (Figure 7).




Figure 5. Effect of D-glucose and/ or quercetin on GR activity in HepG2 cells after 48 (A) and 72 (B) hr treatment. Results are given as mean \pm SD (n=4). * shows a significant difference compared to the respective control group ($p < 0.05$).Glu: D-glucose; Q: Quercetin; and GR: Glutathione reductase.




Figure 6. Effect of D-glucose and/ or quercetin on GSH level in HepG2 cells after 48 (A) and 72 (B) hr treatment. Results are given as mean±SD (n=4). * shows asignificant difference compared to the respective control group (p<0.05); Φ shows asignificant difference compared to 30 mM D-glucose treated group (p<0.05); Θ shows asignificant difference compared to 50 mM D-glucose treated group (p<0.05). Glu: D-glucose; Q: Quercetin; and GSH: Glutathione.

Discussion

The current study aimed to investigate the effects of quercetin on hyperglycemia and hyperglycemia-induced disruption of antioxidant defense system in HepG2 cells. There are several studies indicating that HepG2 cells are a well-characterized cell line to study hyperglycemia *in vitro* (Chandrasekaran et al., 2010a; Alia et al., 2006).

Chandrasekaran et al. used 30 and 50 mM D-glucose for 72 hr to show toxic effects of hyperglycemia on HepG2 cells (Chandrasekaran et al., 2010). Similarly, Cordero-Herrera et al. used different D-glucose concentrations ranging from 20-60 mM for 24 hr, to assess hyperglycemia effects in human HepG2 cells (Cordero-Herrera et al., 2014). Moreover, Buranasin et al. used 25 and 50 mM D-glucose for 72 hr to create hyperglycemic conditions in human gingival fibroblasts (Buranasin et al., 2018).




Figure 7. Effect of D-glucose and/ or quercetin on MDA level in HepG2 cells after 48 (A) and 72 (B) hr treatment. Results are given as mean±SD (n=4). * shows asignificant difference compared to the respective control group (p<0.05); Φ shows asignificant difference compared to 30 mM D-glucose treated group (p<0.05); Θ shows asignificant difference compared to 50 mM D-glucose-treated group (p<0.05). Glu: D-glucose; Q: Quercetin; and MDA: Malondialdehyde.

Our results showed that D-glucose at different concentrations and time-points had different effects on cellular antioxidant enzyme patterns, which may be explained by this fact that the amount and time of the exposure to produce oxidative conditions may influence the antioxidant enzyme defense capacity. Oxidative stress resulted from higher concentrations and longer time periods of D-glucose exposure (50 mM for 72 hr) may overcome the antioxidant enzyme defenses capacity in HepG2 cells compared to the cells treated with 30 mM D-glucose for 48 hr.

ROS are generated at low quantities during normal cell metabolism. However, ROS overproduction has been observed in many diseases, such as diabetes and many types of cancers (Matsuda and Shimomura, 2013). A high level of D-glucose promotes generation of ROS during diabetes that plays a crucial role in D-glucose-induced cellular dysfunction and oxidative stress (Chandrasekaran et al., 2010b). Our data demonstrated that treatment of HepG2 cells with D-glucose evoked a notable increase in SOD activity, and quercetin significantly counterbalanced it, which is compatible with the study of Ceriello et al. showing that hyperglycemia increased the activity and mRNA expression of Cu, Zn-SOD and Mn-SOD in human endothelial cells (Ceriello et al., 1996). Similarly, Weidig et al. showed that incubation of rat coronary endothelial cells with 22 mM D-glucose significantly increased Cu, Zn-SOD activity and free radical scavenger mercaptopropionylglycine reduced it to that of the control group (Weidig et al., 2004). Our results showed a significant decrease in SOD activity after 72 hr of treatments with 50 mM D-glucose, but quercetin significantly raised it. This is compatible with Uchimura et al. study where they showed significantly lower levels of Cu, Zn-SOD and Mn-SOD activities in neutrophils and lymphocytes of patients with diabetes mellitus compared to healthy controls, which can be explained by this theory that adaptive or toxicity responses

may depend on time and concentration of exposure (Uchimura et al., 1999). In our experiment, the activity of CAT significantly increased after 48 hr exposure to 30 mM D-glucose and quercetin significantly reduced this increase compared to the corresponding value. However, after 72 hr treatment, CAT activity significantly decreased in both 30 and 50 mM D-glucose-treated cells. These results are compatible with the Dave et al. study in which hyperglycemia decreased CAT and GPx activity in both types of diabetes (Dave and Kalia, 2007). Consistent to our study, Alam et al. showed that CAT, SOD, glutathione-S-transferase and GSH levels were significantly increased in type-2 diabetic mice treated with quercetin for three weeks compared to normal controls (Alam et al., 2014). The current study demonstrated that treatment of HepG2 cells with different concentrations of D-glucose caused a significant reduction in GPx activity after both 48 and 72 hr respectively, and quercetin increased GPx activity, but it was significant only in HepG2 cells that were treated with 30 mM D-glucose plus quercetin for 72 hr. These data are compatible with the study conducted by Amaral et al. They showed that hyperglycemia reduces GPx activity in testicular cells isolated from diabetic rats (Amaral et al., 2006). Moreover, Jeon et al. showed that consumption of quercetin by type 2 diabetic *db/db* mice, increases GPx activity in the liver after six weeks of treatment (Jeon et al., 2012).

Furthermore, our results showed that GR enzyme activity and levels of GSH decreased in D-glucose-treated HepG2 cells, especially in a concentration- and time-dependent manner. It may be primarily due to intracellular oxidation and increase of oxidative stress in cellular environments (Chandrasekaran et al., 2010b; Hall, 1999). Moreover, we observed that quercetin treatment of HepG2 cells prevents D-glucose-induced oxidation effects; our results were comparable to those of other studies which demonstrated

that quercetin exhibits a preventative effect against hyperglycemia-induced mitochondrial impairment in liver cells (Zou et al., 2014).

In conclusion, quercetin can attenuate oxidative stress caused by hyperglycemia and protect human HepG2 cells by modulating antioxidant defense system. However, further pharmacological studies are required to establish its beneficial effects in patients with disrupted oxidative balance such as diabetes mellitus.

Acknowledgment

The current study was supported by a grant (number 16110) from the office of Vice Chancellor for Research at Shiraz University of Medical Sciences, Shiraz, Iran.

Conflicts of interest

The authors have declared that there is no conflict of interest.

References

- Ader P, Wessmann A, Wolfram S. 2000. Bioavailability and metabolism of the flavonol quercetin in the pig. *Free Radic Biol Med*, 28: 1056-1067.
- Aebi H. 1984. Catalase in vitro. *Methods Enzymol*, 105:121-126.
- American Diabetes A. 2014. Standards of medical care in diabetes. *Diabetes Care*, 1: 14-80.
- Amin F, Roohbakhsh A, Memarzia A, Kazerani H, Boskabady MH. 2020. Paraquat-induced systemic inflammation and increased oxidative markers in rats improved by *Zataria multiflora* extract and carvacrol. *Avicenna J Phytomed*, 10:513.
- Anand David AV, Arulmoli R, Parasuraman S. 2016. Overviews of Biological Importance of Quercetin: A Bioactive Flavonoid. *Pharmacogn Rev*, 10: 84-89.
- Bai X, Zhang H, Ren S. 2013. Antioxidant activity and HPLC analysis of polyphenol-enriched extracts from industrial apple pomace. *J Sci Food Agric*, 93: 2502-2506.
- Buranasin P, Mizutani K, Iwasaki K, et al. 2018. High glucose-induced oxidative stress impairs proliferation and migration of human gingival fibroblasts. *PloS one*, 13: 0201855.
- Carlberg I, Mannervik B. 1985. Glutathione reductase. *Methods Enzymol*, 113: 484-490.
- Ceriello A, dello Russo P, Amstad P, Cerutti P. 1996. High glucose induces antioxidant enzymes in human endothelial cells in culture: evidence linking hyperglycemia and oxidative stress. *Diabetes*, 45: 471-477.
- Chandrasekaran K, Swaminathan K, Chatterjee S, Dey A. 2010. Apoptosis in HepG2 cells exposed to high glucose. *Toxicol In Vitro*, 24: 387-396.
- Chis IC, Muresan A, Oros A, Nagy AL, Clichici S. 2016. Protective effects of Quercetin and chronic moderate exercise (training) against oxidative stress in the liver tissue of streptozotocin-induced diabetic rats. *Physiol Int*, 103: 49-64.
- Choi KC, Son YO, Hwang JM, Kim BT, Chae M, Lee JC. 2017. Antioxidant, anti-inflammatory and anti-septic potential of phenolic acids and flavonoid fractions isolated from *Lolium multiflorum*. *Pharm Biol*, 55: 611-619.
- Cordero-Herrera I, Martín MÁ, Goya L, Ramos S. 2014. Cocoa flavonoids attenuate high glucose-induced insulin signalling blockade and modulate glucose uptake and production in human HepG2 cells. *Food Chem Toxicol*, 64: 10-19.
- Dave GS, Kalia K. 2007. Hyperglycemia induced oxidative stress in type-1 and type-2 diabetic patients with and without nephropathy. *Cell Mol Biol*, 53: 68-78.
- Di Luzio NR, Hartman AD. 1967. Role of lipid peroxidation in the pathogenesis of the ethanol-induced fatty liver. *Fed Proc*, 26: 1436-1442.
- Evcimen ND, King GL. 2007. The role of protein kinase C activation and the vascular complications of diabetes. *Pharmacol Res*, 55: 498-510.
- Hamidi Alamdari D, Bagheri Moghaddam A, Amini S, Hamidi Alamdari A, Damsaz M, Yarahmadi A. 2020. The Application of a Reduced Dye Used in Orthopedics as a Novel Treatment against Coronavirus (COVID-19): A Suggested Therapeutic Protocol. *Arch Bone Jt Surg*, 8: 291-294.
- Kalousova M, Skrha J, Zima T. 2002. Advanced glycation end-products and advanced oxidation protein products in patients with diabetes mellitus. *Physiol Res*,

- 51: 597-604.
- Kazemi E, Mortazavi SM, Ali-Ghanbari A, Sharifzadeh S, Ranjbaran R, Mostafavi-pour Z, Zal F, Haghani M. 2015. Effect of 900 MHz Electromagnetic Radiation on the Induction of ROS in Human Peripheral Blood Mononuclear Cells. *J Biomed Phys Eng*, 5: 105-114.
- Lodhi S, Jain A, Jain AP, Pawar RS, Singhai AK. 2016. Effects of flavonoids from *Martynia annua* and *Tephrosia purpurea* on cutaneous wound healing. *Avicenna J Phytomed*, 6: 578.
- Mashhoody T, Rastegar K, Zal F. 2014. Perindopril may improve the hippocampal reduced glutathione content in rats. *Adv Pharm Bull*, 4: 155-159.
- Misra HP, Fridovich I. 1972. The role of superoxide anion in the autoxidation of epinephrine and a simple assay for superoxide dismutase. *J Biol Chem*, 247: 3170-3175.
- Mosmann T. 1983. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods*, 65: 55-63.
- Mostafavi-Pour Z, Zal F, Monabati A, Vessal M. 2008. Protective effects of a combination of quercetin and vitamin E against cyclosporine A-induced oxidative stress and hepatotoxicity in rats. *Hepatol Res*, 38: 385-392.
- Mulholland PJ, Ferry DR, Anderson D, Hussain SA, Young AM, Cook JE, Hodgkin E, Seymour LW, Kerr DJ. 2001. Pre-clinical and clinical study of QC12, a water-soluble, pro-drug of quercetin. *Ann Oncol*, 12: 245-248.
- Nimse SB, Pal D. 2015. Free radicals, natural antioxidants, and their reaction mechanisms. *RSC Adv*, 5: 27986-28006.
- Racker E. 1955. Glutathione reductase from bakers' yeast and beef liver. *J Biol Chem*, 217: 855-865.
- Razali N, Mat Junit S, Ariffin A, Ramli NS, Abdul Aziz A. 2015. Polyphenols from the extract and fraction of *T. indica* seeds protected HepG2 cells against oxidative stress. *BMC Complement Altern Med*, 15: 438.
- Roden M. 2016. Diabetes mellitus: definition, classification and diagnosis. *Wien Klin Wochenschr*, 2: 37-40.
- Saboonchian F, Jamei R, Sarghein SH. 2014. Phenolic and flavonoid content of *Elaeagnus angustifolia* L.(leaf and flower). *Avicenna J Phytomed*, 4: 231.
- Srivastava S, Somasagara RR, Hegde M, Nishana M, Tadi SK, Srivastava M, Choudhary B, Raghavan SC. 2016. Quercetin, a Natural Flavonoid Interacts with DNA, Arrests Cell Cycle and Causes Tumor Regression by Activating Mitochondrial Pathway of Apoptosis. *Sci Rep*, 6: 24049.
- Weidig P, McMaster D, Bayraktutan U. 2004. High glucose mediates pro-oxidant and antioxidant enzyme activities in coronary endothelial cells. *Diabetes Obes Metab*, 6: 432-441.
- Weydert CJ, Cullen JJ. 2010. Measurement of superoxide dismutase, catalase and glutathione peroxidase in cultured cells and tissue. *Nat Protoc*, 5: 51-66.
- Yarahmadi A, Khademi F, Mostafavi-Pour Z, Zal F. 2018. In-vitro analysis of glucose and quercetin effects on m-TOR and Nrf-2 Expression in HepG2 cell line (diabetes and cancer connection). *Nutr Cancer*, 70: 770-775.
- Yarahmadi A, Zal F, Bolouki A. 2017. Protective effects of quercetin on nicotine induced oxidative stress in 'HepG2 cells'. *Toxicol Mech Methods*, 27: 609-614.
- Bagheri Yazdi H, Hojati V, Shiravi A, Hosseinian S, Vaezi G. 2020. The role of *Artemisia turanica* extract on renal oxidative and biochemical markers in STZ-induced diabetes in rat. *Avicenna J Phytomed*, "In press"
- Zal F, Mahdian Z, Zare R, Soghra B, Mostafavi-Pour Z. 2014a. Combination of vitamin E and folic acid ameliorate oxidative stress and apoptosis in diabetic rat uterus. *Int J Vitam Nutr Res*, 84: 55-64.
- Zal F, Taheri R, Khademi F, Keshavarz E, Rajabi S, Mostafavi-Pour Z. 2014b. The combined effect of furosemide and propranolol on GSH homeostasis in ACHN renal cells. *Toxicol Mech Methods*, 24: 412-416.
- Zal F, Yarahmadi A, Totonchi H, Barazesh M, Sarabi MM. 2020. Nicotine attenuates global genomic DNA methylation by influencing DNMTs gene expression in human endometrial stromal cells. *Genes Environ*, 42: 1-8.
- Zizkova P, Stefek M, Rackova L, Prnova M, Horakova L. 2017. Novel quercetin derivatives: From redox properties to

promising treatment of oxidative stress
related diseases. Chem Biol Interact, 265: 36-46.