

Prevalence of *cdt* Gene in *Campylobacter jejuni* Strains Isolated from Surface Waters of Rasht, Iran

Keyvan Roshanjo (MSc)

Department of Microbiology,
Rasht Branch, Islamic Azad
University, Rasht, Iran

Leila Asadpour (MSc)

Department of Microbiology,
Rasht Branch, Islamic Azad
University, Rasht, Iran

Mohammad Reza Shiri

Shahsavari (MSc)
Department of Cellular and
Molecular Nutrition, Faculty of
Nutritional Sciences and Dietetics,
Tehran University of Medical
Sciences, Tehran, Iran

Arash Hemmati (MSc)

Department of Microbiology,
Islamic Azad University, Rasht
Branch, Rasht, Iran

Corresponding author: Leila
Asadpour

Tel: 09113383860

Email: asadpour@iaurasht.ac.ir

Address: Islamic Azad University,
Rasht Branch, Rasht, Iran

Received : 16 Feb 2015

Revised: 30 Jul 2015

Accepted: 01 Aug 2015

ABSTRACT

Background and Objective: *Campylobacters* are infectious zoonotic agents, and among the main bacterial causes of gastroenteritis in humans. Studies have shown that *Campylobacter jejuni* is of the main causes of infection among humans. Detection of these infectious agents in water resources is of great importance for maintaining the health of humans. Therefore, the aim of this study was molecular detection of *C. jejuni* strains in surface water samples collected from Rasht, Iran.

Methods: This cross-sectional descriptive study was performed on 45 surface water samples collected from the city of Rasht. After culture and isolation of bacteria, the molecular detection of *C. jejuni* was carried out using *hipO*-specific primers. Presence of cytolethal distending toxin (*cdt*) gene in positive samples was evaluated by polymerase chain reaction using *cdtC*-specific primers.

Results: Of 45 samples, seven (15.5%) were positive for *C. jejuni* contamination, five of which (71.4%) had the *cdtC* gene.

Conclusion: The prevalence of toxin-producing *C. jejuni* in surface waters of Rasht is notable. Therefore, it is recommended to take necessary measures for controlling the spread of this microorganism.

Keywords: *Campylobacter jejuni*, Surface water, *cdt* gene, PCR.

INTRODUCTION

Campylobacter is a genus of gram-negative microaerophilic bacilli, and one of the most important causes of gastroenteritis in humans (1). *Campylobacter jejuni* is the most common human clinical isolate, and the main cause of childhood diarrhea around the world (2). It is also an important factor in occurrence of Guillain-Barré syndrome (3). In many European countries, the rate of campylobacteriosis has been higher than salmonellosis (4, 5). Cytotoxic distending toxin (CDT) is one of the virulence factors produced by *Campylobacter*s. Some gram-negative bacteria including the *Campylobacter* spp. can produce this toxin. In 1988, Johnson and Lori were the first that reported CDT production by *Campylobacter* spp.

In a study, 41% of 718 *Campylobacter* isolates produced CDT (6). The toxin consists of three protein subunits: cdtA, cdtB, and cdtC (7) with molecular weight of 30, 29 and 21 kDa, respectively. These subunits have a tendency to associate with the outer membrane of the bacteria and contribute to delivery and activity of the toxin by simultaneous presence at the target site (6, 8). CdtA is the host-cell binding agent, while cdtB is the active part of halotoxin with deoxyribonucleic activity, requiring the presence of the cdtC subunit (9, 10). This subunit leads to formation of active toxins by connecting the other two subunits. The association between the presence of the *cdt* gene and the tissue invasion potential of *C. jejuni* has been demonstrated (11). Cdt causes morphological changes in the cell, damages the DNA, blocks cell proliferation in the G2 phase and prevents transition of cell cycle into the mitosis phase, leading to apoptosis (6, 12, 13). Some studies have reported the presence of the *cdt* gene in almost all *C. jejuni* isolates (14).

Campylobacteriosis is a common disease with a worldwide distribution among humans and animals (15). Poultry and poultry products are the most important sources of *C. jejuni* (16). Epidemiological studies have shown the high prevalence (40 to 100%) of *Campylobacter* spp. in chicken, duck and turkey (17-20). *C. jejuni* is isolated from flowing water, rivers, turkey slaughterhouse wastewater and even seawater (16, 18). The prevalence of *C. jejuni* in rivers and ponds has been variable, reaching up to 70% (8). Molecular screenings indicate that the presence of *Campylobacter* in water is

due to contamination by wild and domestic animal feces (17, 19, 20). Considering the importance of the disease and bacterial screening in surface waters, and the role of *cdt* in the pathogenesis of this bacterium, it is essential to determine the frequency of this gene in *C. jejuni* isolates from high-risk areas. Therefore, the aim of this study was to determine the level of contamination with *C. jejuni* in surface water samples collected from the suburbs of Rasht, and evaluate the frequency of the *cdt* gene among positive samples.

MATERIAL AND METHODS

Sample collection and bacterial isolation

This cross-sectional descriptive study was conducted on 45 water samples collected from the rivers around the city of Rasht in autumn, 2014. One-liter samples were collected from the river (distance of one meter and depth of 30 cm) in sterile containers. For isolation of bacteria, the samples were centrifuged at 6000 rpm for 15 minutes. The resulting precipitate was transferred to Preston agar, and then incubated under microaerophilic conditions at 42 °C. After 48 hours, 100 µl of the medium was transferred to the charcoal agar medium. The plates were incubated under microaerophilic conditions at 42 °C for 48 hours. Then, all suspected colonies were examined microscopically. Biochemical tests including oxidase, catalase and hippurate hydrolysis were performed on the microbial colonies to detect *Campylobacter* spp. (21, 22).

Extraction of bacterial genomic DNA

After culture of bacteria in Preston broth without antibiotics, and incubation at 42 °C on rotary incubator at 60 rpm for 24 hours, extraction and purification of genomic DNA were performed using AccuPrep® kit (Bioneer) according to the manufacturer's instructions. The purity of the extracted genome was assessed by electrophoresis on 1% agarose gel.

Polymerase chain reaction (PCR)

A primer specific for the *hipO* chromosomal locus of the bacterium was used for the molecular identification of *C. jejuni* (23). The characteristics of the primers used, the amount of reagents required for the PCR reaction and the thermocycler conditions for the amplification of the *hipO* gene are given in

Tables 1-3, respectively. Positive samples for *C. jejuni* were examined for the presence of *cdtC* gene using a specific primer for the gene (24) (Table 1). The amount of reagents used for the PCR reaction and the temperature conditions of the thermocycler were similar to the previous step, except for the annealing temperature of the *cdtC* primer (42 °C).

RESULTS

Gram-negative bacilli with wet, shiny and

flattened grayish colonies. These motile, catalase positive, oxidase positive and hippuricase positive isolates were identified as *C. jejuni*. A 735 bp fragment was produced in the PCR process using the *C. jejuni*-specific primer, which confirmed the results of the biochemical tests. Presence of the *cdtC* gene was detected in five of the seven (71.4%) *C. jejuni*-positive samples by identifying a 555 bp PCR product (Figure 2).

Table 1- Characteristics of the forward and reverse primers used in the study

Gene	Nucleotide sequence of primers	Length of PCR product
<i>hipO</i>	F:5'GAAGAGGGTTTGGGTGGTG3' R:5'AGCTAGCTTCGCATAATAACTTG3'	735bp
<i>cdtC</i>	F:5'TGGATGATAGCAGGGGATTTTAAC3' R:5'TTGCACATAACCAAAAGGAAG3'	555bp

Table 2- The amounts of reagents used for the PCR reaction

Reagents	Amount
dNTP mix (10mM)	1µl
MgCl ₂ (50mM)	1µl
10X PCR Buffer	2.5µl
Primer Forward	1µl
Primer Reverse	1µl
Taq DNA Polymerase	0.8 µl
DNA Template	3µl
D.D.W	14.7 µl
Total Volume	25 µl

Table 3- Thermocycler temperature programs for the amplification of the *hipO* gene

Step	Process	Temperature	Time	Number of cycles
First	Initial denaturation	95 °C	5 min	1
Second	Denaturation	94 °C	1 min	35
	Annealing	53 °C	1 min	
	Extension	72 °C	1 min	
Third	Final extension	72 °C	1 min	1

Figure 2- Detection of the *cdtC* after gel electrophoresis of PCR products. Column 1: 100 bp marker, Column 2: positive control, Column 3: negative control (PCR without DNA template), Columns 4, 5, 6, 8, 10: samples containing the *cdtC* gene, Columns 7 and 9: samples lacking the *cdtC* gene.



DISCUSSION

We determined the frequency of *C. jejuni* in rivers around the city of Rasht, and the frequency of *cdtC* gene among the identified bacteria. Bacterial culture and PCR of *hipO* chromosomal locus of the bacterium were used to identify *C. jejuni* in the samples. Hippurate hydrolysis test has a high specificity for detection of the bacterium, while positive results are observed in isolates containing the *hipO* gene.

In this study, 15.5% of the examined water samples were contaminated with *C. jejuni*. Limited number of studies in Iran has investigated the prevalence of *Campylobacter* in surface waters. In study of Ghane et al., *C. jejuni* was identified in 2.66% of samples collected from the Caspian Sea using culture and PCR methods (21). The prevalence of this bacterium in water resources vary in different parts of the world. According to study of Dungan et al. in the United States, the level of contamination with *C. jejuni* in pond water samples is reported to be 70% (25), while study of Meinersmann et al. in Georgia found that only 7.5% of the samples are contaminated with *C. jejuni* (26). Van Dyke et al. evaluated the prevalence of *Campylobacter* in 344 surface water samples collected in Canada, and reported that 69.8% of the samples were positive for *Campylobacter* spp. in the PCR method (18).

According to our results, 71.4% of the positive samples for *C. jejuni* contained the *cdtC* gene. Several studies have reported the high prevalence of the *cdt* gene in *C. jejuni* isolates.

REFERENCES

1. Nachamkin I, Szymanski CM, Blaser MJ. *Campylobacter*. ASM Press. 2008.
2. Olsen KN, Lund M, Skov J, Christensen LS, Hoorfar J. Detection of *Campylobacter* bacteria in air samples for continuous real-time monitoring of *Campylobacter* colonization in broiler flocks. *Appl Environ Microbiol*. 2009; 75(7): 2074-2078. doi: 10.1128/AEM.02182-08.
3. Guarino M, Casmiro M, D'Alessandro R. *Campylobacter jejuni* infection and Guillain-Barre syndrome: a case-control study. *Emilia-Romagna Study Group on Clinical and Epidemiological problems in neurology*. *Neuroepidemiology*. 1998; 17(6): 296-302.
4. De Cesare A, Sheldon BW, Smith KS, Jaykus L-A. Survival and persistence of *Campylobacter* and *Salmonella* species under various organic loads on food contact surfaces. *J Food Prot*. 2003; 66(9): 1587-94.

of *cdtA*, *B*, and *C* genes to be 58.3% in 10 clinical isolates of *C. jejuni* (27). In study of Talukder et al. in Bangladesh, 97.5% of *C. jejuni* isolates had the *cdtA*, *B*, *C* genes. The mentioned study reported that the genes are present in strains causing diarrhea in humans, indicating the pathogenic potential of *Campylobacter* spp. (28). In a study in Brazil, 83.3% of *C. jejuni* isolates from poultry and vegetable samples contained the *cdt* gene (29). In Iran, Shojaei et al. detected the *cdt* gene in 93% of *C. jejuni* isolates from intestine of poultry, in 65% of *C. jejuni* isolates from turkey intestine, and in 67% of isolates from the intestine of quail (30).

CONCLUSION

The results of this study indicate that the frequency of potentially virulent *C. jejuni* is high in the rivers of the suburbs of Rasht, Iran. Considering the role of surface waters in the dissemination of campylobacteriosis, it is recommended that some preventive measures be taken for controlling the spread of this microorganism.

ACKNOWLEDGMENTS

The authors would like to thank the Islamic Azad University of Rasht for the financial support.

CONFLICT OF INTEREST

All contributing authors declare no conflicts of interest.

5. Ica T, Caner V, Istanbul O, Nguyen HD, Ahmed B, Call DR, Beyenal H. Characterization of mono- and mixed-culture *Campylobacter jejuni* biofilms. *Applied and environmental microbiology*. 2011; 78(4):1033-1038.
6. Johnson WM, Lior H. A new heat-labile cytolethal distending toxin (CLDT) produced by *Campylobacter* spp. *Microb Pathog*. 1988; 4(2): 115-126.
7. Nakanishi S, Tazumi A, Moore JE, Millar BC, Matsuda M. Molecular and comparative analyses of the full-length cytolethal distending toxin (*cdt*) gene operon and its adjacent genetic loci from urease-positive thermophilic *Campylobacter* (UPTC) organisms. *Br J Biomed Sci*. 2010; 67(4): 208-215.
8. Dungan RS, Klein M, Leytem AB. Quantification of bacterial indicators and zoonotic pathogens in dairy wastewater ponds. *Appl Environ Microbiol*. 2012; 78(22): 8089-8095.

9. Lara-Tejero M, Galán JE. *CdtA, CdtB, and CdtC form a tripartite complex that is required for cytolethal distending toxin activity*. Infection and immunity. 2001; 69(7): 4358-4365.
10. Mao X, DiRienzo JM. *Functional studies of the recombinant subunits of a cytolethal distending holotoxin*. Cellular microbiology. 2002; 4(4): 245-255.
11. Purdy D, Buswell C, Hodgson A, McAlpine K, Henderson I, Leach S. *Characterisation of cytolethal distending toxin (CDT) mutants of Campylobacter jejuni*. Journal of medical microbiology. 2000; 49(5): 473-479.
12. Whitehouse CA, Balbo PB, Pesci EC, Cottle DL, Mirabito PM, Pickett CL. *Campylobacter jejuni cytolethal distending toxin causes a G2-phase cell cycle block*. Infection and immunity. 1998, 66(5): 1934-1940.
13. De Rycke J, Oswald E. *Cytolethal distending toxin (CDT): a bacterial weapon to control host cell proliferation?* FEMS microbiology letters. 2001; 203(2): 141-148.
14. Talukder KA, Aslam M, Islam Z, Azmi IJ, Dutta DK, Hossain S, et al. *Prevalence of virulence genes and cytolethal distending toxin production in Campylobacter jejuni isolates from diarrheal patients in Bangladesh*. Journal of clinical microbiology. 2008; 46(4): 1485-1488.
15. Southern JP, Smith RM, Palmer SR. *Bird attack on milk bottles: possible mode of transmission of Campylobacter jejuni to man*. Lancet. 1990; 336(8728): 1425-1427.
16. Pepe T, De Dominicis R, Esposito G, Ventrone I, Fratafico PM, Cortesi ML. *Detection of Campylobacter from poultry carcass skin samples at slaughter in Southern Italy*. J Food Prot 2009; 72(8): 1718-1721.
17. Debretsion A, Habtemariam T, Wilson S, Nganwa D, Yehualaeshet T. *Real-time PCR assay for rapid detection and quantification of Campylobacter jejuni on chicken rinses from poultry processing plant*. Mol Cell Probes. 2007; 21(3): 177-181.
18. Van Dyke MI, Morton VK, McLellan NL, Huck PM. *The occurrence of Campylobacter in river water and waterfowl within a watershed in southern Ontario, Canada*. J Appl Microbiol. 2010; 109(3): 1053-1066.
19. Kwan PS, Barrigas M, Bolton FJ, French NP, Gowland P, Kemp R, et al. *Molecular epidemiology of Campylobacter jejuni populations in dairy cattle, wildlife, and the environment in a farmland area*. Applied and environmental microbiology. 2008; 74(16): 5130-5138.
20. Emtiazi F, Schwartz T, Marten SM, Krolla-Sidenstein P, Obst U. *Investigation of natural biofilms formed during the production of drinking water from surface water embankment filtration*. Water research. 2004, 38(5):1197-1206.
21. Ghane M, Moein FG, Massoudian S. *The First Isolation of Campylobacter jejuni*. Advanced Studies in Biology. 2012; 4(9): 407-418.
22. Scates P, Moran L, Madden RH. *Effect of incubation temperature on isolation of Campylobacter jejuni genotypes from foodstuffs enriched in Preston broth*. Applied and environmental microbiology. 2003; 69(8): 4658-4661. doi: 10.1128/AEM.69.8.4658-4661.2003.
23. Zhao C, Ge B, De Villena J, Sudler R, Yeh E, Zhao S, et al. *Prevalence of Campylobacter spp., Escherichia coli, and Salmonella serovars in retail chicken, turkey, pork, and beef from the Greater Washington, D.C., area*. Appl Environ Microbiol. 2001; 67(12): 5431-5436.
24. Bang DD, Nielsen EM, Scheutz F, Pedersen K, Handberg K, Madsen M. *PCR detection of seven virulence and toxin genes of Campylobacter jejuni and Campylobacter coli isolates from Danish pigs and cattle and cytolethal distending toxin production of the isolates*. J Appl Microbiol. 2003; 94(6):1003-1014.
25. Peyrat MB, Soumet C, Maris P, Sanders P. *Phenotypes and genotypes of campylobacter strains isolated after cleaning and disinfection in poultry slaughterhouses*. Vet Microbiol. 2008; 128(3-4): 313-326.
26. Meinersmann R, Berrang M, Little E. *Campylobacter spp. recovered from the Upper Oconee River watershed, Georgia in a 4-year study*. Microbial ecology. 2013; 65(1): 22-27.
27. Ghorbanalizadgan M, Bakhshi B, Kazemnejad Lili A, Najar-Peerayeh S, Nikmanesh B. *A molecular survey of Campylobacter jejuni and Campylobacter coli virulence and diversity*. Iran Biomed J. 2014; 18(3): 158-164.
28. Talukder KA, Aslam M, Islam Z, Azmi IJ, Dutta DK, Hossain S, Nur EKA, Nair GB, Cravioto A, Sack DA, et al. *Prevalence of virulence genes and cytolethal distending toxin production in Campylobacter jejuni isolates from diarrheal patients in Bangladesh*. J Clin Microbiol. 2008; 46(4): 1485-1488.
29. Carvalho AF, Silva DM, Azevedo SS, Piatti RM, Genovez ME, Scarcelli E. *Detection of CDT toxin genes in Campylobacter spp. strains isolated from broiler carcasses and vegetables in São Paulo, Brazil*. Braz J Microbiol. 2013; 44(3): 693-699.
30. Shojaei Kavan R, Hassanzadeh M, Bozorgmehri Fard MH, Pourbakhsh SA, Akhondzadeh Basti A, Barin A, et al. *Detection of cytolethal distending toxin (cdt) Genes in Campylobacter jejuni and Campylobacter coli isolated from the intestinal of commercial broiler chickens, turkey and quail of Iran*. Iranian Journal of Veterinary Medicine. 2015; 9(2): 109-116.