

Parasites of the eyes of fresh and brackish water fishes in Iran

Barzegar, M.¹; Raeisi, M.²; Bozorgnia, A.³
and Jalali, B.^{4*}

¹Senior Expert of Fisheries Sciences, Consultancy Service PVT, Tehran, Iran; ²Department of Food Hygiene and Aquatic Animal Disease, Faculty of Veterinary Medicine, Shahrekord Branch, Islamic Azad University, Shahrekord, Iran; ³Department of Fisheries, Faculty of Natural Resources, Ghaemshahr Branch, Islamic Azad University, Ghaemshahr, Iran; ⁴Department of Aquatic Animal Disease, Faculty of Veterinary Science, Science and Research Branch, Islamic Azad University, Tehran, Iran

*Correspondence: B. Jalali, Department of Aquatic Animal Disease, Faculty of Veterinary Science, Science and Research Branch, Islamic Azad University, Tehran, Iran. E-mail: behiar_jalali@yahoo.com

(Received 18 Sept 2007; revised version 23 Apr 2008; accepted 6 May 2008)

Summary

In the present investigation a total of six parasite species, some eye-specific and others non-organ specific parasites were found in the eyes of 48 examined fish species, collected from fresh and brackish waters of Iran during 2004-2006. *Ichthyophthirius multifiliis* was isolated from the external surface of the eyes of *Oncorhynchus mykiss*, *Gyrodactylus stankovici* from *Cyprinus carpio* and *Lernaea cyprinacea* from *Cyprinus carpio* and *Hypophthalmichthys molitrix*. In the vitreous humor parasite species were *Tylodelphys clavata* (metacercaria) observed in *H. molitrix*, *Alburnus alburnus*, *Carassius auratus*, *Cyprinus carpio*, *Chondrostoma regium*, *Ctenopharyngodon idella* and *Capoeta capoeta*, and *Ornithodiplostomum* sp. (metacercaria) in *Aphanius vladykovi*. *Diplostomum spathaceum* (metacercaria) was found in the lens of eyes of 40 out of 48 fish species. Among parasite species identified, *Tylodelphys clavata* and *Ornithodiplostomum* sp. are recorded in Iran for the first time. Additionally, the geographical distribution and host range of *Diplostomum spathaceum* metacercaria is also presented in this study.

Key words: Parasites, Eye, Fish, Fresh and brackish water, Iran

Introduction

Previously, little attention was paid to the eye parasites of fishes in Iran and the records were limited mostly to infection and disease caused by *Diplostomum spathaceum* metacercaria in the lens of eyes of several freshwater fish species. However, extensive studies have recently been carried out on the parasites of eyes in riverine and lacustrine fishes inhabiting waters in the Zagros mountain area in western Iran. Therefore, our knowledge about the parasites of fish eyes has increased enormously and several new species have been identified (Barzegar and Jalali, 2002, 2006; Jalali and Barzegar, 2005, 2006; Raeisi *et al.*, 2006).

The economic significance of the eye diseases of cultured fish, is associated with specific effects or non-specific side effects of

parasites, including impairment of vision that leads to exophthalmus, cataract and even complete collapse of the eye, which may be the cause of growth inhibition and death of significant portions of cultured fishes.

In the present study, parasites of the eyes of fishes from fresh and brackish waters of Iran, with special attention to the host range of *D. spathaceum* metacercaria are reported. Of these parasites, two new digenean metacercaria species (*Tylodelphys clavata* and *ornithodiplostomum* sp.) found in the vitreous humor are recorded in Iran for the first time.

Materials and Methods

The present study was carried out in several areas of three different zoogeographical regions (Sarmatian,

Mesopotamian and Oriental) of Iran (Fig. 1) during 2004-2006. The fresh fishes were caught and transported to the laboratory, where their eye balls were completely separated and examined with naked eye and under a stereomicroscope at magnification of $\times 4$ to 40. The identification of fish hosts was carried out by an Iranian ichthyologist according to Coad (1992), Berg (1964-65) and Abdoli (1999). In doubtful cases, the whole specimens were fixed in 4% formalin and preserved for more investigation.



Fig. 1: Collection sites of examined fish species in Iran (Coad, 1992)

Methods used for collecting, fixing, staining and mounting of parasite specimens were as follows:

Protozoa: for collection of *Ichthyophthirius multifiliis* specimens, eye conjunctiva were finely scraped onto a microslide, and covered carefully with a coverslip. The samples were exposed to Bouin's fluid for about 15 min and then mounted in Canada balsam after dehydration in accordance with Lom and Dykova (1992).

Digenea: metacercaria was collected in a 0.9% saline solution. The sample was placed with a little saline on a glass slide and appropriate pressure was applied; it was fixed and washed in 90 and 70% alcohol, respectively, and then stained with alum haematoxylin according to Roberts (2001).

Crustacea: specimens of *Lernaea* sp. were collected from the eyes of the infected fish and cleaned in saline. The samples were

preserved in 70% alcohol, stained and cleared with polyvinyl lactophenol and mounted in Canada balsam according to Fernando *et al.* (1972).

Results

Six parasite species, found on the surface or in the eyes of the examined fishes are listed in Table 1.

Discussion

Various parasite species at different stages of life span have been found in the eye and associated structures of fish. Roberts (2001) stated that few *Myxobolus* spp. infected the sclera (*M. hoffmani*, *M. scleroperca*), anterior chamber and iris (*M. couseii*) of both fresh- and seawater fishes of Canada. Vitreous body, lens and retina are the predilection site of several metacercaria of Strigeidida order, where pressing against the cornea and other orbital locations cause collapse of the eye. Probably the most significant damage, called worm cataract and subsequent growth retardation, caused by *Diplostomum* spp. metacercaria; this parasite is now widespread throughout Iran. Hoglund (1991) reported that at least 125 fish species are considered as the second intermediate host of *D. spathaceum* in the world, of these, more than 40 fish species are recorded to be infected with *D. spathaceum* (metacercaria) in Iran (Table 1). Common carp and Chinese carp in ponds are seriously threatened by the eye parasites either due to diplostomiasis, the causative agent of blindness and subsequent emaciation or by spending more time at the surface of pond waters where they can be readily eaten by the piscivorous bird species. In addition to *D. spathaceum* metacercaria which is the most well known fish digenean parasite in Iran, in this study two new digenean metacercaria, namely *Tylodelphys clavata* and *Ornithodiplostomum* sp. are reported for the first time in Iran. These two new parasites were found in the vitreous humor of the eyes of exotic and native cyprinid fishes inhabiting Chaghakhour and Gandoman Lagoon. *Tylodelphys* sp. was

Table 1: Parasites found on or in the eyes of freshwater fishes of Iran

Parasites			Host(s)	Locality(s)	Region(s)	Reference(s)
Parasite group	Species	Family	Species			
Protozoan	<i>Ichthyophthirius multifiliis</i> (Fouquet, 1876)	Salmonidae	<i>Oncorhynchus mykiss</i>	Haraz Ri	East of Caspian	Present study
Monogenean	<i>Gyrodactylus stankovici</i> (Ergens, 1970)	Cyprinidae	<i>Cyprinus carpio</i>	Gilan fish hatcheries	West of Caspian	Jalali, 1998
Digenean	<i>Diplostomum spathaceum</i> (Rudolphi, 1819) (metacercaria) Fig. 2	Acipenseridae	Acipenser fish	Shahid Beheshti fish hatchery	West of Caspian	Ghoroghi, 1991
				Gilan fish hatcheries	West of Caspian	Mokhayer, 1989; Jalali, 1998; Nezam abadi and Abdi, 2002
		Clupeidae	<i>Alosa caspia persica</i>	Caspian Sea*	Caspian Sea	Present study
		Cyprinidae	<i>Abramis brama</i>	Gilan fish hatcheries	West of Caspian	Nezam abadi and Abdi, 2002
			<i>Alburnus alburnus</i>	Chaghakhour Lag	Tigris	Present study
				Gandoman Lag	Tigris	Present study
			<i>Alburnoides bipunctatus</i>	Saryson Ri	West of Caspian	Rahanandeh, 2006
			<i>Aspius vorax</i>	Khuzestan fish hatcheries	Karoon	Jalali, 1998
			<i>Barbus lacerta</i>	Vahdat Res	Tigris	Barzegar and Jalali, 2006
			<i>Barbus sharpeyi</i>	Khuzestan fish hatcheries	Karoon	Jalali, 1998
			<i>Barbus</i> sp.	Khuzestan fish hatcheries	Karoon	Jalali, 1998
			<i>Blicca bjoerkna</i>	Amirkalayeh Lag	East of Caspian	Khara <i>et al.</i> 2005
				Golestan fish hatcheries	East of Caspian	Mokhayer, 1989
			<i>Capoeta aculeata</i>	Chaghakhour Lag	Tigris	Present study
			<i>Capoeta damascina</i>	Zayande-rud Ri	Esfahan	Present study
			<i>Capoeta capoeta</i>	Makoo Res	Azərbayjan	Abdolmaleki, 2000
				Baroun Res	Azərbayjan	Masoumian <i>et al.</i> 2005
				Ghala-jough	Azərbayjan	Pazooki <i>et al.</i> 2007
				Golestan fish hatcheries	East of Caspian	Mokhayer, 1989
				Sanandaj fish hatcheries	Tigris	Khancheh-sepehrredin, 2000
			<i>Carassius auratus gibelio</i>	Makoo Res	Azərbayjan	Abdolmaleki, 2000
				Hamoon Lag	Sistan	Sharif Rohani, 1994
				Chaghakhour Lag	Tigris	Present study
				Boojagh Lag	West of Caspian	Khara <i>et al.</i> 2004
				Telar Ri	East of Caspian	Present study
				Amirkalayeh Lag	East of Caspian	Khara <i>et al.</i> 2005
			<i>Chalcalburnus mossulensis</i>	Kaftar La	Neyriz	Barzegar and Jalali, 2002
			<i>Chalcalburnus</i> sp.	Zarivar La	Tigris	Jalali and Barzegar, 2006
			<i>Chondrostoma regium</i>	Zayande-rud Ri	Esfahan	Present study
				Kaftar La	Neyriz	Barzegar and Jalali, 2002
				Chaghakhour Lag	Tigris	Present study
			<i>Ctenopharyngodon idella</i>	Hamoon Lag	Sistan	Sharif Rohani, 1994
				Kaftar La	Neyriz	Barzegar and Jalali, 2002
				Chaghakhour Lag	Tigris	Present study
				Gilan fish hatcheries	West of Caspian	Rahanandeh, 2006
				Mazandaran and Gilan fish hatcheries	Caspian	Mokhayer, 1989 and Jalali, 1998
			<i>Cyprinus carpio</i>	Vahdat Res	Tigris	Barzegar and Jalali, 2006
				Hamoon Lag	Sistan	Sharif Rohani, 1994
				Hoorolazim Lag	Karoon	Moghainemi, 1995
				Kaftar La	Neyriz	Barzegar and Jalali, 2002
				Chaghakhour Lag	Tigris	Present study
				Hatcheries fish in Gilan pro	West of Caspian	Rahanandeh, 2006
				Aras Res	Azərbayjan	Masoumian <i>et al.</i> 2005
				Boojagh Lag	West of Caspian	Khara <i>et al.</i> 2004
				Mazandaran and Gilan fish hatcheries	Caspian	Mokhayer, 1989
				Sanandaj fish hatcheries	Tigris	Khancheh-sepehrredin, 2000
			<i>Hypophthalmichthys molitrix</i>	Zarivar La	Tigris	Jalali and Barzegar, 2006
				Chaghakhour Lag	Tigris	Present study

Ri = River, La = Lake, Lag = Lagoon, Pro = Province, Spi = Spring water, Res = Reservoir and *Brackish water

Table 1 Continued

Parasites		Host(s)		Locality(s)	Region(s)	Reference(s)		
Parasite group	Species	Family	Species					
Digenean	<i>Diplostomum spathaceum</i> (Rudolphi, 1819) (metacercaria)	Cyprinidae	<i>Hypophthalmichthys molitrix</i>	Gilan fish hatcheries	West of Caspian	Rahanandeh, 2006		
				Mazandaran and Gilan fish hatcheries	Caspian	Mokhayer, 1989 and Jalali, 1998		
				Sanandaj fish hatcheries	Tigris	Khancheh-sepehrredin, 2000		
			<i>Leuciscus lepidus</i>	Zayande-rud Ri	Esfahan	Present study		
				Sanandaj fish hatcheries	Tigris	Khancheh-sepehrredin, 2000		
			<i>Hypophthalmichthys nobilis</i>	Gilan fish hatcheries	West of Caspian	Rahanandeh, 2006		
				Mazandaran and Gilan fish hatcheries	Caspian	Mokhayer, 1989 and Jalali, 1998		
			<i>Rutilus rutilus caspicus</i>	Boojagh Lag	West of Caspian	Khara <i>et al.</i> 2004		
				Sijoal	East of Caspian	Present study		
				Amirkalayeh Lag	East of Caspian	Khara <i>et al.</i> 2005		
				Caspian Sea	Caspian Sea	Masoumian <i>et al.</i> 2001		
			<i>Rutilus rutilus kutum</i>	Boojagh Lag	West of Caspian	Khara <i>et al.</i> 2004		
				Shahid Rajaei fish hatchery	East of Caspian	Present study		
				Kileh Spi	East of Caspian	Present study		
				Mazandaran and Gilan fish hatcheries	Caspian	Mokhayer, 1989		
			<i>Schizothorax zarudnyi</i>	Hamoon Lag	Sistan	Sharif Rohani, 1994		
			<i>Schizothorax pelzami</i>	Hamoon Lag	Sistan	Sharif Rohani, 1994		
			<i>Tinca tinca</i>	Amirkalayeh Lag	East of Caspian	Khara <i>et al.</i> 2005		
			Cyprinodontidae	<i>Aphanius vladykovi</i>	Chaghakhour Lag	Tigris	Present study	
					Gandoman Lag	Tigris	Present study	
			Esocidae	<i>Esox lucius</i>	Amirkalayeh Lag	East of Caspian	Khara <i>et al.</i> 2005	
					Boojagh Lag	West of Caspian	Khara <i>et al.</i> 2004	
					Amirkalayeh Lag	East of Caspian	Khara <i>et al.</i> 2005	
					Gilan fish hatcheries	West of Caspian	Nezam abadi and Abdi, 2002	
			Mugilidae	<i>Liza abu</i>	Hoorolazim Lag	Karoon	Moghainemi, 1995	
			Mastacembelidae	<i>Mastacembelus mastacembelus</i>	Zarivar La	Tigris	Jalali and Barzegar, 2006	
			Percidae	<i>Perca fluviatilis</i>	Amirkalayeh Lag	East of Caspian	Khara <i>et al.</i> 2005	
				<i>Sander lucioperca</i>	Gillan fish hatcheries	West of Caspian	Nezam abadi and Abdi, 2002	
			Salmonidae	<i>Oncorhynchus mykiss</i>	Makoo Res	Azarbaijan	Masoumian <i>et al.</i> 2005	
					Fish farms in Azerbaijan	Azarbaijan	Nekoei fard and Dini talatapeh, 2000	
					Urmia fish Hatcheries	Azarbaijan	Asadzadeh Mangili and Ghorbanzadeh, 1998	
					Urmia fish Hatcheries	Azarbaijan	Naghili, 2001	
			Siluridae	<i>Silurus glanis</i>	Amirkalayeh Lag	East of Caspian	Khara <i>et al.</i> 2005	
			Percidae	<i>Perca fluviatilis</i>	Amirkalayeh Lag	East of Caspian	Khara <i>et al.</i> 2004	
			<i>Tylodelphys clavata</i> (Nordman, 1832) (metacercaria) Fig. 3	Cyprinidae	<i>Alburnus alburnus</i>	Chaghakhour Lag	Tigris	Present study
					<i>Capoeta aculeata</i>	Chaghakhour Lag	Tigris	Present study
						Gandoman Lag	Tigris	Present study
					<i>Carassius auratus</i>	Chaghakhour Lag	Tigris	Present study
						Gandoman Lag	Tigris	Present study
					<i>Chondrostoma regium</i>	Chaghakhour Lag	Tigris	Present study
					<i>Ctenopharyngodon idella</i>	Chaghakhour Lag	Tigris	Present study
					<i>Cyprinus carpio</i>	Chaghakhour Lag	Tigris	Present study
					<i>Hypophthalmichthys molitrix</i>	Chaghakhour Lag	Tigris	Present study
<i>Barbus grypus</i>	Hoor-o-Azim Lag	Tigris			Moghainemi, 1995			
<i>Cyprinus carpio</i>	Hoor-o-Azim Lag	Tigris			Moghainemi, 1995			
	Hoor-o-Azim Lag	Tigris			Moghainemi, 1995			
Mugilidae	<i>Liza abu</i>	Hoor-o-Azim Lag			Tigris	Moghainemi, 1995		
<i>Orinithodiplostomum</i> sp. (metacercaria) Fig. 4	Cyprinodontidae	<i>Aphanius vladykovi</i>			Chaghakhour Lag	Tigris	Present study	
					Gandoman Lag	Tigris	Present study	
Crustacean	<i>Lernaea cyprinacea</i> (adult)	Cyprinidae			<i>Cyprinus carpio</i>	Mazandaran fish hatcheries	East of Caspian	Jalali, 1987
			<i>Hypophthalmichthys molitrix</i>	Mazandaran fish hatcheries	East of Caspian	Jalali, 1987		

Ri = River, La = Lake, Lag = Lagoon, Pro = Province, Spi = Spring water, Res = Reservoir and * Brackish water



Fig. 2: *Diplostomum spathaceum* (×400)




Fig. 3: *Tylodelphys clavata*




Fig. 4: *Ornithodiplostomum* sp.

first reported from the eyes of *Liza abu*, *Cyprinus carpio* and *Barbus grypus* in Hoor-o-Azim by Moghainemi (1995); however in this study, the parasite is identified to species level and in a new locality. The occurrence of *Ornithodiplostomum* sp. (metacercaria) is also reported for the first time in Iran.

From epidemiological point of view, both parasites are Palearctic species therefore, they may be native or translocated by introduced fishes (common carp or silver carp) from the Caspian Sea to Chaghakhour Lagoon. Investigation on the relationship between the host-parasite systems may

elucidate the sources of two digenean metacercaria in Chaghakhour Lagoon. Additional investigations, including DNA analysis are the best option for specific recognition of metacercaria differences or similarities between species found in either Chaghakhour Lagoon or Caspian Basin and European species. *Ichthyophthirius multifiliis*, the most common external holotrich parasite, can be frequently seen on the eyes of fishes, particularly in fry and fingerlings during heavy infection. *Gyrodactylus stankovici* may sometimes infect the exterior part of fish eyes (Jalali, 1998).

Finally, *Lernaea cyprinacea*, the most common copepodid parasite in the freshwater aquaculture in Iran, is very pathogenic to small fish due to its relatively large size. In our finding, premetamorphosed females penetrate the margin of the eye ball of common carp fingerlings and cause exophthalmus.

In conclusion, most of the eye-specific parasites found in Iranian fresh and brackish water fishes cause various degrees of blindness. Although, the infected fishes are not killed directly by the parasite, however, related growth retardation, behavioural changes and associated secondary invaders (piscivorous bird, bacteria and external protozoa) may lead to death of the infected fish.

Acknowledgement

The authors express their thanks to Prof. K. Molnar for his valuable suggestion on site preference of *Gyrodactylus stankovici*.

References

- Abdoli, A (1999). *The inland water fishes of Iran*. 1st Edn., Nature Museum and World Wild of Iran. P: 377 (In persian).
- Abdolmaleki, Sh (2000). *Survey of parasites of fishes of Makoo reservoir*. Dept. of Fisheries and Sciences. Iran. P: 21 (In persian).
- Asadzadeh Manjili, A and Ghorbanzadeh, A (1998). Survey on infestation of rainbow trout to *Diplostomum spathaceum* in west part of Azarbaijan province. Iran. Sci. Fish. J., 4: 103-110 (In persian).
- Barzegar, M and Jalali, B (2002). Parasites of

- Kaftar lake fishes, their geographical distribution and economical importance. Journal of the School of Veterinary Medicine. Shahid Chamran University of Ahvaz. 5: 52-64 (In persian).
- Barzegar, M and Jalali, B (2006). Helminthes, Acanthocephala and Crustacean parasites of fishes in Vahdat reservoir. Iran. J. Vet. Sci., 3: 229-234 (In Persian).
- Berg, LS (1964-65). *Freshwater fishes of USSR and adjacent countries*. Vol. 3, 1st Edn., Nauka, Moscow, USSR. P: 496 (In russian).
- Coad, BW (1992). *Freshwater fishes of Iran. A checklist and bibliography*. Ichthyology Section. Canadian Museum of Nature. Ottawa, Ontario, Canada. P: 66.
- Fernando, CH; Furtado, JJ; Gussev, AV; Hanek, G and Kakong, SA (1972). *Methods for the study of freshwater fish parasites*. 1st Edn., University of Waterloo, Biology Series. P: 76.
- Hoglund, J (1991). Ultrastructure observations and radiometric assay on cercarial penetration and migration of the *Diplostomum spathaceum* in the rainbow trout *Oncorhynchus mykiss*. Parasitol. Res., 77: 283-289.
- Ghoroghi, A (1991). Report on *Diplostomum spathaceum* in acipenserid fingerlings in Shahid Beheshti hatchery. Fisheries Research Organization. P: 54 (In persian).
- Jalali, B (1987). Lernaeosis in cyprinid cultured fish in Iran. Master Thesis. Godolo University. Hungary. P: 21.
- Jalali, B (1998). *Parasites and parasitic diseases of fresh water fishes of Iran*. 1st Edn., Fisheries Co. of Iran. P: 564 (In persian).
- Jalali, B and Barzegar, M (2005). A survey on parasites of gills of fishes of Vahdat reservoir. Iran. J. Vet. Sci., 3: 41-50.
- Jalali, B and Barzegar, M (2006). Fish parasites in Zarivar lake. J. Agr. Sci. Technol., 8: 47-59.
- Khara, H; Nezami, ShA; Sattari, M; Mirhasheminasab, SF and Mousavi, SA (2005). An investigation on fish infection *Diplostomum spathaceum* in Amirkalayeh wetland. Iran. Sci. Fish. J., 14: 49-66.
- Khara, H; Sattari, M; Nezami, Sh; Mirhasheminasab, SF; Mousavi, SA; Taati, R and Tatina, M (2004). Parasites and incidence of parasites of intestine of some economic fishes in Bojagh Lagoon. Iran. Sci. Fish. J., 15: 9-18 (In persian).
- Khancheh-sepehrredin, K (2000). Survey on *Diplostomum spathaceum* in warm water fish cultured and natural water resources in Sanandaj-Kurdistan. Thesis of DVM, Veterinary Dept. Urmia Islamic Azad University. No. 394 (In persian).
- Lom, J and Dykova, I (1992). *Protozoan parasites of fishes*. 1st Edn., Amsterdam, Netherlands, Elsevier Science Publisher. P: 315.
- Masoumian, M; Mehdizadeh, J and Mokhayer, B (2001). Study on parasitic infestation of *Rutilus rutilus caspius* in south east of Caspian Sea. Iran. J. Fish. Sci., 4: 61-74 (In persian).
- Masoumian, M; Pazooki, J; Yahyazadeh, M and Teymornezhad, A (2005). Protozoan from freshwater fishes from north west of Iran. Iran. J. Fish. Sci., 4: 31-42.
- Moghainemi, R (1995). *Survey on parasites of fishes of Hoorolazim Lagoon (Dashte Azadegan)*. Dept. of Fisheries and Sciences. Iran. P: 107.
- Mokhayer, B (1989). Diplostomiasis in Iran. J. Fac. Vet. Med., University of Tehran. 44: 17-24 (In persian).
- Naghili, H (2001). Survey on *Diplostomum spathaceum* in cold water fishes in west Azarbaijan province. Thesis of DVM Veterinary Dept. Urmia Azad University. No. 497 (In persian).
- Nekoe fard, A and Dini talatapeh, H (2000). Survey on disease caused by *Diplostomum spathaceum* and *Ichthyophthirius multifiliis* in cultured rainbow trout hatchery in west Azarbaijan province. First symposium of health and disease of aquatic animal. Ahvaz. 14-16 Feb. (In persian).
- Nezam abadi, H and Abdi, K (2002). Review of diplostomiasis in fishes. Aquatic Health and Disease Office. Veterinary Organization of Iran. Final Report. P: 44 (In persian).
- Pazooki, J; Masoumian, M; Yahyazadeh, HM and Abbasi, K (2007). Metazoan parasites from freshwater fishes of northeast Iran. Iran. J. Fish. Sci., 9: 25-33 (In persian).
- Rahanandeh, M (2006). *Reports on diplostomiasis in Gilan province*. Vet. Dept., Sciences and Research Branch, Islamic Azad University. P: 27.
- Raeisi, M; Barzegar, M; Jalali, B and Alimardani, K (2006). A survey on monogenean parasites of gills of fishes in Chaghakhor lake. Iran. J. Vet. Sci. 1: 411-418.
- Roberts, RJ (2001). *Fish pathology*. 1st Edn., London, UK, Harcourt Publisher Fishing News Books Limited. P: 472.
- Sharif Rohani, M (1994). Survey on parasites and parasitic diseases in Sistan region. *Proceeding of the 2nd symposium of Iranian veterinary clinics, Tehran*. Iran. P: 109.