

Short Paper

The effect of sexuality on some haematological parameters of the yellowfin seabream, *Acanthopagrus latus* in Persian Gulf

Karimi, Sh.¹; Kochinian, P.² and Salati, A. P.^{2*}

¹Graduated from Faculty of Marine Natural Resources, Khorramshahr University of Marine Science and Technology, Khorramshahr, Iran; ²Department of Fisheries, Faculty of Marine Natural Resources, Khorramshahr University of Marine Science and Technology, Khorramshahr, Iran

*Correspondence: A. P. Salati, Department of Fisheries, Faculty of Marine Natural Resources, Khorramshahr University of Marine Science and Technology, Khorramshahr, Iran. E-mail: apsalati@kmsu.ac.ir

(Received 8 Nov 2011; revised version 1 Sept 2012; accepted 23 Sept 2012)

Summary

Acanthopagrus latus is one of the major commercially exploited species in the Persian Gulf. In recent years, it has gained importance as a candidate for aquaculture. Haematological parameters of wild-caught specimens were studied during the pre-spawning period to determine if they are influenced by the sex of fish. Blood samples were collected from 55 Fish (30 female and 25 male) captured from the Musa Creek in the north-west Persian Gulf. RBC counts were higher in male than in female fish ($P < 0.05$) and other parameters such as WBC, Hct, Hb, MCV, MCH, MCHC and leukocyte differential count did not show a significant difference between male and female fish.

Key words: Persian Gulf, *Acanthopagrus latus*, Haematology, Sex

Introduction

Haematologic assessment is an important tool for evaluation of fish physiological status such as other domestic animals (Kori-Siakpere *et al.*, 2005). Analysis of haematological characteristics for detection of stress or disease is a common practice. Fish blood parameters vary in different species and the individual variation has also been reported, which could be attributed to genetic variation, nutritional status, sex, age, stress caused by capture, handling and sampling procedures (Kori-Siakpere *et al.*, 2005). Hence, it is necessary to establish normal values as different parameters can affect haematologic parameters of fish. One of these parameters is sex (Mulcahy, 1970; Joshi, 1980).

Acanthopagrus latus (Subfamily: Sparidae) is a protandrous hermaphrodite fish distributed in southern Japan, southeastern China, Taiwan, southeastern

Asia, the Persian Gulf and Australia and in the Indian Ocean to southeastern Africa (Hayashi, 1993). It is also one of the leading fish species with high mariculture potential in the Persian Gulf. Effect of pollutants on fish blood has been studied (Hedayati *et al.*, 2010; Hedayati *et al.*, 2011; Hedayati, 2012), but haematologic characteristics have not been succinctly documented. Therefore, this study was done to catalogue the basic values of *A. latus* in order to establish any statistical variability between male and female in a population subjected to the same environmental conditions.

Materials and Methods

Fifty five *A. latus* were captured from the Musa Creek (north-west Persian Gulf) during the pre-spawning months (October-December). Careful netting and handling was implemented to minimize stress. The specimens were anesthetized with clove oil

and the peripheral blood was collected by puncture of the caudal vein with a heparinized syringe. Fish were weighed and measured and blood was sampled from the caudal vein with heparinized syringes. After sampling, the fish was killed by a sharp blow to the head and sex was determined according to histomorphology of gonads. Thirty were females (total length 25.58 ± 3.52 , weight 349.1 ± 126.27) and 25 were male (total length 23.99 ± 2.63 , weight 258.01 ± 128.10). Blood samples were transferred at 4°C to laboratory for analysis. RBCs and WBCs were manually counted after dilution with Natt-Herrick's. Hematocrit value was determined by the standard microhematocrit method. Hb was determined by cyanomethemoglobin method (Mostaghni *et al.*, 2005). A differential white cell count was done with giemsa stained blood smear to determine the number of different types of WBC. Each sample was analysed in duplicate. Mean corpuscular hemoglobin (MCH), mean corpuscular hemoglobin concentration (MCHC) and mean corpuscular volume (MCV) were calculated by standard formula (Coles, 1986). Data are expressed as mean \pm SE. Differences in haematological parameters between male and female fish were statistically analysed by Student's t-test.

Result

The mean values, standard error and range of haematologic parameters are

summarized in Table 1. The haematological profile of *A. latus* indicated a significant difference ($P<0.05$) in RBC counts between male and female. RBC counts were higher in males compare to females but this pattern was not recorded for other RBC related parameters such as Hb and Hct. WBC counts did not show any significant difference between male and female fish. In differential counts of WBCs eosinophil and basophil were not observed in blood smear. Other WBCs did not show a significant difference between male and female.

Discussion

There is a growing interest in the study of haematological parameters of fish as an important tool in assessing the physiological conditions for aquaculture purposes. This study showed the normal values of haematologic parameters of *A. latus* in the wild during the pre-spawning period. Dacie and Lewis (1991) believed gender has a great influence on haematology of fish and included it among factors influencing fish haematology. There are reports of sex-related differences in haematologic parameters of fish (Gabriel *et al.*, 2004; Akinrotimi *et al.*, 2007). Sex could be associated with differences in blood cell constituents that could be caused by higher metabolic activity of male versus female fish (Collazos *et al.*, 1998). The RBC count and related parameters were higher in male cyprinid fish (Orun *et al.*, 2003). Number of RBCs in *A. latus* was similar to values

Table 1: Hematological and biological parameters in male and female *A. latus*. Values are given as mean \pm SD. Different letters denote significant differences ($P<0.05$) between treatments

Parameters	Mean \pm SD		Max		Min	
	Male	Female	Male	Female	Male	Female
Weight (g)	285.01 \pm 128.1	349.1 \pm 169.27	690	796.94	182.7	191.43
Total length (cm)	23.99 \pm 2.63	25.58 \pm 3.52	30	34	20	21
RBC ($\times 10^6 \text{ mm}^3$)	3.37 \pm 0.6 ^a	3.02 \pm 0.67 ^b	4.69	5.46	2.55	2.09
WBC ($\times 10^3 \text{ mm}^3$)	5.72 \pm 1.99	5.45 \pm 1.89	9.75	9.5	3.75	2.5
Hemoglobin (g/dL)	6.76 \pm 2.83	7.74 \pm 3.83	12.51	14.95	2.68	2.12
Hematocrit (%)	31.18 \pm 5.41	29.07 \pm 5.96	38.46	42	19.35	18.18
MCV (nm^3)	93.69 \pm 25.16	95.32 \pm 32.87	123.68	167.59	53.53	46.77
MCH (p.g)	20.54 \pm 9.53	26.20 \pm 13.71	38.5	53.83	8.53	6.53
MCHC (g/dL)	23.01 \pm 11.68	30.52 \pm 16.58	48.45	65.81	8.62	8.81
Lymphocytes (%)	72.87 \pm 4.64	74.22 \pm 6.60	79	80	65	64
Neutrophils (%)	19.62 \pm 3.29	17.28 \pm 1.97	23	20	15	15
Monocytes (%)	7.51 \pm 2.5	8.5 \pm 2.51	11	11	5	5

reported for *Rhamdia quelen* (Borges *et al.*, 2004), *Dicentrarchus labrax* L. (Ozretic *et al.*, 2001) and *Cichlasoma dimerus* (Vazquez and Guerrero, 2007), but higher compared to *Anguilla anguilla* (Sahan *et al.*, 2007) and *Brycon amazonicus* (Tavares-Dias *et al.*, 2008).

Hb and Hct values for the *A. latus* in the present study were higher than reported for *Heterotis niloticus* (Fagbenro *et al.*, 2000) and obtained for *Parachanna obscura* (Kori-Siakpere *et al.*, 2005), but lower than *C. gariepinus* (Sowunmi, 2003) and *Synodontis membranacea* (Owolabi, 2011). Hemoglobin content is affected by endogenous factors such as age and general condition of the fish and also exogenous factors, handling, transport and sampling (Kori-Siakpere *et al.*, 2005) that make it difficult to compare results.

In fish, as in other vertebrates, WBC count is frequently used as indicators of health status. These cells are key components of innate immune defense and are involved in the regulation of immunological function in the organism (Ballarin *et al.*, 2004). WBC values did not show a significant difference between male and female fish. Differences in the WBC counts could be attributed to biotic (age, maturity, sex, pathogens) and abiotic (water temperature, pH, dissolved oxygen content) factors and in particular to stress (Pavlidis *et al.*, 2007).

Lymphocyte and neutrophil constituted the most peripheral WBCs like other reports in *Tilapia zilli* (Ezzat *et al.*, 1973) and *Oreochromis aureus* (Silveira and Rigores, 1989). Eosinophils and basophils were not observed in blood smear. Absence of eosinophils and in particular basophils, has been reported in some species (Rowley *et al.*, 1988).

Our findings showed that there is a similarity between haematologic indices in male and female *A. latus*, but complementary studies such as cytochemistry and ultra structure should be done.

References

Akinrotimi, OA; Gabriel, UU; Anyanwu, PE and Anyanwu, AO (2007). Influence of sex, acclimation methods and period on Haemato-

- logy of *Sarotherodon melanotheron*. Res. J. Biol. Sci., 2: 348-352.
- Ballarin, L; Dall'Oro, M; Bertotto, D; Libertini, A; Francescon, A and Barbaro, A (2004). Haematological parameters in *Umbrina cirrosa* (Teleostei, Sciaenidae): a comparison between diploid and triploid specimens. Comp. Biochem. Physiol. A., 138: 45-51.
- Borges, A; Scotti, LV; Siqueira, DR; Jurinitz, DF and Wassermann, GF (2004). Hematologic and serum biochemical values for jundiá (*Rhamdia quelen*). Fish Physiol. Biochem., 30: 21-25.
- Coles, EH (1986). *Veterinary clinical pathology*. 3rd Edn., Philadelphia, W. B. Saunders Co., P: 394.
- Collazos, ME; Ortega, E; Barriga, C and Rodriguez, AB (1998). Seasonal variation in hematological parameters male and female *Tinca tinca*. Mol. Cell Biochem., 183: 165-168.
- Dacie, JV and Lewis, SN (1991). *Practical haematology*. 5th Edn., Edinburgh, Churchill Livingstone, P: 390.
- Ezzat, AA; Shabana, MB and Farghaly, AM (1973). Studies on the blood characteristics of *Tilapia zilli* (Gervais). I. Blood cells J. Fish Biol., 6: 1-12.
- Fagbenro, O; Adedire, CO; Ayotunde, EO and Faminu, EO (2000). Haematological profile, food composition and enzyme assay in the gut of the African bony-tongue fish, *Heterotis (Clupisudis) niloticus* (Cuvier 1829) (Osteoglossidae). Trop. Zool., 13: 1-9.
- Gabriel, UU; Ezeri, GNO and Pabunmi, OO (2004). Influence of sex, source, health status and acclimation on the haematology of *Clarias gariepinus* (Burch, 1822). Afr. J. Biotechnol., 3: 463-467.
- Hayashi, M (1993). Sparidae. In: Nakabo, T (Ed.), *Fishes of Japan with pictorial key to the species*. 1st Edn., Tokyo, Tokai University Press. PP: 746-749.
- Hedayati, A (2012). Effect of marine mercury toxicity on immunological responses of seabream. Asian J. Anim. Sci., 6: 1-12.
- Hedayati, A; Safahieh, A; Savari, A and Marammazi, JG (2010). Assessment of aminotransferase enzymes in yellowfin sea bream under experimental condition as biomarker of mercury pollution. World J. Fish Mar. Sci., 2: 86-92.
- Hedayati, A; Safahieh, A; Savari, A; Movahedinia, A; Zare, P and Bagheri, T (2011). Serum biochemical change induced by *in vitro* sub chronic mercury chloride in yellowfin sea bream (*Acanthopagrus latus*). Iranica J. Eng. Environ., 2: 208-214.
- Joshi, BD (1980). Sex related variations in some

- hematologic values of certain freshwater teleosts. *Jpn. J. Ichthyol.*, 27: 139-143.
- Kori-Siakpere, O; Ake, JEG and Idoge, E (2005). Haematological characteristics of the African snakehead, *Parachanna obscura*. *Afr. J. Biotechnol.*, 4: 527-530.
- Mostaghni, K; Moghaddam Jafary, A and Badiei, K (2005). Effects of experimental proximal duodenal obstruction on clinical, hematological, serum and urine biochemical changes in sheep. *Iranian J. Vet. Res.*, 6: 1-8.
- Mulcahy, MF (1970). Blood values in the pike, *Esox lucius*. *J. Fish Biol.*, 21: 203-209.
- Orun, I; Dorucu, M and Yazlak, H (2003). Haematological parameters of three cyprinid fish species from Karakaya Dam Lake, Turkey. *Online J. Biol. Sci.*, 3: 320-328.
- Owolabi, OD (2011). Haematological and serum biochemical profile of the upside-down catfish, *Synodontis membranacea* Geoffroy Saint Hilaire from Jebba Lake, Nigeria. *Comp. Clin. Pathol.*, 20: 163-172.
- Ozretic, MK; Ozretic, B; Petrovic, S and Nikolic, T (2001). Seasonal variations of some blood parameters in farmed sea bass (*Dicentrarchus labrax* L.). *Perid. Biol.*, 103: 67-75.
- Pavlidis, M; Futter, WC; Kathario, P and Divanach, P (2007). Blood cells of six Mediterranean mariculture fish species. *J. Appl. Ichthyol.*, 23: 70-73.
- Rowley, AF; Hunt, TC; Page, M and Mainwaring, G (1988). Fish. In: Rowley, AF and Ratcliffe, NA (Eds.), *Vertebrate blood cells*. (1st Edn.), Cambridge, Cambridge University Press. PP: 19-127.
- Sahan, A; Altun, T; Cevik, F; Cengizler, I; Nevsat, E and Genc, E (2007). Comparative study of some haematological parameters in European eel (*Anguilla anguilla* L., 1758) caught from different regions of Ceyhan river (Adana, Turkey). *E.U.J. Fish. Aqua. Sci.*, 24: 167-171.
- Silveira, R and Rigores, C (1989). Características hematológicas normales de *Oreochromis aureus* en cultivo. *Rev. Latinoam. Acuic. Lima*, 39: 54-56 (abst. in English).
- Sowunmi, AA (2003). Haematology of the African catfish, *Clarias gariepinus* (Burchell, 1822) from *Eleiyele reservoir*, Ibadan, Southwest Nigeria. *Zool.*, 2: 40-44.
- Tavares-Dias, M; Moraes, FR and Imoto, ME (2008). Hematological parameters in two neotropical freshwater teleost, *Leporinus macrocephalus* (Anostomidae) and *Prochilodus lineatus* (Prochilodontidae). *Biosci. J.*, 24: 96-101.
- Vazquez, GR and Guerrero, GA (2007). Characterization of blood cells and haematological parameters in *Cichlasoma dimerus* (Teleostei, Perciformes). *Tissue Cell*. 39: 151-160.

Archive