

## Short Paper

## Naturally occurring ehrlichiosis in Egyptian dogs

Salem, N. Y. \*; Rakha, G. H. and Baraka, T. A.

Department of Clinical Sciences, Faculty of Veterinary Medicine, Cairo University, Giza, Egypt

\*Correspondence: N. Y. Salem, Department of Clinical Sciences, Faculty of Veterinary Medicine, Cairo University, Giza, Egypt. E-mail: nemovet2011@gmail.com

(Received 8 Jun 2013; revised version 25 Nov 2013; accepted 30 Nov 2013)

## Summary

Canine ehrlichiosis has emerged as one of the most clinically important tick-borne diseases affecting dogs. Eighty-five naturally infected dogs have been investigated, the most consistent clinical signs recorded were fever, emaciation and lymphadenopathy, anemia, monocytosis, thrombocytopenia, hypoalbuminemia, elevation in liver enzymes and total bilirubin were the most remarkable changes associated with canine ehrlichiosis in Egyptian dogs; microscopic examination failed to provide definitive diagnosis of canine ehrlichiosis. Season did not greatly influence the disease; the type of ticks involved in the disease transmission in Egypt was *Rhipicephalus sanguineus*.

**Key words:** Canine ehrlichiosis, Fever, Anemia, Liver enzymes, PCR

## Introduction

Canine ehrlichiosis has emerged as one of the most important infectious diseases affecting dogs (Moreira *et al.*, 2003); infecting mainly macrophages, monocytes and granulocytes (Jadhav *et al.*, 2011), transmitted chiefly by *Rhipicephalus sanguineus* (Troy and Forrester, 1990). The pathogenesis involves an 8-20 day incubation period and 3 clinical aspects: acute, subclinical and chronic phases (Alleman, 2005). Fever, weight loss, hepatosplenomegaly, lymphadenopathy and hemorrhage were described (Troy and Forrester, 1990). Thrombocytopenia, anemia and leucopenia are common hematologic abnormalities (Scorpio *et al.*, 2008). Hypoalbuminemia, hyperglobulinemia, elevated liver enzymes, hyperbilirubinemia have been reported (Rungsipat *et al.*, 2009; Stephanie *et al.*, 2010). Morulae are uncommonly detected and its absence cannot rule out the infection (Alleman, 2005). PCR has proven to be a more specific, sensitive and reliable tool for definitive diagnosis (Breitschwerdt *et al.*, 1998). The aim of this study is to describe naturally occurring ehrlichiosis in Egyptian dogs by assessment of clinical and laboratory findings as well as PCR detection of *Ehrlichia* spp. This study pinpoints the effect of the season and other risk factors on disease epidemiology in this area of the world.

## Materials and Methods

Eighty five dogs of different ages, sexes, and breeds were used in this study. The dogs were referred to Small Animal Medicine Teaching Hospital, Faculty of Veterinary Medicine, Cairo University. Signs were recorded; complete physical examination and clinical hematology were done. Serum samples were analysed

for AST, ALT, ALP, TP, albumin, total bilirubin, BUN and creatinine with respective kits (Stanbio® Inc., USA), ticks were collected and identified (Nuttal *et al.*, 1908). Blood films were examined under oil immersion lens.

DNA was extracted from whole blood using Blood DNA Preparation Kit (Jena Bioscience GmbH, Jena, Germany) according to manufacturer's instructions. Purified DNA was tested with primers that amplify a portion of 16S RNA gene.

ECC (5'AGAACGAACGCTGGCGGCAAGC-3') and ECB (5'CGTATTACCGCGGCTGCTGGCA-3') primers amplified all *Ehrlichia* spp. (Dawson *et al.*, 1996); PCR cycle was performed according to Murphy *et al.* (1998), and a previously confirmed sample was used as a positive control.

Student's t-test (STATISTICA for Windows, version 5.1, StatSoft, Inc.) was used.

## Results

Dogs were divided into 4 groups: <1 year (3/85, 3.52%), between 1-<3 years (15/85, 17.64%), between 3-<5 years (49/85, 57.64%) and >5 year (18/85, 21.17%). Male (46/85, 54.11%) and female (39/85, 45.88%) were almost equally infected.

German shepherd dogs were the most affected breed 50/85 (58.82%), Labrador 12/85 (14.11%) and the other breeds 23/85 (27.05%). The seasonal results were: summer (27/85, 31.76%), autumn (21/85, 24.70%), winter (17/85, 20.01%), and spring (20/85, 23.52%). The ticks were identified as *Rhipicephalus sanguineus* according to morphologic characteristics. The most consistent signs (Fig. 1) were pyrexia (67/85, 78.82%), pale mucous membrane (39/85, 45.88%), presence of ticks at time of admission (54/85, 63.52%) Fig. 1A, hepatosplenomegaly (26/85, 30.58%), lymphadenopathy



**Fig. 1:** 1A) Severe tick infestation in head of Great Dane infected dog. 1B) Severe emaciation in German shepherd infected dog. 1C) Yellowish discoloration in oral mucosa of Rottweiler infected dog. 1D) Analysis of PCR products by 1.5% agarose gel electrophoresis and ethidium bromide staining showing the positive samples after the primary amplification cycle, DNA ladder (100 bp DNA ladder, Jena Bioscience, Jena, Germany) was loaded with the samples simultaneously

**Table 1:** Hematologic and serum biochemical findings of 85 dogs with ehrlichiosis<sup>1</sup>

Parameter	Patient data	Control data <sup>a2</sup>
RBCs ( $\times 10^6/\mu\text{l}$ )	4.7 $\pm$ 0.32*	7.41 $\pm$ 1.19
Hemoglobin (g/dl)	12.06 $\pm$ 0.6*	15.74 $\pm$ 2.17
PCV (%)	35.69 $\pm$ 6.35*	44.14 $\pm$ 6.14
MCV (fl)	76.62 $\pm$ 14.7*	69.14 $\pm$ 6.15
MCH (pg)	22.85 $\pm$ 2.3	23.10 $\pm$ 2.02
MCHC (%)	32 $\pm$ 3.77	33.56 $\pm$ 0.77
WBCs ( $\times 10^3/\mu\text{l}$ )	8.354 $\pm$ 2.69*	11.850 $\pm$ 1.16
Neutrophil ( $\times 10^3/\mu\text{l}$ )	3.574 $\pm$ 0.393*	6.728 $\pm$ 0.19
Lymphocyte ( $\times 10^3/\mu\text{l}$ )	3.612 $\pm$ 0.340	3.928 $\pm$ 0.27
Monocyte ( $\times 10^3/\mu\text{l}$ )	1.050 $\pm$ 0.124*	0.775 $\pm$ 0.23
Platelets ( $\times 10^3/\mu\text{l}$ )	103.93 $\pm$ 16.67*	270.93 $\pm$ 14.12
Total protein (g/dl)	7.75 $\pm$ 0.51*	9.66 $\pm$ 0.48
Albumin (g/dl)	2.256 $\pm$ 0.65*	4.38 $\pm$ 0.66
ALT (u/l)	118.84 $\pm$ 24.00*	48.87 $\pm$ 7.25
AST (u/l)	124.82 $\pm$ 13.8*	35.103 $\pm$ 6.346
ALP (u/l)	213.21 $\pm$ 15.2*	90.159 $\pm$ 17.912
BUN (mg/dl)	21.004 $\pm$ 8.27*	11.93 $\pm$ 3.65
Creatinine (mg/dl)	0.99 $\pm$ 0.408	1.07 $\pm$ 0.31
Total bilirubin (mg/dl)	0.909 $\pm$ 0.4*	0.46 $\pm$ 0.15

\* Show statistically significant difference at P<0.05. <sup>1</sup> From Small Animal Internal Medicine Teaching Hospital, Faculty of Veterinary Medicine, Cairo University, and <sup>2</sup> A control data was taken from healthy dogs in the same locality

(38/85, 44.70%), emaciation (17/85, 20%) Fig. 1B, icterus (2/85, 2.35%) Fig. 1C, epistaxis (2/85, 2.35%), polyarthritis (3/85, 3.52%), peripheral edema (1/85, 1.17%), morulae were detected in 5/85 (5.88%) films.

The mean hematologic and serum biochemistry values are shown in Table 1.

Reduction in RBCs, HB content, PCV percentage with normocytic normochromic anemia, leucocytic count, and neutrophils along with increase in monocytes count were observed and 95% of dogs had thrombocytopenia. Significant increase in ALP, ALT, AST with hypoalbuminemia was observed. 488 bp were amplified as shown in Fig. 1D.

### Discussion

Canine ehrlichiosis is a widely known tick-borne disease. The highest occurrence was at 3-5 years followed by dogs over 5 years; Poitout *et al.* (2004) found the highest prevalence in dogs between 4-13 years. German shepherd was the most affected breed; the defective cell-mediated immunity may be associated with the severity of the disease in this breed (Nyindo *et al.*, 1980).

Seasonal similarity was found, although Mosallanejad *et al.* (2010) reported the highest prevalence was in summer. The global climatic changes made ticks more adaptive (Leschnik *et al.*, 2008); the dynamicity of ticks depends on the climatic condition followed by changes in seasonal patterns (Friedhoff, 1988). The weather in Egypt is warm throughout the year, which may have an impact on the pattern in Egypt.

Ehrlichiosis causes multisystem involvement, the organism multiplies inside mononuclear cells spreading to liver, spleen and lymph nodes (Woody and Hoskins, 1991). These circulating infected MNCs adhere to vascular epithelium inducing vasculitis and subendothelial tissue infection (Iqbal and Rikihisa, 1994). Fever and leucopenia were common in canine ehrlichiosis and should be added as differential diagnosis of pyrexic dogs (Unver *et al.*, 2005).

Morulae were detected in 5/85 (5.88%); the detection is ultimate for diagnosis, but it's seldom found and mostly unrewarding (Alleman, 2005).

Reduction in RBCs, HB content, and PCV percentage with normocytic normochromic anemia have been recorded. *Ehrlichia* is thought to cause bone marrow destruction leading to mild to severe non-regenerative anemia (Rungsipat *et al.*, 2009). Davoust *et al.* (1996) believed the anemia was due to immune-mediated destruction of RBCs, as antibodies would be fixed at the membrane causing their lyses by effector cells of the immune system.

Decrease in leucocytes count and neutrophils were observed, and the depletion of granulocytic precursors or damage of myeloid cells may be implicated (Codner and Farris-Smith, 1986). Monocytosis has been documented in ehrlichiosis (De Castro *et al.*, 2004); this elevation is expected in inflammatory disease with high demand for macrophages, hemolytic or hemorrhagic and immune-mediated diseases (Raskin *et al.*, 2004).

Hypoproteinemia and hypoalbuminemia were detected along with blood loss or decreased protein production due to concurrent mild liver disease. Reardon and Pierce (1981) observed development of many expanding foci in hepatic sinusoids, with the compression leading to necrosis of adjacent hepatocytes with increased enzymes and bilirubin activities.

Elevated BUN levels were observed in patients without similar rise in creatinine levels; catabolic state is likely to occur in anemic patients due to tissue hypoxia, and pyrexia (Michel *et al.*, 1997).

PCR detection of *Ehrlichia* is possible; recent studies showed efficacy of PCR in *Ehrlichia* detection (Iqbal and Rikihisa, 1994). The difference between direct microscopic examination and PCR revealed the low capability of microscopic examination in establishing a definitive diagnosis (Lakshmanan *et al.*, 2007).

This study showed season did not greatly influence the disease; the type of tick involved was *Rhipicephalus sanguineus*. Pyrexia, anemia, leucopenia, thrombocytopenia, and elevated liver enzymes are the base for tentative diagnosis but for definitive diagnosis, PCR is found to be more efficient.

## Acknowledgements

The authors thank the members of the Biotechnology Research Unit (ARRI), Egypt, for supplying facilities for molecular analysis. Special thanks to Dr. H. Farag, assistant lecturer of infectious disease for his help in tick identification.

## References

- Alleman, AR** (2005). The diagnosis and treatment of tickborne disease in dogs. In: *Proceedings of the North American Veterinary Conference* (NAVC, 2005), Orlando, Florida, USA. PP: 472-477.
- Breitschwerdt, EB; Hegarty, BC and Hancock, SI** (1998). Sequential evaluation of dogs naturally infected with *Ehrlichia canis*, *Ehrlichia chaffeensis*, *Ehrlichia equi*, *Ehrlichia ewingii*, or *Bartonella vinsonii*. *J. Clin. Microbiol.*, 36: 2645-2651.
- Codner, EC and Farris-Smith, L** (1986). Characterization of the subclinical phase of ehrlichiosis in dogs. *J. Am. Vet. Med. Assoc.*, 189: 47-50.
- Davoust, B; Parzy, D; Pubert, D; Martet, G; Deparis, X and Ott, D** (1996). Hematological symptoms of acute canine ehrlichiosis. *Rev. Méd. Vét.*, 147: 69-74.
- Dawson, JE; Biggie, KL; Warner, CK; Cookson, K; Jenkins, S; Levine, JF and Olson, JG** (1996). Polymerase chain reaction evidence of *Ehrlichia chaffeensis*, an etiologic agent of human disease, in dogs from southeast Virginia. *Am. J. Vet. Res.*, 57: 1175-1179.
- De Castro, MB; MacHado, RZ; De Aquino, LPT; Aless, AC and Costa, MT** (2004). Experimental acute canine monocytic ehrlichiosis: clinico-pathological and immunopathological findings. *Vet. Parasitol.*, 119: 73-86.
- Friedhoff, KT** (1988). *Transmission of Babesia. Babesiosis of domestic animals and man*. 1st Edn., Boca Raton, USA, CRC Press. PP: 23-52.
- Iqbal, Z and Rikihisa, Y** (1994). Re-isolation of *Ehrlichia canis* from blood and tissues of dogs after doxycycline treatment. *J. Clin. Microbiol.*, 32: 1644-1649.
- Jadhav, RK; Kumari, RR; Javed Jameel, A and Kumar, P** (2011). Emergence of arthropod transmitted infections in kennel dogs. *Vet. World*. 4: 522-528. doi: 10.5455/vetworld.2011.522-528.
- Lakshmanan, B; John, L; Gomathinayagam, S and Dhinakarraj, G** (2007). Molecular detection of *Ehrlichia canis* from blood of naturally infected dogs in India. *Vet. Archiv.*, 77: 165-166.
- Leschnik, M; Kirtz, G; Tichy, A and Leidinger, E** (2008). Seasonal occurrence of canine babesiosis is influenced by local climate conditions. *Int. J. Med. Microbiol.*, 298: 243-248.
- Michel, KE; King, LG and Ostro, E** (1997). Measurement of urinary nitrogen content as an estimate of the amount of total urinary nitrogen loss in dogs in intensive care unit. *JAVMA*. 210: 356-359.
- Moreira, SM; Bastos, CV; Araújo, MS and Passos, LMF** (2003). Estudo retrospectivo (1998-2001) da erliquiose canina em Belo Horizonte [Retrospective study (1998-2001) of canine ehrlichiosis in Belo Horizonte]. *Arquivo Brasileiro de Medicina Veterinária e zootecnia*. 55: 141-147.
- Mosallanejad, AR; Jalali, BR and Alborzi, MH** (2010). Sero-prevalence of *Ehrlichia canis* in dogs referred to veterinary hospital of Shahid Chamran University of

- Ahvaz, Iran. Archiv. Razi Inst., 65: 21-26.
- Murphy, GL; Ewing, SA; Whitworth, LC; Fox, JC and Kocan, AA** (1998). A molecular and serologic survey of *Ehrlichia canis*, *Ehrlichia chafeensis* and *E. ewingii* in dogs and ticks from Oklahoma. Vet. Parasitol., 79: 325-339.
- Nuttal, GHF; Warburton, C; Cooper, WF and Robinson, LE** (1908). *Ticks. Part 1. A monograph of the Ixodoidea*. England, Cambridge University Press. PP: 1-104.
- Nyindo, M; Huxsoll, DL; Ristic, M; Kakoma, I; Brown, JL; Carson, CA and Stephenson, EH** (1980). Cell mediated and humoral immune response of German shepherd dogs and Beagles to experimental infection with *Ehrlichia canis*. Am. J. Vet. Res., 41: 250-254.
- Poitout, MF; Shinozaki, JK; Stockwell, PJ; Holland, CJ and Shukla, SK** (2004). Genetic variants of *Anaplasma phagocytophilum* infected dogs in western Washington State. J. Clin. Microbiol., 43: 796-801.
- Raskin, RE; Latimer, KS and Tvedton, H** (2004). Leucocyte disorder. In: Willard, MD and Tvedten, H (Eds.), *Small animal clinical diagnosis by laboratory methods*. (4th Edn.) St. Louis, Missouri, W. B. Saunders Co., PP: 63-91.
- Reardon, MJ and Pierce, KR** (1981). Acute experimental canine ehrlichiosis. Sequential reaction of the hemic and lymphoreticular systems. Vet. Pathol. J., 18: 48-61.
- Rungspipat, A; Oda, M; Kumpoonsiri, N; Wangnaitam, S; Poosoontham, R; Komkaew, W; Suksawat, F and Ryoji, Y** (2009). Clinicopathological study of experimentally induced canine monocytic ehrlichiosis. Vet. Clin. Pathol., 18: 13-22.
- Scorpio, DG; Wachtman, LM; Tunin, RS; Barat, NC; Garyu, JW and Dumler, JS** (2008). Retrospective clinical and molecular analysis of conditioned laboratory dogs with serologic reactions to *Ehrlichia canis*, *Borrelia burgdorferi* and *Rickettsia rickettsii*. J. Am. Assoc. Lab. Anim. Sci., 47: 23-28.
- Stephanie, NG; Lim, S; Daub, A; Saini, SS and Kaushik, AK** (2010). Benign mononuclear gammopathy in a dog exhibiting a low antibodies titer to *E. canis*. J. Clin. Immunol. Immunopathol. Res., 2: 9-14.
- Troy, GC and Forrester, SD** (1990). *Ehrlichia canis*, *E. equi* and *E. risticii* infections. In: Greene, GE (Ed.), *Infectious diseases of the dog and cat*. (1st Edn.), Philadelphia, W. B. Saunders Co., PP: 48-59.
- Unver, A; Rikihisa, Y; Borku, K; Ozkanlar, Y and Hanedan, B** (2005). Molecular detection and characterization of *Ehrlichia canis* from dogs in Turkey. Berl. Munch. Tierarztl. Wochenschr., 118: 300-304.
- Woody, BJ and Hoskins, JD** (1991). Ehrlichial diseases of dogs. Vet. Clin. North Am. Small Anim. Pract., 21: 75-98.

Archive of SID

مقاله کوتاه:

## وقوع طبیعی ارلیشیوزیس در سگ‌های مصری

نوحا یوسف سالم، جمال حسن رخا و طاهر احمد باراکا

گروه علوم درمانگاهی، دانشکده دامپزشکی دانشگاه قاهره، جیزه، مصر

(دریافت مقاله: 18 خرداد 1392، پذیرش نهایی: 9 آذر 1392)

ارلیشیوزیس سگ‌ها به عنوان یکی از مهمترین بیماری‌های منتقل شونده از کنه به سگ‌ها شناخته شده است. 85 سگ آلوده شده طبیعی مورد بررسی قرار گرفتند. پایدارترین علائم بالینی ثبت شده عبارت بود از تب، لاغری و لنفادنوپاتی. کم خونی، افزایش مونوسیت‌ها، کاهش پلاکت‌ها، کاهش آلبومین خون، افزایش آنزیم‌های کبدی و بیلیروبین کل، برجسته‌ترین تغییرات مرتبط با ارلیشیوزیس سگ‌های مصری بود. بررسی‌های میکروسکوپی در تشخیص قطعی بیماری ارلیشیوزیس سگ‌ها مؤثر باشد. فصل تأثیر زیادی بر بیماری نداشت. کنه عامل انتقال این بیماری در مصر ریپی سفالوس ساگینوس بود.

**واژه‌های کلیدی:** ارلیشیوزیس سگ‌ها، تب، کم خونی، آنزیم‌های کبدی، واکنش زنجیره‌ای پلیمرز