The Effect of Different Deheating Processes on Residual Myrosinase Activity, Antimicrobial Properties and Total Phenolic Contents of Yellow Mustard (*Sinapis alba*)

M. Mizani^{a*}, M. Yousefi^b, S. Rasouli^c, A. Sharifan^d, M. Bamani Moghaddam^e

^a Associate Professor of the Department of Food Science and Technology, Science and Research Branch, Islamic Azad University, Tehran, Iran.

^b Instructor of the Department of Food Science and Technology, Science and Research Branch, Islamic Azad University, Tehran, Iran.

^c Associate Professor of the Departement of Nanomaterials and Nanocoatings, Institute for Color Science and Technology (ICST), Tehran, Iran.

^d Assistant Professor of the Department of Food Science and Technology, Science and Research Branch, Islamic Azad University, Tehran, Iran.

^e Associate Professor of the Department of Statistics, Faculty of Mathematical Sciences and Computer, Allameh Tabatabai University, Tehran, Iran.

Received: 12 September 2015

Accepted: 17 March 2016

ABSTRACT: Mustard is a natural multi-functional additive with strong hot flavor. In this research, powdered yellow mustard (*Sinapis alba*) was heated by 21 different thermal processes, statistically designed by Response Surface Methodology to evaluate the effects of combinations of heating media (hot air, hot water and steam) and heating time (0-20 min) on residual myrosinase activity, the minimum inhibitory concentration (MIC), color characteristic and also the major phenolic constituents of each treated sample. The optimum process, for the desirable function was 5 min of heating by hot air. Under optimal conditions, myrosinase activity was reduced by 44%, MIC and lightness (L*) were approximately unchanged, redness (a*) and yellowness (b*) were increased as compared to the control. In spite of the fact that the total phenolic content were reduced severely (>90%) due to deheating processes, but the chromatography spectrum indicated that some compound (caffeic acid) were newly formed or existed in higher concentration (Sinapine) particularary in steam treated samples.

Keywords: Phenolic Component, Thermal Heating, Yellow Mustard.

Introduction

Cruciferous vegetables are vegetables of the family Brassicaceae (also called Cruciferae). The family contains over 330 genera and about 3,700 species such as cabbage, broccoli, cauliflower, turnip, rapeseed, mustard, radish, horseradish, wasabi and watercress. Mustard is the third most commonly consumed spice, after salt and pepper. In the second half of the twentieth century, world usage of mustard more than doubled, from 75,000 tons to over

170,000 tons and it becomes more popular spice in fast-food restaurants. It is usually found in three types: yellow or white mustard (Sinapis alba), brown mustard (Sinapis nigra) and oriental (Sinapis juncea) (Smith & Hong, 2003). White mustard is the most widely used- type of mustard in food industry and contains valuable active components including isothiocyanates, complex polysaccharides (mucilagenous substance), phenolic compounds, dithiolthiones & dietary fibers. Sinalbin is a thiocvanate glycoside that exists in white mustard, which is hydrolyzed by myrosinase

^{*}Corresponding Author: mizani1_2000@yahoo.com

to form a flavor- active Isothiocyanate. Isothiocyanates is responsible for the known "hot" flavor of white mustard and also its antimicrobial and preservative function (Lin Glucosinolate et al., 2000). and/or isothiocyanates may display diverse and interesting biological properties. Some are hepatotoxic or goitrogenic, whereas others anti-bacterial, anti-fungal, have antinematocidal and protozoal. or anticarcinogenic activities (Herzallah & Holley, 2012). Thiocyanate and oxazolidinethione anions are known to compete with iodine in two ways, by inhibiting its uptake through competition the sodium-iodide with symporter (the common term for combined transporters), and binding of iodine to tyrosine residues of thyroglobulin at high concentrations. Hence one of the common symptoms of glucosinolate and /or isothiocyanate exposure is the impairment of thyroid function, resulting the in а hypertrophy of this endocrine gland (goiter). The toxicity of glucosinolate is generally attributed the isothiocvanates. to thiocyanate, oxazolidinethione and nitriles, originating from enzymatic cleavage of glucosinolates by myrosinase. In addition to the above mentioned negative aspects of myrosinase activity, it has also positive functions due to antimicrobial activities. The mechanism of this antimicrobial action is uncertain, and different theories have been approved. Turgis et al. (2009), suggested that the antibacterial activity of mustard powder is probably due to the ability of its components (i.e. isothiocyanates) to damage the microorganism cell membrane, lysing and releasing of the outer membraneassociated materials especially ATP from the cells. Investigations of Nilson & Holley, (2012), showed that isothiocyanates may sulfhydryl the groups bind to of microorganism cell enzymes and interfere in its functions. Another valuable component of mustard is phenolic compounds which are main source of antioxidant activity through

2

acting as chain breakers and free radical scavenger (Shahidi *et al.*, 1995). Unfortunately this activity may be severely loss during different processing (Amin & Yee Lee, 2005).

The main objective of this research was to apply 21 different statistically designed thermal processes on white mustard powder and evaluate the positive effects (myrosinase partial inactivation: hot flavor and toxicity reduction) and also the negative effects (loss of antimicrobial and the total phenol contents) of each thermal treatment and finally determine the optimum deheating condition by statistical analysis through response surface methodology (RSM) and desirability function.

Materials and Methods

- Materials

Yellow mustard powder (code no: 201) were provided by Pars Siavashan, the local agent of G.S.DUNN (Ontario, Canada). Four microorganisms including *Bacillus subtilis* (PTCC 1254), *Bacillus coagulans* (NBRC 12583), *Saccharomyces cerevisiae* (PTCC 5269) and *Zygosaccharomyces rouxi* (PTCC 5206) were provided by Iranian Research Organization for Science and Technology (IROST). Sabrose Dextrose agar (SDA) and Nutrient agar (NA) (Merck KGaA) were used as growth media for yeasts and bacteria, respectively. Glucose Assay Kit was provided by Zist-Chime (Tehran, Iran)

- Methods

- Heating processes

Mustard powder was spread in a thin (2cm) layer in an aluminum pan. Three different thermal enzyme inactivation media, hot air, hot water and steam, have been used by heat processing of the samples in oven, water bath and autoclave, respectively. All of the processes applied at the same temperature (100 ± 5 °C) for (0 -20 min) (Eapen *et al.*, 1968; Herzallah & Holley, 2012). Different combinations of heating

time and methods were designed by response surface methodologies, Design-Expert 7.0.0 software (Table 1).

- Residual myrosinase activity

10 g of the heated mustard powder was homogenized in 90 ml distilled cold water. The homogenate was then centrifuged at 3000g for 15 min at 4°C (Wang *et al*, 1994). Glucose assay kit was used to determine the residual myrosinase activity in supernatant according to the amount of released glucose by UV spectrophotometer at 500nm (Lin *et al.*, 2000).

- Minimum inhibitory concentration

Antimicrobial activity of the heated mustard samples were studied according to their minimum inhibitory concentrations (MIC). Surface culture was accomplished in the specific culture media for the selected yeasts and bacteria, and then 4 holes were created in each plate. Suspensions of natural mustard powder and the heated mustard samples were prepared with different. concentrations (0.5, 1, 1.5, 2, 2.5, 3, and 3.5 %) and injected in the holes. The plates were incubated for 24-48 h at 37 °C, and then the minimum inhibitory concentrations of the treated and untreated samples on the microorganisms were determined (Saadoun et al., 2007).

- Color characteristics

Color of the heated samples were studied by CS 2000 spectroradiometer (KonicaMinolta, Japan) at 380 -780 nm using D65 illuminant. Color characteristics of each treated sample were described in L*, a*, and b*values which are coordinates in visual uniform color space and are related to x, y, z tristimulus values by the following equations:

 $L^{*}= 116 (Y / Yn) 1/3 - 16$ a^{*} = 500 [(X / Xn) 1/3 - (Y / Yn) 1/3] b^{*} = 200[(Y / Yn) 1/3 - (Z / Z n) 1/3]

These factors have been specified by the International Committee of Illumination and were calculated by the software system (CS-S10W).

- Total concentration of phenolic compounds Phenolic compounds have been directly extracted from the heated samples (2 gr) by means of 5 ml methanol/water 80:20 (V/V) solution. The suspensions were homogenized in the ultrasonic bath for 15 min at room temperature, and then centrifuged at 5000 g for 25 min. An aliquot of the supernatant was filtered and analyzed by HPLC (Young Lin Acme 2000) according to International Olive Council (2009).

- Statistical analysis

Statistically the design of thermal treatments was accomplished by response surface methodologies, Design-Expert 7.0.0 software and one- way analysis of variance (ANOVA) ($p \le 0.05$) was applied to analyse the results obtained.

Run	1	2	3	4	5	6	7	8	9	10	11
Heating Media	Steam	Hot air	Hot water	Hot water	Hot air	Steam	Steam	Hot air	Steam	Hot air	Hot air
Heating Time (min)	5	20	12.5	20	5	16.25	12.5	5	5	12.5	16.25
Run	12	13	14	15	16	17	18	19	20	21	
Heating Media	Hot water	Hot water	Hot air	Hot water	Steam	Hot air	Steam	Hot water	Steam	Hot water	
Heating Time (min)	5	5	20	16.25	20	8.75	20	8.75	8.75	20	

Table 1. Different heating processes statistically designed by RSM method

Results and Discussion

- Residual myrosinase activity

The results of the residual enzymatic activity of the natural mustard powder and different treated samples are shown in Table 2. All of the samples showed a decreasing pattern of enzymatic activity with increasing time. The most and the least effective heating media were steam and hot water, respectively. This might be expected, because of higher heating transfer rate of steam (6000-15000W/m^{2°}C) as compared to hot water (1000-6000 W/m^{2°}C) (Kreith, 2000).

Although heat transfer coefficient for hot water is substantially higher than hot air (10- $100 \text{ W/m}^{2^{\circ}}\text{C}$), but the results of this research showed lesser effect on the enzyme inactivation. It seems that forced air

circulation in laboratory oven (Memmert universal oven, model UF55) might increase the heating rate of hot air medium and results in reducing the enzyme activity more severely as compared to still hot water method (Yousefi et al., 2014). In the beginning stages (up to 5 minutes) of heating by hot water, an increasing pattern of enzyme activity (from 934.20 to 1152.24 µg glucose released / min) might be observed (Table 2). The same result has reported by Diogo, 2010 for been inactivation myrosinase in broccoli (Brassica oleracea var. Italica). It has been reported that myrosinase activity of the low moisture content (<13%) broccoli increased at the moderate temperatures $(40-60^{\circ}C)$ and decreased by further heating up to the higher

Heating media	Heating time	Residual enzymatic activity	Enzymatic activity
ficuring incur	(min)	(μg glucose released / min)	reduction (%)
	0	934.20±15.23°	
	5	657.02±11.70 °	29
Steam	8.75	516.7±20.21 ^e	44
	12.5	411.99±15.01 ^{fg}	55
	16.75	198.41±13.90 ⁱ	78
	20	101.80±10.01 ^j	89
	0	934.20±15.23°	
	5	1157.74±7.00 ^a	-23
Hot water	8.75	$948.02{\pm}11.90^{b}$	-1
	12.5	594.5±9.70 ^d	36
	16.75	$409.67 \pm 3.42^{\text{ fg}}$	56
	20	388.13±13.02 ^g	58
	0	934.20±15.23 ^b	
	5	533.54±5.70 ^e	42
Hot air	8.75	443.04±9.21 ^f	52
	12.5	374.91±7.51 ^g	59
	16.75	296.32±7.01 h	68
	20	223.07±8.80 ⁱ	76

 Table 2. Residual enzyme activities of different heat-treated mustard powder samples

temperatures. Therefore it seems that because of lower heating rate of hot water, mustard powder might be heated slowly in this method that provide an opportunity for activating the enzyme at the initial lower temperatures. Wang et al. (1994) studied the heating rate of fruits in exposure to hot air and hot water and reported that the circulation rate of the heating medium is an effective factor but it is more important for hot air processes. Therefore this parameter might be the reason of the steeper slope for hot air blanching as compared to hot water, in spite of its lower heat transfer coefficient. Eapen et al. (1968) investigated on thermal inactivation of myrosinase from canola seeds and showed that the residual activity was more than 70% after oven heating for 30 min at 105°C. According to the results of Table 2, the inactivation rate was much higher in the present research therefore myrosinase of mustard powder retained its activity about 25% after being heated in the oven for 20 min. Therefore source of the enzyme is an important factor that might cause differences in thermal resistance of the enzyme and also in thermal conductivity and/or internal heat resistance of the product. This result is in agreement with the report by Ghawi *et al.* (2012) who observed lower heat stability of myrosinase in mustard than in canola. Another main reason for this difference is related to the particle size of the product (Wang *et al.*, 1994) that leads to slower heat penetration rate into the particles of whole canola seeds as compared to mustard fine powder.

- Minimum inhibitory concentrations (MICs)

The MICs of different treated mustard samples obtained for *Bacillus subtilis*, *Bacillus coagulans*, *Saccharomyces cerevisiae* and *Zygosaccharomyces rouxi* are given in Table 3.

Dun	Microorganisms species					
Kun -	Bacillus subtilis	Bacillus coagulans	Saccharomyces cerevisiae	Zygosaccharomyces rouxi		
1	_*		1.5	3		
2	-		3	-		
3	-	-	1.5	3.5		
4	-	-	2	-		
5			1.5	3.5		
6		-	3	-		
7	-	-	2	-		
8		-	1.5	3.5		
9	-	-	1.5	3		
10	-	-	2	-		
11	_	-	2.5	-		
12	3.5	3.5	0.5	2		
13	3.5	3.5	0.5	2		
14	-	-	3	-		
15	-	-	2	-		
16	-	-	-	-		
17	-	-	1.5	3.5		
18	-	-	-	-		
19	3	3.5	1	2.5		
20	-	-	1.5	3.5		
21	-	-	2	_		

 Table 3. Minimum Inhibitory Concentration (%) for different treated mustard samples*

*MIC for the untreated mustard powder against the above mentioned microorganisms were as follows: *Bacillus subtilis* (-), *Bacillus coagulans* (-), *Saccharomyces cerevisiae* (1%), *Zygosaccharomyces rouxi* (3%). *without inhibitory effect (MIC>3.5%).

As it may be observed, only three of the samples (Runs 12, 13 and 19) showed antimicrobial activity against bacterial strains with MICs 3 and 3.5% for Bacillus subtilis and Bacillus coagulans, respectively, whereas yeasts were more sensitive and inhibited with lower concentrations of mustard (0.5-3%). Shofran et al. (1998), and Nilson & Holley, (2012) studied the antimicrobial effect of mustard on different microorganisms and reported higher activity against yeasts as compared to the bacterial species. Lengthening the heating time (from 5 to 20min) of each method (hot air, steam or hot water) resulted in severe degradation of antimicrobial compounds and therefore higher inhibitory effect was obtained. The results of antimicrobial activity (Table 3) and remained enzymatic activity (Table 2) confirm each other, therfore deheated mustard samples with the same amount of residual enzymatic activity (such as runs 7, 10 and 15 where 400 µg glucose / min was released) showed equal antimicrobial activity against the same strain (i.e. Saccharomyces cerevisiae with MIC=2%). This might be expected because the antimicrobial activity of the samples is actually related to the amount of benzyl isothiocyanate and this compound is the product of enzymatic degradation of glucosinolates. Therefore by decreasing the residual enzymatic activity or, in other words, by lowering the amount of benzyl isothiocyanate, the antimicrobial activity will be reduced. The same pattern might be observed among other samples such as runs 12, 13 and 19. This result is in agreement with Luciano et al. (2011) who investigated the antimicrobial effect of both hot (natural enzyme) and cold (reduced enzyme activity) mustard on E.coli and reported that by heating mustard and reducing its enzyme activity, the antimicrobial effect might be decreased. Lara -Lledo et al. (2012) studied

the antimicrobial activity of mustard powder and sinigrin, and approved that mustard powder is more active than sinigrin because of presence of myrosinase in mustard. According to the results of Table 3, higher concentrations of the treated samples (MICs) might be necessary inactivate to Zygosaccharomyces rouxi than Saccharomyces cerevisiae. It means the latter yeast is more sensitive. This result is in agreement with the investigation of Shofran et al. (1998).

- Color characteristics

Values of three main color characteristics (L*, a* and b*) of 21 treated mustard samples are presented in Table 4.

All of the heating methods resulted in changing the color of the samples significantly and made them darker as compared to the control. Among different heating media, steam showed the highest effect on color quality of mustard and caused approximately 19% decrease in L* and 1.8% increase in b* after 20min heating of mustard powder. However data of the redness (a*) of the samples were very interesting. Therefore this parameter in runs 16, 18 is about 6 times greater than the control. These results confirmed more intense heating by steam. It seems that decreasing L* and increasing a* and b* during heating might be the result of undergoing of millard reactions in the presence of mustard proteinaceous compounds (24.94%) (dwb)and carbohydrates (39.49%) (dwb)and producing brown nitrogen-containing high molecular weight pigments, melanoidins (USDA, 1976; Bastos et al., 2012).

- Total phenolic contents

The total phenolic contents of heattreated samples of mustard powder have been shown in Table 5.

Run	a*	b*	L*
1	3.04±0.15 °	34.41±0.01 ^{ij}	67.82±1.28 ^{defg}
2	$1.08\pm0.06^{\text{hi}}$	$35.35 \pm 0.03^{\text{f}}$	$72.04 \pm 0.42^{\text{ abcde}}$
3	$2.03{\pm}0.03^{g}$	36.65 ± 0.01^{ab}	$66.02 \pm 0.86^{\text{fgh}}$
4	3.24±0.01 ^e	36.72±0.02 ^a	63.18 ± 0.08 ^{ghi}
5	-0.61 ± 0.07^{m}	$34.14{\pm}0.06^{kl}$	74.39±0.67 ^a
6	5.15±0.02 ^b	35.48 ± 0.03^{e}	$63.54 \pm 0.75^{\text{ ghi}}$
7	4.3±0.00 °	34.53 ± 0.03^{i}	65.61 ± 0.75 fgh
8	-0.61 ± 0.07^{m}	34.08 ± 0.02^{1}	74.28 ± 0.29^{a}
9	3.04 ± 0.15^{e}	34.14 ± 0.06^{kl}	68.04 ± 3.02^{defg}
10	-0.31 ± 0.01^{1}	34.71 ± 0.02^{h}	73.21±0.05 ^{abc}
11	-0.03±0.00 ^k	35.07±0.07 ^g	$72.78 \pm 0.09^{\text{ abcd}}$
12	0.87 ± 0.00^{ij}	$36.39 \pm 0.01^{\circ}$	68.81±2.39 ^{cdef}
13	0.89 ± 0.00^{ig}	36.39±0.01 °	69.1±0.84 ^{bcdef}
14	1.08 ± 0.06^{hi}	35.36±0.01 ^{ef}	72.31±1.21 ^{abcd}
15	2.42 ± 0.01^{f}	36.67±0.01 ^{ab}	64.83 ± 0.45 ^{fgh}
16	6.18 ± 0.01^{a}	35.67 ± 0.02^{d}	59.49±1.86 ⁱ
17	-0.55 ± 0.01^{m}	34.35±0.00 ^k	73.84±1.54 ^{ab}
18	6.19±0.00 ^a	35.67 ± 0.02^{d}	61.25±1.13 ^{hi}
19	1.19±0.02 ^h	36.56 ± 0.00^{b}	$67.74 \pm 1.42^{\text{defg}}$
20	3.6 ± 0.01^{d}	34.59 ± 0.02^{j}	67.12±1.35 ^{efg}
21	3.22 ± 0.01^{e}	$35.46{\pm}0.02^{a}$	$63.24 \pm 0.75^{\text{ghi}}$

J. FBT, IAU, Vol. 6, No. 2, 1-12, 2016

Table 4.	Color parameters	of treated	mustard	samples

* Color properties (a*, b* and L*) for untreated mustard powder were -0.85, 34.02 and 74.34, respectively.

Table 5. Total phenolic contents of different heat-treated mustard powder samples

Heating modia	Heating time	Total phenolic contents	Phenolic contents
neating media	(min)	(mg/kg)	reduction (%)
	0	55455.61±22.1 ^a	_
	5	4962.35 ± 44.0 ^{cd}	91.01
Steam	8.75	4688.22±25.4 °	91.54
	12.5	4584.16±17.5 °	91.70
	16.75	4431.49±5.7 ^{fg}	92.02
	20	4351.35±2.4 ^g	92.15
	0	55455.61±22.1 ^a	_
	5	5421.04 ± 12.8^{b}	90.22
Hot water	8.75	5324.01±31.9 ^b	90.03
	12.5	5058.28±3.1 ^{cd}	90.87
	16.75	4922.17 ± 17.0^{d}	91.12
	20	4621.91±16.4 ^e	91.66
	-		
	0	55455.61±22.1 ^a	_
	5	5358.22±22.7 ^b	90.33
Hot air	8.75	5090.95±14.0 ^c	90.82
	12.5	4732.43±112.6 ^e	91.46
	16.75	4629.36±17.2 ^e	91.56
	20	4561.23±67.5 ^{ef}	91.77

Untreated mustard powder had the highest content of phenolic compounds (55455.61 mg/kg). Mustard seed is usually considered as a rich source of polyphenolic

compounds (37000 - 46000 mg/kg) among different cultivars (Amin & Yee Lee, 2005; Ildiko *et al.*, 2006). As might be expected, in all of the blanching methods, a reduction in

M. Mizani et al.

the total phenolic contents of heat -treated mustard powder samples was observed. Therefore, the lowest and the highest reduction rates belonged to hot water (90.22%-91.66%) and steam (91.01%-92.15%) heat processes during (5-20 min). It has been previously shown that at the same temperature, wet (steam) heating methods had more severe effect on the reduction of total phenolic compounds as compared to dry (hot air) methods (Ghaderi Ghahfarokhi et al., 2011). But the interesting result of our research is between two different wet heating media (steam and hot water), hot water might act substantially milder than the other and even dry (hot air) heating medium. This result is positively coordinated with the reduction of myrosinase activity; therefore more phenolic contents might be saved in the samples with higher residual enzyme activity. It might be concluded that in blanching process, the circulation rate of the heating medium that affect its heat transfer coefficient should be adjusted to provide optimum enzymatic activity as well as the total phenolic contents.

- The effect of different heating processes on the phenolic compounds

Tables 6, 7 and 8 present the effect of different deheating processes on the phenolic compounds of white mustard powder.

Table 6. The effect of not air nearing process on the concentration (mg/kg) of different phenotic compo	Table 6. The effect of h	ot air heating process	s on the concentration	n (mg/kg) of d	lifferent phenolic	compounds
--	--------------------------	------------------------	------------------------	----------------	--------------------	-----------

Phenolic			Heating ti	ime (min)		
component	0	5.00	8.75	12.50	16.25	20.00
Hydroxytyrosol	653.89 ± 24.37^{a}	535.64±9.33 ^b	317.04±16.14°	211.50±10.54 ^d	89.92±9.67 ^e	19.09 ± 2.23^{f}
Tyrosol	$325.38{\pm}18.18^{a}$	$317.30{\pm}10.84^{a}$	322.44±6.49 ^a	287.94 ± 4.38^{b}	251.26 ± 8.80^{b}	208.15±12.56°
Sinapine	$2505.0{\pm}117.50^d$	2606.71 ± 50.8^d	2741.13±59.60 ^{dc}	$3008.4{\pm}34.40^{cb}$	$3228.49{\pm}55.0^{ba}$	3450.98±83.9 ^a
Caffeic acid	0 ± 0.00^{c}	0 ± 0.00^{c}	$0{\pm}0.00^{c}$	6.37±2.06 ^c	15.76 ± 3.06^{b}	51.39±4.36 ^a
Syringic acid	126.51±3.94 ^a	121.46 ± 5.95^{a}	67.86±5.33 ^b	23.95±1.62°	0 ± 0.00^{d}	$0{\pm}0.00^{d}$
Sinapic acid derivative	316.66±5.83 ^a	$306.88{\pm}7.84^{a}$	268.66±7.32 ^b	$244.65 {\pm} 9.38^{b}$	$207.17 \pm 8.00^{\circ}$	187.45±9.07 °
Flavanomarein	34.325±3.79ª	35.87 ± 4.33^{a}	32.45 ± 3.16^{ab}	30.47 ± 3.98^{abc}	20.27 ± 3.09^{bc}	18.40±2.43 °
Sinapic acid	66.270 ± 2.36^{a}	61.91±2.27 ^{ab}	58.91 ± 3.50^{ab}	57.75 ± 3.34^{ab}	52.62±3.12 ^b	50.63 ± 4.60^{b}

Values in same row with same superscript are not significantly (P<0.05) different.

Table 7. The effect of seam heating process on the concentration (hig/kg) of different phenome compounds							
Phenolic		Heating time (min)					I
component	0	5.00	8.75	12.50	16.25	20.00	ľ
Hydroxytyrosol	653.89±24.37ª	515.11b±3.85 ^b	314.07±6.85°	194.91±9.22 ^d	87.98±9.83 ^e	10.85 ± 2.19^{f}	
Tyrosol	$325.38{\pm}18.18^{a}$	$302.66{\pm}15.80^{a}$	$308.94{\pm}4.89^{a}$	285.17 ± 7.41^{ab}	239.46±11.95 ^{bc}	195.42±11.87 °	
Sinapine	2505.0±117.50 ^c	2496.34 ± 7.0^{0c}	2591.90±112.10 ^c	$2869.84 \!\pm\! 181.8^{bc}$	$3015.19 \!\pm\! 47.70^{ab}$	$3314.10{\pm}17.8^{a}$	
Caffeic acid	0±0.00 °	0±0.00 °	3.03±0.28 °	6.04±0.10 ^c	25.74±4.47 ^b	49.74±3.34 ^a	
Syringic acid	126.51±3.94 ^a	105.43 ± 7.06^{b}	63.58±7.71 ^b	9.11 ± 3.84^{d}	$0.51\!\pm\!0.73^d$	$0{\pm}0.00^{d}$	
Sinapic acid	216 66 5 928	204 25 1 5 91ab	277 55 16 10 ^b	210 12 + 0 44°	108 41 + 12 72° d	166 74+6 50 d	
derivative	510.00±3.85	304.33±3.81	277.35±0.19	219.13±9.44	198.41±12.72	100./4±0.39	
Flavanomarein	34.325±3.79 ^a	38.66 ± 2.05^{ab}	26.01 ± 1.58 bc	22.56±3.76 ^{cd}	17.38 ± 0.57^{cd}	14.16 ± 0.74^{d}	
Sinapic acid	66.270 ± 2.36^{a}	63.08±3.84 ^a	60.59±2.37 ^{ab}	51.05±1.52 bc	48.52±3.82°	43.42±2.12°	

Table 7. The effect of steam heating process on the concentration (mg/kg) of different phenolic compounds

Values in same row with same superscript are not significantly (P<0.05) different.

Phonolia component			Heating ti	me (min)		
r nenone component	0	5.00	8.75	12.50	16.25	20.00
Hydroxytyrosol	653.89±24.37 ª	626.56±9.11 ^a	518.46±15.67 ^b	421.59±15.86°	280.04±15.00 ^d	115.50±11.32 ^e
Tyrosol	325.38 ± 18.18 ^a	$335.43{\pm}10.47^{ab}$	324.16 ± 7.40^{ab}	$308.28{\pm}6.43^{ab}$	$277.21 \pm \! 17.09^{bc}$	249.32±14.73°
Sinapine	$2505.0{\pm}~117.50^{b}$	$2562.00{\pm}11.70^{b}$	$2535.8{\pm}8.00^{b}$	2621.78 ± 14.90^{b}	$2649.84{\pm}68.70^{b}$	$2889.83{\pm}14.8^{a}$
Caffeic acid	$0\pm0.00^{\circ}$	$0\pm 0.00^{\circ}$	$0\pm0.00^{\circ}$	$1.74 \pm 0.56^{\circ}$	12.04±2.39 ^b	19.96 ± 2.16^{a}
syringic acid	126.51±3.94 ^a	126.29±5.59 ^a	110.39± ^a	89.87±2.91 ^b	71.74±3.27 °	63.07±2.47 °
sinapic acid derivative	316.66±5.83 ^a	321.81±5.54 ª	314.65±6.05 ^a	$295.34{\pm}8.54^{ab}$	275.85±6.31 ^b	$245.19 \pm 7.46^{\circ}$
Flavanomarein	34.325±3.79 ^a	39.88±2.61 ^{ab}	37.49±0.75 ^{ab}	33.32±1.11 ^{a b}	28.47±1.65 ^{bc}	20.38±3.40°
sinapic acid	66.270 ± 2.36^{a}	67.22 ± 1.68^{ab}	$66.93 {\pm} 1.41^{ab}$	60.60 ± 0.97^{abc}	64.16±1.47 ^{bc}	58.26±1.61 ^c

Table 8. The effect of hot water heating process on the concentration (mg/kg) of different phenolic compounds

Values in same row with same superscript are not significantly (P<0.05) different.

As shown in the results sinapine and hydroxytyrosol are the major phenolic compounds present in the mustard powder at the levels of 2505.02 mg/kg and 653.89 mg/kg. respectively. During heating processes different variations in phenolic compounds has been observed. Hydroxytyrosol was decreased bv approximately 82%, 97% and 98% after heating for 20 minutes by hot water, steam and hot air, respectively. Hydroxytyrosol and tyrosol showed high heat sensitivity that might be related to the simple structure and low molecular weights of these phenolic compounds (Baldioli et al., 1996).

On the other hand Flavanomarein with high molecular weight (450.36 g·mol-1) and complex structure (Cerretani et al., 2009) showed a good heat stability and its concentration did not change significantly during different deheating processes. Sinapine (colin ester of sinapic acid) is one of the phenolic acid that its concentration in mustard powder sample was interestingly increased during processing by all of the heating media (Tables 6, 7 and 8). It might be emphasized that hot air and hot water showed the most and the least severe effect by approximately 37% and 15% increase in sinapine concentration, respectively.

There are also some newly formed phenolic compounds. Caffeic acid [(3,4-Dihydroxy-phenyl) -2-propenoic acid] was not existed in the original mustard powder but formed after heat treatments and its concentration significantly increased with time. Therefore after 12 min of hot water and air heating its level reached 1.74 mg/kg, 6.37 mg/kg and steam has similar effect in shorter time (8 min). Caffeic acid is an organic compound, classified as hydroxycinnamic acid and is one of the active compounds exhibiting biological activities. Numerous studies investigated the pharmacological properties of caffeic acid including inhibition on cancer cell proliferation (Chuu et al., 2012), powerful antioxidant (Chen et al.. 2001), immunomodulatory (Park et al., 2004), antimicrobial (Stojkovic et al., 2013), antiaging (Jeong, et al., 2011), and antiinflammatory (LeBlanc et al., 2012) and antidiabetic activity (Dhungyal et al., 2014). It might be concluded that heating by steam and hot air might positively have effect on mustard powder by forming a nutritional valuable phenolic compound. Ee et al. (2011) investigated the effect of roasting on the antioxidant activity of different types of seeds and reported that oxidative stability of the roasted seeds might be increased because of the release of phenolic compounds from the partially destroyed cell structure or progressing maillard reactions and formation of some type of compounds with antioxidant activity. Vujasinovi et al. (2012) confirmed that the chemical composition of seeds might be changed during roasting and might result in increasing the oxidative stability. Shrestha *et al.* (2013) reported that heat stability of mustard might be increased during thermal processes due to formation of new phenolic compound, however, the chemical mechanisms of this formation have not been yet clearly evaluated and remain ambiguous.

- Process Optimization by Response Surface Method (RSM)

Response surface methodology was applied to determine the optimum conditions for deheating powdered yellow mustard seed produce a product with the best to antimicrobial activity, color and total phenolic contents. According to the statistical analysis 5min of hot air heating in the autoclave might be introduced as the optimum process as shown in Table 9.

 Table 9. Functional properties of the optimal deheated mustard powder

Properties	Expected results
MIC against	
Bacillus subtilis	
Bacillus coagulans	- 1
Saccharomyces cerevisiae	1.6
Zygosaccharomyces rouxi	3.1
Color Characteristics	
a*	2.09
b*	34.50
L*	67.94
Residual enzyme activity	643.37
(µg glucose released / min)	
Total Phenolic contents	5376.42
(mg/kg)	

*without inhibitory effect (MIC>3.5%)

Conclusion

Hot/pungent flavor of mustard has usually limited its usage in the food industry. In this study, different combinations of the heating time and methods were applied to partially inactivate myrosinase and save antimicrobial activity simultaneously. One of the main factors that significantly

influenced the residual enzymatic activity was the moisture content of blanched product that seems to be a limiting parameter for hot water blanching of the relatively dried products such as plant seeds. According to the statistical analysis of the results, five minutes of hot air heating might be introduced as the optimum blanching process to produce a deheated yellow mustard powder with approximately 31% lesser enzyme activity (less hot flavor) and also suitable color characteristics and antimicrobial function and the least change in phenolic compounds. The effects of this multifunctional additive, produced under optimal conditions, on physicochemical and microbiological properties of foodstuffs such as different kinds of sauces need further studies.

Acknowledgement

The authors gratefully acknowledge and thank Pars Siavashan Trading Co. for providing yellow mustard seed samples and Mahmoudi (Color Science Dr. and Technology (ICST), Tehran, Iran) and Mr. Kazem Zadeh (Technoazma Company, Tehran, Iran) for their helpful assistance concerned with color analyses and accurate phenolic of compounds measurments respectively.

References

Ahn, E. S., Kim, Y. S. & Shin, D. H. (2001). Observation of bactericidal effect of allyl isothiocyanate on *Listeria monocytogenes*. *Food Science and Biotechnology*, 10, 31–35.

Amin, I. & Yee Lee, W. (2005). Effect of different blanching times on antioxidant properties in selected cruciferous vegetables. *Journal of Science of Food Agriculture*, 85, 2314–2320.

Appendini, P. & Hotchkiss, J. H. (2002). Review of antimicrobial food packaging. *Innovative Food Science and Emerging Technologies*, 3, 113–126.

Baldioli, M., Servili, M., Perretti, G. & Montedoro, G. F. (1996). Antioxidant activity of tocopherols and phenolic compounds of virgin olive oil. *Journal of the American Oil Chemists' Society*, 73, 11, 1589–1593.

Bastos, D. M., Monaro, R. & Séfora, M. (2012). Maillard reaction products in processed food: Pros and Cons. Nutrition Department, School of Public Health, So Paulo University, Brazil.

Cerretani, L., Bendini, A. & Rodriguez-Estrada, M. (2009). Microwave heating of different commercial categories of olive oil: Part I. Effect on chemical oxidative stability indices and phenolic compounds. *Food Chemistry*, 115, 1381–1388.

Chen, Y. J., Shiao, M. S. & Wang, S. Y. (2001). The antioxidant caffeic acid phenethyl ester induces apoptosis associated with selective scavenging of hydrogen peroxide in human leukemic HL-60 cells. *Anticancer Drugs*, 12, 143–149.

Chuu, C. P., Lin, H. P., Ciaccio, M. F., Kokontis, J. M., Hause, R. J., Hiipakka, R. A., Liao, S. & Jones, R. B. (2012). Caffeic acid phenethyl ester suppresses the proliferation of human prostate cancer cells through inhibition of p70S6K and Akt signaling networks. *Cancer Prevention Research (Phila).* 5, 788 – 797.

Dhungyal, P., Koirala, P. & Sharma, C. (2014). Caffeic Acid, Potent Phytocompound against Diabetes Mellitus. Skim Medicine University, *Medical Journal*, 1, 2, 152-161.

Eapen, K. E., Tape, N. W. & Sims, P. A. (1968). New process for the production of better-quality rapeseed oil and meal. I. Effect of heat treatment on enzyme destruction and color of rapeseed oil. *American Oil Chemists Society*, 45(3) 194-196.

Ee, K. Y., Agboola, S., Rehman, A. & Zhao, J. (2011). Characterisation of phenolic components present in raw and roasted wattle (Acacia victoriae Bentham) seeds. *Food Chemistry*, 129(3), 816-821.

Ghaderi Ghahfarokhi, M., Mamashloo, S., Sadeghi Mahoonak, A. R. & Alami, M. (2011). Evaluation of antioxidant activity, reducing power and free radical scavenging of different extract of *Artemisia annua L. Journal on Plant Science Researches*, 21, 1, 46-57 [In Persian].

Ghawi, S. K., Methven, L., Rastall, A. & Niranjan, K. (2012). Thermal and high hydrostatic pressure inactivation of myrosinase

from green cabbage: A kinetic study. Food Chemistry, 131, 1240–1247.

Han, J. H. (2000). Antimicrobial food packaging. *Food Technology*, 54 (3), 56–65.

Herzallah, S. & Holley, R. (2012). Determination of sinigrin, sinalbin, allyl- and benzyl isothiocyanates by RP-HPLC in mustard powder extracts. *Food Science and Technology*, 1-7.

Ildiko, S. G., Kla'ra, K. A., Marianna, T. M., A' gnes, B., Zsuzsanna, M. B. & Ba'lint, C. (2006). The effect of radio frequency heat treatment on nutritional and colloid-chemical properties of different white mustard (*Sinapis alba L.*) varieties. *Innovative Food Science and Emerging Technologies*, 7, 74 – 79.

International Olive Council. (2009). Determination of biophenols in olive oils by HPLC, COI/T.20/Doc No 29.

Jeong, C. H., Jeong, H. K., Choi, G. N., Kim, D. O., Lee, U. & Heo, H. J. (2011). Neuroprotective and anti-oxidant effects of caffeic acid isolated from Erigeron annuus leaf. *Chinese Medicine*, 6, 25-29.

Kreith, F. (2000). Handbook of Thermal Engineering. *CRC Press*, 3, 225-226.

Lara-Lledó, M., Olaimat, A. & Holley, R. A. (2012). Inhibition of *Listeria monocytogenes* on bologna sausages by an antimicrobial film containing mustard extract or sinigrin. *International Journal of Food Microbiology*, 2, 321-328.

LeBlanc, L. M., Pare, A., Jean-François, J., Hebert, M. J. & Touaibia, M. (2012). Synthesis and Antiradical/Antioxidant Activities of Caffeic Acid Phenethyl Ester and Its Related Propionic, Acetic, and Benzoic Acid. Analogues. *Molecules*, 17, 14637-14650.

Lin, C. M., Kim, J., Du, W. X. & Wei, C. I. (2000). Bactericidal activity of isothiocyanate against pathogens on fresh produce. *Journal of Food Protection*, 63, 25–30.

Luciano, F. B. & Holley, R. A. (2009) Enzymatic inhibition by allyl isothiocyanate and factors affecting its antimicrobial action against *Escherichia coli* O157:H7.*International Journal* of Food Microbiology, 131, 240–245.

Luciano, F. B., Belland, J. & Holley, R. A. (2011). Microbial and chemical origins of the bactericidal activity of thermally treated yellow mustard powder toward *Escherichia coli*

O157:H7 during dry sausage ripening. International Journal of Food Microbiology, 145, 69–76.

Nadarajah, D., Han, J. H. & Holley, R. A. (2005). Use of mustard flour to inactivate *Escherichia coli* O157:H7 in ground beef under nitrogen flushed packaging. *International Journal of Food Microbiology*, 99, 257–267.

Nilson, A. M. & Holley, R. A. (2012). Use of deodorized yellow mustard powder to control *Escherichia coli* O157:H7 in dry cured Westphalian ham. *Food Microbiology*, 30, 400-407.

Park, J. H., Lee, J. K., Kim, H. S., Chung, S. T., Eom, J. H., Kim, K. A., Chung, S. J. & Paik, S. Y. (2004). Immunomodulatory effect of caffeic acid phenethyl ester in Balb/c mice. *International Immunopharmacol*, 4, 429 – 436.

Saadoun, I., Hamed, K. M., Momani, M. & Ababneh, F. (2007). Effect of Three Orobanche spp. Extracts on Some Local Phytopathogens, Agrobacterium and Erwinia. *Turkish Journal of Biology*, 32, 113-117.

Shahidi, F., Wanasundara, U. N. & Amarowictis, R. (1994). Natural antioxidants from low-pungency. *Food Research International*, 27, 489-493.

Shahidi, F., Wanasundara, U. & Amarowicz, R. (1995). Isolation and partial characterization of oilseed phenolics and evaluation of their antioxidant activity. *Elsevier Science*, 1088 - 1099.

Shofran, B. G., Purington, S. T., Breidt, F. & Fleming, H. P. (1998). Antimicrobial properties of sinigrin and its hydrolysis products. *Journal of Food Science*, 63, 4, 620-624.

Shrestha, K., Gemechu, F. & De Meulenaer, B. (2013). A novel insight on the high oxidative

stability of roasted mustard seed oil in relation to phospholipid, Maillard type reaction products, tocopherol and canolol contents. *Food Research International*, 54, 587-594.

Smith, J. & Hong, S. (2003). Food additives data book. Oxford, Black Well Science.

Stojkovic, D., Petrovic, J., Sokovic, M., Glamoclija, J., Kukic-Markovic, J. & Petrovic S. (2013). In situ antioxidant and antimicrobial activities of naturally occurring caffeic acid, *p*coumaric acid and rutin, using food systems. *Journal of Science of Food and Agriculture*, 93, 3205 – 3208.

Turgis, M., Han, J., Caillet, S. & Lacroix, M. (2009). Antimicrobial activity of mustard essential oil against *Escherichia coli* O157:H7 and *Salmonella typhi*. *Food Control*, 20, 1073–1079.

USDA. (1992). Grading manual for tomato catsup. United States Department of Agriculture.

Vujasinovic, V., Djilas, S., Dimic, E. & Radocaj, O. (2012). The effect of roasting on the chemical composition and oxidative stability of pumpkin oil. *European Journal of Lipid Science Technology*, 114, 568–574.

Wang, X., Wiesenborn, D., Lindley, J. & Backer, L. (1994). Thermal inactivation of myrosinase in yellow mustard seed. *American Society of Agricultural Engineers*, 37, 879-886.

Yousefi, M., Mizani, M., Rasouli, S., Sharifan, A. & Bameni Moghadam, M. (2014). Optimization of deheating process of yellow mustard (Sinapis alba) by response surface methodology and desirability function. *International Journal of Biosciences*, 5, 99, 147-155.