

## **Survey on Physical, Chemical and Microbiological Characteristics of PAH-Contaminated Soils in Iran**

*\*M Arbabi, S Nasser, A R Mesdaghinia, S Rezaie, K Naddafi, G H Omrani and M Yunesian*

*Center for Environmental Research, Tehran University of Medical Sciences, Iran*

### **ABSTRACT**

Polycyclic aromatic hydrocarbons (PAHs) are one of the important groups of organic micro pollutants (Xenobiotics) due to their widespread distribution and low degradability in the environment (atmosphere, water and soil). Some PAHs exhibit carcinogenic and/or mutagenic properties and are listed by the United States Environmental Protection Agency (USEPA) and European Commission (EC) as priority pollutants. In this research three petroleum contaminated sites in Iran were selected in order to separate and classify PAH-degrading microorganisms. Samples were analysed for: soil physico-chemical properties, soil particle size distribution, Ultrasonic extraction of PAH (phenanthrene) and microbial analysis. Ultrasonic extraction method was shown to be a reliable procedure to extract a wide range of PAH concentrations from different soils, e.g. clay, silt, and clay-silt mixtures. Results showed that the extraction rate of phenanthrene in mentioned different soils was in the range of 85 – 100 percent. Results showed that two of three selected sites were contaminated with phenanthrene in the range of 10 – 100 mg/kg of soil, and had a reasonable population of PAH-degrading bacteria, which were enable to adaptate and degradate a concentration range of phenanthrene between 10 and 1000 mg/kg of soil. According to results, it can conclude that, the bioremediation of contaminated soils in Iran may be considered as a feasible practice.

**Keywords:** *Polycyclic aromatic hydrocarbons (PAHs), Bioremediation, PAH-degrading microorganisms, Ultrasonic extraction.*

### **INTRODUCTION**

Polycyclic aromatic hydrocarbons (PAHs) comprise a large and heterogeneous group of organic contaminants which are formed and emitted as a result of the incomplete combustion of organic material (Mulder et al., 2001). Anthropogenic contamination sources, such as road traffic and combustion of fossil fuels predominate, but there are also natural sources, e.g. volcanic eruptions and forest fires

(Lee et al., 2001). Polycyclic aromatic hydrocarbons (PAHs) are one of the important groups of organic micropollutants (xenobiotics) due to their widespread distribution in the environment (atmosphere, water and soil). They are produced by incomplete combustion of fossil fuels as well as by diagenetic processes during fossil fuel formation (Khodadoust et al., 2000). In smaller amounts, PAHs are naturally produced by forest fires and possibly microbiological synthesis (Turlough, 1999). PAHs consist of fused benzene rings in linear, angular or clustered arrangements, and contain by definition only carbon and hydrogen atoms. However, nitrogen, sulfur and oxygen atoms

**\*Corresponding Author:** Tel & Fax: +98 21 66112762, P.O. Box: 14185-766, E-mail: [marbabi@razi.tums.ac.ir](mailto:marbabi@razi.tums.ac.ir)

may readily substitute in the benzene rings to form heterocyclic aromatic compound which are commonly grouped with the PAHs (Feng, and Aldrich, 2000; Lundstedt, 2003; Slavica et al., 2003). Some PAHs exhibit carcinogenic and/or mutagenic properties and are listed by the United States Environmental Protection Agency (USEPA) and European Commission (EC) as priority pollutants (Amellal et al., 2001). Many PAHs have toxic, mutagenic and/or carcinogenic properties.


Several microorganisms have been isolated that are capable of degrading PAHs. However, PAHs are relatively persistent and recalcitrant in soils and are more difficult to be degraded than many other organic contaminants under natural conditions (Wilson and Lones, 1993).

Since PAHs are hydrophobic compounds with low solubility in water, they have a greater tendency to bind with organic matter or soil, limiting their availability to microorganisms (Lee et al., 2001; Yun et al., 2003).

Consequently, our environment has been become highly polluted with different PAHs. Phenanthrene, a three-ring angular PAH, is known to be a human skin photosensitizer and mild allergen and is mutagenic to bacterial system under specialized conditions (Smanta et al., 2002; Yun et al., 2003).

Many PAHs contain a "bay-region" and a "K-region". The bay-and-K-region epoxides, which can be formed metabolically, are highly reactive both chemically and biologically. These epoxides are suspected to be ultimate carcinogens (Turlongh, 1999; Goldman et al., 2001). As phenanthrene contains bay- and K-region (Fig 1), it is also used as a model substrate for studies on the metabolism of bay-region and K-region-containing carcinogenic PAHs such as benzo [a] pyrene, benzo [a] anthracene and chrysene (Cerniglia, 1989; Mastrangela et al., 1997; Falahtpisheh et al., 2001). There are several reports of the degradation of phenanthrene by different bacteria (Bucker et al., 1979; Weis et al., 1998; Stegeman et al., 2001).

**Fig.1:** Formation of CO<sub>2</sub> from the degradation of



phenanthrene(Cerniglia, 1989)

There are several treatment methods for PAHs contaminated soils including: incineration, fixation, solvent extraction wet oxidation, land-fill and bioremediation. Other than incineration, bioremediation is the only practical consideration for complete degradation of organic contaminants (Norris et al., 1994).


Other methods except bioremediation needed to energy, transportation and monitoring.

There is an increasing interest in the cleaning of soil contaminated with organic compounds using bioremediation (Wilson and Lones, 1993). Bioremediation has become an intensive area for research and, as result; rapid progress has been made in developing effective microbial bioremediation processes (Fig 2) (Smanta et al., 1999). The objectives of this study were as follows:

Separation of PAH degrading micro organisms from contaminated soils in petroleum contaminated sites in Iran;

Extraction of PAH using ultrasonication method and;

Analysis of extracted samples by HPLC-UVD using a PAH special column.



**Fig.2:** Fate, toxicity and remedition of polycyclic aromatic hydrocarbons (PAHs) from the environment.

## MATERIALS AND METHODS

### Initial site investigation

In order to separate PAH-degrading microorganisms, the soil samples were collected from three different petroleum contaminated sites:

#### (1) Tehran Oil Refinery site

The oil Refinery of Tehran is located at south part of Tehran, which produces a variety of oil products from crude oil since 19 years.

According to field studies, the site is polluted by oily water effluents and emission by oil combustion stacks. This site was selected as oil refinery site.

#### (2) Bahregan Oil Zones

Bahregan oil zone is one of the most important rude oil extraction zones of Iran from Persian Gulf.

This site was selected as an oil extraction and transportation from sea to land; the soil samples were collected from oil sludges that are in

Bahregan oil zone shore. This site is located between Genaveh and Daylam port, near the Imam Hassan village.

#### (3) Oil Products Reservoir site

This site is located at northwest of Tehran, where oil reservoirs are provided in order to store the oil products for emergency conditions. This site was selected as oil reservoir site.

### Soil sampling

Baseline soil samples for contaminant concentrations, soil chemistry and microbiology were collected from aged contaminated soils around the above mentioned three sites. The samples were collected in the range of 3-4 kg from the surface and 10 cm deep layer of soil. Prior conducting any analysis, the coarse pieces and stones were separated and the rest were mixed well.

The sub samples were kept cold (3-5) °C for microorganisms' extraction. Microbial analyses were conducted within 24-48 h after sampling.

### Remediation strategy

#### Soil physico-chemical analyses

Fandamental soil physico-chemical properties were analysed through following standard methods:

- Soli size particle distribution determination based on Unified Soil Classification (ASTM-D-2487);
- Soil moisture content (APHA);
- Organic carbon (ISO, DIS 10694 @11277);
- Polycyclic aromatic hydrocarbon, phenanthrene, ( USEPA methods #3550B, #8310 and NIOSH #5506 by PHLC with UV detection (EPA, 2003; NIOSH, 2003).

#### PAH analysis

Phenanthrene concentration was measured by HPLC, with chromatographic conditions as follows:

- Analytical column: C<sub>18</sub> Ultra Sep ES PAH QC Speica, 60 × 2 mm ID
- Folwrate (ml/min): 0.5
- Injection rate: 50 µl
- Elute: Acetionitrl/water: 40-100%
- UV detector wavelength: 254 <sup>nm</sup>

According to pre-test results the optimum elute condition for phenanthrene was determined at 60/40 (acetonitril/water) (Herbert, 2003).

#### *PAH extraction methods*

There are four important methods for solid phase extraction (SPE) of PAHs from contaminated soil as below:

- Soxhlet Extraction
- Sonication
- Supercritical Fluid Extraction using CO
- Microwave Assisted Extraction

In this study, extraction was conducted by sonication method using ultrasonic instrument (RK31H, Bandelin Electronic, Sonorex, 35 kHz, Germany), because of fast and easy extraction process compared with other methods.

Certified phenanthrene contaminated soil samples (1g) were dried at air temperature and after sieving (through 50 meshes sieve), were suspended in 10 ml of acetonitril and extracted in ultrasonic bath at 40-45°C for 45- 60 minutes. Extracts were settled for 10 min, and centrifuged at 5000-6000 rpm for 15 min. Elutes were filtered by 25mm, 2µm pore size, PTFE membrane filter and submitted to HPLC-UV analysis.

#### *Soil microbial analyses*

Microorganisms were extracted from the contaminated soils by mixing of 1g soil with 10 ml of sterile Na<sub>2</sub>P<sub>2</sub>O<sub>7</sub> solution (2.8g/ liter) in 50 ml erlen mayer for 2h on a shaker (250 rpm ) (Kastner et al., 1998). The soil particles were allowed to sediment for 30 min. the supernatant was diluted and plated on solid media.

Initial microbial analysis was conducted both for fungi and bacteria. Fungi determination test was carried out on solid media containing cloroamphenicole and cyclohexamide.

Total CFU were determined using plate count agar or HPC (heterotrophic plate count) media. Also MPN analyses were conducted using lactose broth media with 15 tubes method (APHA, 1998).

Microorganisms were extracted from the contaminated soils, and supernatant after dilution by 10<sup>-3</sup> - 10<sup>-5</sup> times was plated on solid

media. After separation of bacteria, the stock samples of separated microbes were prepared by adding of 500 µl of polluted sample and 500µl of glycerin together. The stock samples were holed at - 80°C for future pilot study.

## RESULTS

### *Soil physical analyses*

Regarding to next studies on effects of soil structure in PAH degradation in contaminated soil, the soil particle size analysis was carried out on different soil samples, which collected from three before mentioned sites.

The soil particle size determination analysis was conducted using the stack of sieves from 4 to 200 meshes opening size. The results for determination of particle size of contaminated soils of oil refinery of Tehran, Bahregan oil zone, and oil products reservoir sites are presented in Tables 1, 2 and 3 respectively.

After determination particle size distribution of soils, to determine the type of soil the percentage of gravel, sand, silt and clay were measured.

Properties of contaminated soils are presented in Table 4.

**Table 1:** Determination of particle size of contaminated soil from oil refinery of Tehran

Sieve No. (mesh)	Openin g (mm.)	Mass retained (grams) Mr	Retained% (Mr/M)×10 0	Finer (%)
4	4.75	0	0	100
10	2.00	0.9	0.225	99.8
35	0.50	98.93	24.73	75.07
50	0.30	48.39	12.1	62.97
60	0.25	25.26	6.31	56.66
70	0.21	25.1	8.78	47.88
80	0.18	1.76	0.44	47.44
100	0.15	18.76	4.69	42.75
120	0.12	39.19	9.8	32.96
200	0.075	91.7	22.92	10.04
Pan	--	40	10	----
Total weight	--	400 g	100	----

**Table 2:** Determination of particle size of contaminated soil from Bahregan oil zone

Sieve No. (mesh)	Opening (mm)	Mass retained (grams) Mr	Retained % (Mr/M)×100	Finer (%)
4	4.75	3.73	1.24	98.76
10	2.00	9.28	3.09	95.7
35	0.50	21.90	7.30	88.4
50	0.30	5.65	1.88	86.52
60	0.25	5.24	1.75	84.8
70	0.21	80.07	26.69	58.11
80	0.18	11.61	3.87	54.24
100	0.15	35.16	11.72	42.52
120	0.12	102.6	34.2	8.32
200	0.075	4.11	1.37	6.9
Pan		20	6.90	--
Total weight	----	300 g	100	----

**Table 3:** Determination of particle size of contaminated soil from oil reservoir site (Northwest part of Tehran)

Sieve No. (mesh)	Opening (mm)	Mass retained (grams) Mr	Retained % (Mr/M)×100	Finer (%)
4	4.75	13.49	4.5	95.5
10	2.00	67.23	22.41	73.09
35	0.50	78.65	26.2	46.89
50	0.30	73.51	24.5	22.39
60	0.25	10.85	3.6	18.79
70	0.21	22.35	7.45	11.35
80	0.18	0	0	11.35
100	0.15	4.47	1.48	8.86
120	0.12	11.95	3.98	4.86
200	0.075	4.5	1.5	4.38
Pan		13	4.38	-
Total weight	-	300 g	100	-

**Table 4:** Physico – chemical properties of contaminated soils

Sample Locations			
Item	Oil refinery of Tehran	Behragan Oil Zone	Oil Reservoir Site (Tehran)
Gravel (%)	0	1.24	4.5
Sand (%)	90	91.86	91.12
Silt (%)	--	6.9	--
Clay (%)	10	--	4.38
Moisture (%)	6.4	1	4
pH	6.8	7.1	7
Organic carbon (%)	7.89	7.312	6.72
Bulk density (g/ml)	0.7	0.6	0.75
Uniformity coefficient (Uc)	3.70	1.79	7.8
Coefficient of curvature (Cc)	0.49	0.73	0.61
Effective size(mm)	0.075	0.12	0.16
Type of soil *	SP-SC	SP-SM	SP-SC

**\* Note:**

- SP-SC: Clayey – sands, poorly graded sand – clay mixture.
- SP-SM: Silty – sands, poorly graded sand – silt mixture.
- Soil classification has been done based on Unified Soil Classification for coarse-grained soils.

**Soil microbiology**

Initial analysis of soils for fungi determination; do not show any growth on solid media. But CFU and MPN analysis results for different three types of contaminated soils are presented in Table 5.

**Table 5:** Initial Soil Microbiology

Type of Soil	Initial soil PAH Conc. (mg/kg)	Ultrasonic extraction Conc. (mg/kg)	Recovery (%)	No. of samples
CLAY	100	90-96	90-96	20
	500	425-470	85-94	20
	1000	850-950	85-95	20
SILT	100	90-100	90-100	20
	500	450-500	90-100	20
	1000	850-970	85-97	20
CLAY & SILT	100	85-95	85-95	20
	500	425-475	85-95	20
	1000	850-950	85-95	20
SILT	2000	1800-2000	90-100	5
	5000	4500-4750	90 – 95	5
	10000	9000-9500	90 – 95	5
CLAY	2000	1800-1980	90-99	5
	5000	4350-4750	87 – 95	5
	10000	9000-9500	90 – 95	5

### PAH analysis

In this study phenanthrene is selected as 3-reings PAHs prior the contaminated soils analysis for phenanthrene, it was necessary the rate of extraction of PAH using ultrasonication method to be measured. In this regards, ultrasonic extraction procedure was carried out with different soil types, e.g. clay, silt and clay and silt mixture and different initial soil phenanthrene concentrations. The results for ultrasonic extraction rate in different soils with different phenanthrene concentrations are presented in Table 6. Due to sealing of oil reservoirs and leakage control in oil reservoir site of Tehran, the phenanthrene soil contamination was not detected. However, in two other sites the phenanthrene soil contamination was measured because of high pollution of these sites. The results for phenanthrene measurement in contaminated collected soils are shown in Table 7.

**Table 6:** Ultrasonic extraction of PAH (phenanthrene) from contaminated soils

Sample location	Microbial populations	
	MPN/g soil	CFU/g soil (heterutrophic populations)
Oil Refinery of Tehran	$5 \times 10^6$	$6.5 \times 10^5$
Bagregan Oil Zone	$4.5 \times 10^6$	$6.7 \times 10^5$
Oil Reservoir Site (Tehran)	$8.5 \times 10^7$	$7 \times 10^6$

**Table 7:** Ultrasonic extraction of PAH (phenanthrene) from collected soil samples

Sample Locations	Phenanthrene Conc.(mg/kg soil)
Oil refinery of Tehran	60 (approx.)
Behragan Island	12 (approx.)
Oil reservoir site (Tehran)	Zero

## DISCUSSION

According to sieve analysis and uniformity coefficient ( $U_c$ ) and Coefficient of curvature ( $C_c$ ) based on unified soil classification (Table 4), the type of soils were SP-SC (poorly graded, sand-clay mixture) and SP-SM (poorly graded, sand-silt mixture).

Results of physico-chemical, PAH concentration, and microorganism's analysis in three types of petroleum contaminated soils collected from different sites in Iran, showed that:

Two of three selected sites were contaminated with phenanthrene in the range of 10-100 mg/kg of soil. According to soil particle size distribution analysis, the contaminated soils were classified as SP-SC (poorly graded clay-sand mixture) and SP-SM (poorly graded sand-silt mixture). Contaminated soil microbiology,



(CFU analysis), showed that the contaminated soils contains the reasonable populations of bacteria ( $6.5 \times 10^5$  -  $7 \times 10^6$  CFU). Hence, it can be concluded that the bioremediation of contaminated soils in Iran may be considered as a feasible practice.

Inoculation tests results showed that, the separated bacteria demonstrated the same tolerance both with high and low level phenanthrene contaminated soils.

Ultrasonic extraction method was a reliable procedure for extracting of different PAH concentrations from different soils, e.g. clay, silt, and clay-silt mixture. It should be mentioned that, the Ultrasonic extraction procedure needs much more clean up work after injection of elutes to HPLC.

## ACKNOWLEDGEMENTS

Authors present their gratitude to the Center for Environmental Research (CER) of Tehran University of Medical Sciences (TUMS), for its financial support for this research.

## REFERENCES

- Amellal N, Portal J M, Berthelin J (2001). Effect of soil structure on bioavailability of polycyclic aromatic hydrocarbons within aggregates of a contaminated soil. *Applied Geochemistry*, 16:1611-19.
- APHA, AWWA, WEF (1998). Standard methods for the examination of water and wastewater, 20<sup>th</sup> ed. Washington.
- Bucker M (1979). Mutagenicity of phenanthrene and phenanthrene K-region derivatives. *Mutat Res*, 66: 337-48.
- Carmicheal L M, Christman R F, Pfaender F K (1997). Desorption and mineralization kinetics of phenanthrene and chrysene in contaminated soils. *Environ Sci Technol*, 31(1): 126-32.
- EPA (1996). Poltcyclic aromatic hydrocarbons (PAHs): Ultrasonic Extraction, Method 3550B and 8310.
- EPA (2003). Extraction methods of polycyclic aromatic hydrocarbons. <http://www.epa.gov/paoswer/hazwaste/test/sw846.htm>.
- Falahatphisheh M H (2001). Antagonistic interactions among nephrotoxic polycyclic aromatic hydrocarbons. *Journal Toxicol Environ Health*, 62: 543-60.
- Feng D, Aldrich, C (2000). Sonochemical treatment of simulated soil contaminated with diesel. *Advances in Environmental Research*, 4: 103-12.
- Goldman R (2001). Smoking increase carcinogenic polycyclic aromatic hydrocarbons in human lung tissue. *Cancer Res*, 61: 6367 - 71.
- Harayama S (1997). Polycyclic aromatic hydrocarbon bioremediation design. *Current Opinion in Biotechnology*, 8: 268-73.
- Herbert H (2003). Sample preparation and HPLC analysis of PAHs extracted soil samples. *Certified Quality Manegement*, DIN EN ISO 9001, Unexas Application, KNAUER.
- Kastner M, Breuer M J, and Mahro B (1998). Impact of Inoculation protocols, salinity, and pH on the degradation of polycyclic aromatic hydrocarbons (PAHs) and survival of PAH-degrading bacteria introduced into soil. *Appl Environ Microbiol*, 64(1): 359-362.
- Khodadoust A P, Bagchi R, Suidan M T, Brenner R C, and Sellers N G (2000). Removal of PAHs from highly contaminated soils found at prior manufactured gas operations. *Journal of Hazardous Materials*, B80:159-74.
- Lee P K, Ong S K, Golchin J, Nelson G L (2001). Use of solvents to enhance PAH biodegradation of coal tar-contaminated soils. *Wat Res*, 35 (16):3941-49.
- Lundstedt S (2003). Analysis of PAHs and their transformation products in contaminated soil and remedial processes. University of Amsterdam, the Netherlands. [www.google.com](http://www.google.com).

- Mastrangela G (1997). Polycyclic aromatic hydrocarbons and cancer in man. *Environ Health Perspect*, 104: 1166-70.
- Mulder H, Breure AM, Rulkens WH (2001). Prediction of complete bioremediation periods for PAH soil pollutants in different physical states by mechanistic models. *Chemosphere*, 43: 1085-1094.
- NIOSH (1998). Manual of analytical methods (NMAM): Polynuclear aromatic Hydrocarbons by HPLC, Method 5506, Issue 3. 4<sup>th</sup> ed. pp: 1-9.
- Norris R D, Hinchey R E, Brown R, McCarty PL, Semprini L, Wilson J T, Kampbell DH, Reinhard M. and Ward C H (1994). *Handbook of bioremediation*. Lewis Publishers, Boca Raton.
- Samanta S K, Chakraborti A K, Jain R K (1999). Degradation of Phenanthrene by different bacteria: evidence for novel transformation sequences involving the formation of 1-naphthol. *Appl Microbiol Biotechnol*, 53: 98-107.
- Samanta S K, Singh O, Jain R K (2002). Polycyclic aromatic hydrocarbons: environmental pollution and bioremediation. *TRENDS in Biotechnology*, 20 (6): 243-248.
- Slavica S D, Slavica B, Brantner B (2003). A comparison of ultrasonic extraction and soxhlet extraction of polycyclic aromatic hydrocarbons from soil. *Umweltanalytisches labor, Sachenplatz 13, A-1200 Vienna, Austria*.
- Stegeman J J (2001). Cytochrome P 4501 A expression in mid water fishes: potential effects of chemical contaminants in remote oceanic zones. *Environ Sci Technol*, 35: 54-62.
- Turlough F G (1999). The extraction of aged polycyclic aromatic hydrocarbon (PAH) residues from a clay soil using sonication and soxhlet procedure: a comparative study. *Environ Monit*, 1: 63-67.
- Weis L M (1998). Bay and bay like regions of polycyclic aromatic hydrocarbons were potent inhibitors of gas junctional intercellular communication. *Environ Health Perspect*, 106: 17-22.
- Wilson S C, Jones K C (1993). Bioremediation of soil contaminated with polynuclear aromatic hydrocarbons (PAHs): a review. *Environ Pollut*, 81(3): 229-249.
- Yun T, Tianling Z, Xinhong W (2003). PAHs contamination and PAH-degrading bacteria in Xiamen western sea. Chemical speciation and Bioavailability. *Chemical Speciation and Bioavailability*, 14: 25-33.