BIOCHEMICAL STUDY OF FRESHWATER FISH CATLA CATLA WITH REFERENCE TO MERCURY CHLORIDE

¹P. Martin Deva Prasath, ^{*2}S. Arivoli

¹Department of Chemistry, TBML College, Porayar-609 307, Nagai District, Tamil Nadu, India

²Department of Chemistry, Mannai Rajagopalasamy Government Arts College, Mannargudi–614 001, Tamil Nadu, India

Received 16 March 2007; revised 18 September 2007; accepted 15 February 2007

ABSTRACT

Toxic activity of mercury chloride was tested in vivo on fresh water fish *Catla catla*. Acute toxicity tests were conducted to measure the impact of toxicity on the fishes within a short period at the various concentrations of HgCl₂ (0.1, 0.5, 1, 1.5, 2, 2.5 and 3mg/L). The protein and carbohydrate were estimated using Anthrone by standard methods and enzymes such as Na⁺-K⁺, Mg²⁺ and Ca²⁺ adenoxide triphosphatases were determined caloriemetrically. Depletion of protein was observed at all exposure periods. The biochemical estimation values of carbohydrates in muscle, intestine and brain showed significant values with P<0.05. The activity of Mg²⁺ adenoxide triphosphatases in muscle, intestine and brain was studied and the maximum depletion was observed at 96h, 72h and 72h, respectively. Regarding activity of Ca²⁺ adenoxide triphosphatases in muscle, the maximum depletion of this enzyme activity in muscle was observed at all exposure periods, except 96h. In intestine, enzyme activity was gradually decreased in all exposure and an increase was found during 96h. In brain, the enzyme activity was observed at all exposure periods except 96h.

Key words: Toxic, protein, carbohydrate, enzyme, Catla catla

INTRODUCTION

Heavy metals are being passed on into aqueous environments through industrial processes, sewage disposal, soil leaching and rainfall. The concentrations of these heavy metals are sublethal or lethal to aquatic organisms when the duration of exposure to these metals are prolonged (Eisler and Gardener, 1973). It is well documented that effect of heavy metals are dependent upon the physical and chemical conditions of the environment especially water salinity, hardness, pH and dissolved oxygen and can act synergistically. Heavy metals from several industrial mining and other sources enormously contribute to the pollution problem in rivers and streams resulting in adverse impacts on biota including fish.

Fish population is generally considered very sensitive to all kinds of environmental changes to which it is exposed as they are exclusively aquatic with external mode of fertilization. Certain stages

*Corresponding author-Email: *arivu3636@yahoo.com* Telefax: +91 04367 255440 in the life cycle of fresh water fish are more susceptible to environmental and pollution stresses (Von Westernhagan, 1988). Even though many studies are available in toxicity of various fishes (Jagadeesan *et al.*, 1991; Margarat *et al.*, 1991; Jagadeesan *et al.*, 2001; Francis *et al.*, 2002;), there is no report available on *Catla catla*. Hence the present study has been carried out on *Catla catla* with reference to mercury chloride.

MATERIALS AND METHODS

Finger lings of *Catla catla* species were collected and kept in aquarium and the water used was clear and unchlorinated (APHA, 1980). Fishes were fed daily with fish pellets and acclimatized for 30 days. Acute toxicity tests were conducted to measure the impact of toxicity on the fishes within a short period (Sprague, 1969) at the various concentration of HgCl₂ (0.1, 0.5, 1, 1.5, 2, 2.5 and 3mg/L). The tissue was homogenized with 1mL of Tris-HCl buffer. The protein and carbohydrate P. Martin, et al., BIOCHEMICAL STUDY OF FRESHWATER ...

were estimated by standard methods (Carroll *et al.*, 1956) and enzymes such as Na⁺-K⁺, Mg²⁺ and Ca²⁺ adenoxide triphosphatuses were analyzed by caloriemetric method (Fiske *et al.*, 1925).

RESULTS

Protein

Depletion in the protein content in the muscle, intestine and brain of the (*Catla catla*) exposed

to mercury chloride for 24h, 48h, 72h and 96h in 0.1, 0.3 and 0.5m/L sub-lethal concentrations were estimated. Protein content in the muscle of the control, depletion and increase of muscle protein are shown in Fig. 1. Intestine of the mercury chloride treated fish shows a gradually decrease in protein level. Depletion of protein was observed at all exposure periods (Fig. 2). The level of protein in the control fish brain, is presented in Fig. 3.



Fig. 1: Changes in total protein level (mg/g) in muscle tissue of *Catla catla* exposed to different sublethal concentrations of mercury chloride with different exposure period (n=6)



Fig. 2: Changes in total protein level (mg/g) in intestine tissue of *Catla catla* exposed to different sublethal concentrations of mercury chloride with different exposure periods (n=6)



Fig. 3: Changes in total protein level (mg/g) in brain tissue of *Catla catla* exposed to different sublethal concentrations of mercury chloride with different exposure periods (n=6)

Carbohydrates

Depletion of carbohydrate content of the muscle (Fig. 4), intestine (Fig. 5) and brain (Fig. 6) of *catla catla* exposed to the mercury chloride for 24, 48, 72 and 96h in 0.1, 0.3 and 0.5mg/L sublethal concentrations were estimated. Among these, the maximum depletion of carbohydrate was

observed in intestine during 96h. Generally, depletion in carbohydrate content is directly proportional to the exposure period of the toxicant. The obtained biochemical estimation values of the muscle, intestine and brain were subjected to statistical analysis and showed significant values at P<0.05.



Fig. 4: Changes in total carbohydrate level (mg/g) in muscle tissue of *Catla catla* exposed to different sublethal concentrations of mercury chloride with different exposure periods (n=6)



Fig. 5: Changes in total carbohydrate level (mg/g) in intestine tissue of *Catla catla* exposed to different sublethal concentrations of mercury chloride with different exposure periods (n=6)



Fig. 6: Changes in total carbohydrate level (mg/g) in brain tissue of *Catla catla* exposed to different sublethal concentrations of mercury chloride with different exposure periods (n=6)

Enzyme analysis

The activity of Na⁺-K⁺, Ca²⁺ and Mg²⁺ ATPase in muscle (Fig. 7), intestine (Fig. 8) and brain (Fig. 9) were estimated in the experimental fish after 24, 48, 72 and 96h in 0.1, 0.3 and 0.5mg/L sub-lethal concentration of exposure to the mercury chloride. The alteration in the activity of enzymes of Na⁺ K⁺ ATPase were observed. The activity of Mg²⁺ ATPase in muscle (Fig. 10), intestine (Fig. 11) and

brain (Fig. 12) were studied and the maximum depletion was observed in 96h, 72h, and 72h, respectively. The activity of Ca^{2+} ATPase in muscle and depletion of enzyme activity was observed at all exposure periods, except for 96h (Fig. 13). In intestine, enzyme activity was gradually decreased in all exposures and increased during 96h (Fig. 14). In brain, the enzyme activity was also observed at all exposure periods except 96h (Fig.15).



Fig. 7: Alteration in Na⁺ K⁺ ATPase level (mm pi released/mg protein/h) in muscle tissue of *Catla catla* exposed to different sublethal concentrations of mercury chloride with different exposure periods (n=6)



Fig. 8: Alteration in Na⁺ K⁺ ATPase level (mm pi released/mg protein/h) in intestine tissue of *Catla catla* exposed to different sublethal concentrations of mercury chloride with different exposure periods (n=6)



Fig. 9: Alteration in Na⁺ K⁺ ATPase level (mm pi released/mg protein/h) in brain tissue of *Catla catla* exposed to different sublethal concentrations of mercury chloride with different exposure periods (n=6)

Iran. J. Environ. Health. Sci. Eng., 2008, Vol. 5, No. 2, pp. 109-116



Fig. 10: Alteration in Mg²⁺ ATPase level (mm pi released/mg protein/h) in muscle tissue of *Catla catla* exposed to different sublethal concentrations of mercury chloride with different exposure periods (n=6)



Fig. 11: Alteration in Mg²⁺ ATPase level (mm pi released/mg protein/h) in intestine tissue of *Catla catla* exposed to different sublethal concentrations of mercury chloride with different exposure periods (n=6)



Fig. 12: Alteration in Mg²⁺ ATPase level (mm pi released/mg protein/h) in brain tissue of *Catla catla* exposed to different sublethal concentrations of mercury chloride with different exposure periods (n=6)



Fig. 13: Alteration in Ca²⁺ ATPase level (mm pi released/mg protein/h) in muscle tissue of *Catla catla* exposed to different sublethal concentrations of mercury chloride with different exposure periods (n=6)

P. Martin, et al., BIOCHEMICAL STUDY OF FRESHWATER



Fig. 14: Alteration in Ca²⁺ ATPase level (mm pi released/mg protein/h) in intestine tissue of *Catla catla* exposed to different sublethal concentrations of mercury chloride with different exposure periods (n=6)



Fig. 15: Alteration in Ca²⁺ ATPase level (mm pi released/mg protein/h) in brain tissue of *Catla catla* exposed to different sublethal concentrations of mercury chloride with different exposure periods (n=6)

DISCUSSION

Biochemical Analysis

Protein

Proteins are involved in major physiological events therefore the assessment of the protein content can be considered as a diagnostic tool to determine the physiological phases of organism. Proteins are highly sensitive to heavy metal poisoning (Jacobs *et al.*, 1977). Depletion of protein content has been observed in the muscle, intestine and brain of the fish *Catla catla* as a result of mercury chloride toxicity. When an animal is under toxic stress, diversification of energy occurs to accomplish the impending energy demands and hence the protein level is depleted (Neff, 1985).

The depletion of total protein content may be due to breakdown of protein into free amino acid under the effect of mercury chloride at the lower exposure period (Shakoori *et al.*,1994).

Corresponding to it a level of increase in protein content was observed at increased exposure period of 96h at 0.1, 0.3 and 0.5mg/L sublethal

concentration. These indicate that mercury induces proteolysis in the fish even under sublethal toxic stress resulting in elevated levels of protein content; but, the degree of proteolysis appears time-dependent, as the decrease in protein levels progressed significantly at 24h, 48h and 72h but regressed and attained almost normally at 96h of exposure.

Metals could alter the structure, permeability and integrity of lysosomal membranes resulting in the diffusion of their enzyme into cytosol (Sternlib *et al.*, 1976). Hence high activity of protease, a lysosomal enzyme, in the organs of fish might be due to the damage caused by mercury to lysosomes. Elevated protease activity induced proteolysis, the intensity increased with the increase in exposure period from 25h to 72h may be the increase in free amino acid pool due to increased proteolysis would act as an osmotic and ionic effectors to bring the electrostatic equilibrium between the external medium and blood (Schmit-Nielson, 1975; Jurss, 1980). Besides, free amino acids would also serve as precursors for energy production under stress, and for the synthesis of required proteins to face the metal challenge (Sree devi *et al.*, 1991).

Carbohydrate

In the present study, an initial decrease and an increase in the level of total carbohydrate has been noticed in the muscle and intestine tissues. The disturbance in the carbohydrate metabolism was considered as one of the most outstanding biological lesions due to the action of heavy metal (De Bruin, 1976). The decrease in carbohydrate content in the muscle, intestine and brain may be due to glucose utilization to meet excess energy demand imposed by severe anaerobic stress of mercury intoxication (Margarat et al., 1999). Another possible reason for depletion in the tissue may be due to impairment of glycogen synthesis. Under hypoxic conditions; fish derive the energy by anaerobic breakdown of glucose which is available to the cells with the increased glycogenolysis. The observed depletion of carbohydrate in the present study explains the increased demand of these molecules to provide energy for the cellular biochemical process under toxic manifestations. Similar results were observed in Thalmile crenata, Anabas testudienues and Anabas scandens, when exposed to copper, lead, nitrate and mercury chloride, respectively (Villalan et al., 1988; Mary Candravathy et al., 1991). During the initial period, metabolic activity failed to recover indicating the effects of accumulated mercury in the tissues. However, at 96h the metabolites reached to normal levels. It indicates the slow elimination of mercury and resynthesis of metabolities against the toxicants, indicating the reduced rates of glycogenolysis and glycolysis. The recovery could be attributed to restoration of regulatory function of phosporylases by elimination of toxicant (Holcombe et al., 1976; Richert et al., 1979) from the endocrine glands like pancreas and adrenals.

Enzyme analysis

Adenosine triphosphatases (ATPases)

The major target molecules affected by metals include ion dependent ATPases, which lead to disturbances in ion homeostasis. The inhibition of ATPases lead to decreased ATP breakdown and reduced the availability of free energy. The reduced energy supply may affect several metabolic processes (Ramalingam *et al.*, 1999).

$Na^{+}K^{+}$ -ATPase

Na⁺K⁺-ATPase is considered as a marker enzyme to understand the physiological impairment of the cell (Campbell *et al.*, 1974). The inhibition of membrane bound ATPases in the tissues, muscle, intestine and brain could be result of physicochemical alteration of the membrane. The alteration in ionic balance depolarises the nerve and due to depolarisation the nerve cells increase in releasing the neuro transmitter which inturn inhibits Na⁺K⁺-ATPase activity (Kimellberg *et al.*, 1974). Inhibition of the enzyme increases intracellular Ca²⁺ and decreases intracellular Mg²⁺ concentration (Karamer *et al.*, 1991).

Ca^{2+} ATPases

Calcium ions are essential for the transmitter release and the level of calcium is regulation by Ca^{2+} ATPase (Rubin, 1970). Ca^{2+} ATPase maintains low intracellular Ca^{2+} ions than the Ca^{2+} concentration of extra cellular medium (Harrison *et al.*, 1980). Decreased Ca^{2+} ATPase due to the activity of Mercury chloride may lead to high internal Ca^{2+} level (Duncan, 1967). Increase in intracellular accumulation of Ca^{2+} ions there by increases the release of neuro-transmitter from the synaptic vesicles by exocytosis (Yamaguchi *et al.*, 1979) which may inhibit Na⁺K⁺ ATPase.

Mg^{2+} ATPases

Mg²⁺ ATPase is involved in the control of passive permeability (Duncan, 1967). It is also involved in oxidative phosphorylation, the inhibition due to toxicants directly prevents or reduces the oxidative phosphorylation (Nuefeld et al., 1979). Inhibition of Mg²⁺ ATPase may result in effects on energy metabolism and respiration (Desaiah et al., 1977). During the initial exposure period, metabolities activity failed to recover indicating effects of accumulated mercury in the tissues. However on the 96 hr of exposure, the metabolities reached to near normal levels. Generally, more energy is needed to mitigate any stress condition and this may be obtained from carbohydrates and proteins. Increase in structural proteins could help the animal to fortify its organs for developing resistance and increase in soluble fraction from the general intracellular environment and help the animal to adapt to the imposed toxic stress Sivaramakrishnan *et al.*, 1998).

ACKNOWLEDGEMENT

The authors thank the Heads of Department of Zoology and chemistry for their encouragement.

REFERENCES

- APHA., (1980). Standard methods for examination of water and waste water 15th Ed. N. Y. USA.
- Campbell, R. D., Leadem, T. P., Johnson, D. W., (1974). The invivo affect of DDT on Na⁺ K⁺ activated ATPase activity in rainbow trout (Salmo gaird neri). Bull. Env.Contom. Toxi. col., **11** 425–428.
- Carroll, N. V., Longley, R. W., Roe, J. H., (1956). Glycogen determination in liver and muscle by the use of anthrone. J. Biol. chem., **220**: 583–593.
- De Bruin, A., (1976). Biochemical toxicology of environmental agents. Elsevier. North Holland Biochemical press. N. Y. USA., Desaiah, D., Hayes, A. W., Ho, I. K., (1977). Effects of rubratoxin B on adenosine triophosphatase activities in the mouse. Toxical. Appl. Pharmacol., **39**: 71–79.
- Duncan, C. J., (1967). The molecular properties and evaluation if excitable cells. Pergamon press, oxford, *p*. 253.
- Eisler and Gardener, G. R., (1973). Acute toxicology to an estuarine teleost of mixtures of cadmium, copper and zinc salts. J. Fish. Biol., **5**: 131–142.
- Fiske, C., (1925). Subbarrow, Calarimetric determination of Phosporous. J. Biol. Chem., 68: 374–400.
- Francis S., Mohan, K. G., Oommen, V., (2002). Influence of steroid harmones on plasma proteins in fresh water Tilapia Oreochromis mossambicus. Indian J. Exper. Biol., 40: 1206–1208.
- Harrison, R., Lunt, G. G., (1980). Biochemical membrane: Their structure and function. Balckee, London, 75–93.
- Holcombe, G. W., Benoit, D. A., Leonard, E. N., Mekim, J. M., (1976). Long effects of lead exposureon three generation if brook trout, salvelinus fontianalis. J. Fish. Res. Bd. Canada., 33: 1731–1741.
- Jacobs, J. M., Carmicheal, N., Cavanagh, J. B., (1977). Ultra structural changes in the nervous system of rabbits poisoned with methyl mercury. Toxical. Appl. Pharmocol., 39: 249–261.
- Jagadeesan, G., Jebanesan, A., Mathivanan, A., (2001). In vivo recovery of organic constituents in gill tissue of Labco rohita after exposure to sublethal concentrations of mercury. J. Exp. India., 3: 22–29.
- Jagadeesan, G., Mathivanan, A., (1999). Organic constituents changes induced by three different sub-lethal concentrations of mercury and recovery in the liver tissue of labco rohita finegerlings. Poll. Res., **18**: 177–181.
- Jurss., (1980). The effect of changes in external salinity on the free amino acids and amino transferases of white muscle from fasted Salmo gairdneri. Comp. Biochem. Physiol., 65: 501–504.

- Kimellberg, H. K., Papahad, J., (1974). Effect of phospholipids acetyl chain fluidity phase transition and cholesterol in (Na⁺K⁺), stimulated adenosine triphosphatose. J. Biol. Chem., **249**: 1071–1080.
- Karamer, H. J., Meyer–Lehnert, Micheal, H., Predel, H.G., (1991). Endogenous nitriuretics and quabin like factors, their roles in body fluid volume and BP regulation. Am. J. Hypertens., 1: 81.
- Margarat, A., Jagadeesan, G., (1999). Abtodotal recovery ion carbohydrate metabolism of mercury in toxicated mice mus musculus (L.): Uttarpradesh. J. Zoo., 19: 196–198.
- Mary Candravathy, V. M., Reddy, S. L. N., (1991). Lead nitrate exposure changes in carbohydrate metabolism of fresh water fish. J. Env., Biol., **17**: 75.79.
- Neff, J. M., (1985). Use of biochemical measurement to detect polluatant medicated damage to fish. ASTM spec .Tech. publ., 854: 154–183
- Nuefeld, G. J., Prittchard, J. B., (1979). Osmoregulation and gill Na⁺K⁺ ATPhase in the rock crab, cancer irroratus: Response to D.D.T. Comp Biochem. Physiol., 620: 165– 172.
- Ramalingam, V., Arunadevi, R., (1999). Effect of Mercury chloride on testicular enzyme in adult Albino rat. Poll. Res., 18: 441–444.
- Richert, W. L., David, F. A. Malins, D. C., (1979). Uptake and metabolism of Lead and cadmium cohosalmon onchorhyncus Kisutch. Comp. Biochem. Physiol. Comp. Pharmacol., 63: 229–234.
- Schmit–Nielson, B., (1975). Comparitive physiology cellular ion and volume regulation. J. Exp. Zool., 194: 107–220.
- Shakoori, A. R., Javed Iqbal, M., Latif Mughal, A., Syed Shahid A., (1994). Bio chemical changes induced by inorganic mercury on the blood, liver and muscles of fresh Ctenopharyngodon idella. J. Ecotoxicol. Environ. Monit., 4:81–92.
- Sivaramakrishnan, B. and Radhakrishnaiyah, K., (1998). Impact of sublethal concentration of mercury on nitrogen metabolism of the freshwater fish, *Cyprinus carpio* (Linnaeus). J. Environ. Biol., **19**: 111–117.
- Sternlib.I and Goldfisher, S., (1976). Heavy metals and lysosomes. In: Lysosomes in biology pathology. (Ed.J.T.Dingle and R.T.Dean) American Elsevier company, N. Y. USA., 185–200.
- Sprague, J. B., (1969). Measurement of pollutant toxicity to fish I. Bioassay methods for acute toxity. Wat. Res., 8: 793–821.
- Villalan, P., Narayanan, K. R. Ajmal Khan, S. and Natarajan, R., (1988). Proximate composition of muscle, hepatopancrease and gill in the copper exposed estuarine crab Thalamita crenata (Latreille). Proceeding of the second national symposium on eco toxicology, Annamalai University, India.
- Von Westernhagan, H., (1988). Sub lethal effects a pollutant on fish eggs and larvae. In fish physiology VOL.XI PART-A, Academic press N. Y. USA., 253–346.
- Yamaguchi, F., Matsumara and Kadous, A. A., (1979). Hepatachlor epoxide : Effects of Ca medicated transmitter release from brain synaptosomes in rat. Biochem. Pharmacol., 29: 1815–1823.