

Original Article

The Effect of Aerobic Training and Vitamin D Supplementation on Blood Pressure in Elderly Women with Nonalcoholic Fatty Liver and Vitamin D Deficiency

Zahra Hoseini, Nasser Behpour*, Rastegar Hoseini

Department of Exercise Physiology, Faculty of Sport Sciences, Razi University, Kermanshah, Iran

Received: 11 May 2019

Accepted: 10 Jun 2019

Abstract

Background & Objective: The aim of this study was to investigate the effects of aerobic training (AT) and vitamin D (Vit D) supplementation on cardiovascular variables in elderly women with nonalcoholic fatty liver disease (NAFLD) and vitamin D deficiency.

Materials & Methods: A total of forty women (60-65 years) with NAFLD (second or third grade) and vitamin D deficiency were recruited for the study. Participants were randomly assigned to four AT+Vit D, AT, Vit D, and control (C) groups and were allocated to receive either vitamin D (~50000 IU • week for eight weeks) or placebo. The aerobic training protocol consisted of 40-60 minutes of aerobic training at 60% - 75% of HRmax, 3 times a week for eight weeks. Anthropometric indices along with systolic blood pressure (SBP), diastolic blood pressure (DBP), and heart rate (HR) were measured and mean of arterial pressure (MAP) was calculated.

Results: AT+Vit D, Vit D and AT groups demonstrated a significant decrease in SBP, DBP, and HR, ($P < 0.05$) while, C group showed a significant increase in the mentioned variables ($P < 0.05$). Also after 8 weeks, AT + Vit D and AT groups showed significant decrease in the mean of MAP ($P = 0.001$; $P = 0.018$). Compared to the control group, significant differences in the SBP, DBP, and HR were observed in all groups. The mean SBP was significantly lower in AT+Vit D group comparing with AT, Vit D, and C groups. Also, no significant differences were observed between AT and Vit D groups in all variables.

Conclusion: AT developed significant improvement in cardiovascular variables indicating SBP, DBP, MAP, and HR in elderly women with NAFLD and vitamin D deficiency. Meanwhile, AT combined with Vit D decreased SBP more significantly that might be incorporated in the management programs of the patients suffering from NAFLD in order to augment improvement in their blood pressure.

Keywords: Exercise, Vitamin D, Blood Pressure, NAFLD

Introduction

Currently, the elderly represent a large and growing portion of the population (1). According to the 2011 census in Iran, 7.7% of the total population of the country aged over 65, which is projected to increase by 25 to 35% by

2031 (2). The older one becomes, the higher the prevalence of chronic diseases like Non-Alcoholic Fatty Liver Disease (NAFLD) especially in women (3). NAFLD is the leading cause of chronic liver disease, with increased prevalence up to 20–30% worldwide (4). It is characterized by the accumulation of fat more than 5% of the hepatocytes in the liver (5), which encompasses various conditions ranging from simple steatosis, nonalcoholic steatohepatitis, to

*Corresponding Author: Behpour Nasser, Department of Exercise Physiology, Faculty of Sport Sciences, Razi University, Kermanshah, Iran. Email: n_behpour@yahoo.com
<https://orcid.org/0000-0003-4987-9872>

cirrhosis (6). Recent studies suggested that NAFLD is an independent risk factor for hypertension, the major cause of Cardiovascular disease (CVD), and its progression is associated with the development of hypertension (7). The deficiency of vitamin D has been implicated as the predictive factor for the occurrence of NAFLD (8, 9). Furthermore, Vitamin D deficiency is known to be associated with hypertension, obesity, atherosclerosis, type 2 diabetes mellitus, oxidative stress, cancer and CVD (10-13). Moreover, studies suggest that a low level of vitamin D is associated with high blood pressure. They also revealed that a high chance of developing hypertension is correlated with the increased risk of mortality and high levels of cardiovascular events in the future (12, 14, 15). It has been reported that vitamin D regulates calcium and bone homeostasis, and its receptor is present in many tissues, such as vascular endothelium and myocardial tissue. This indicates the biological role and function of vitamin D, which can suppress renin production by affecting the juxtaglomerular system (16, 17). Vitamin D supplementation can lead to the suppression of parathyroid hormone and pro-inflammatory cytokine production. In addition, there are possible pathophysiological pathways that show the effects of vitamin D on vascular protection against hypertension (18). A number of previous studies associated with NLFAD have reported that regular aerobic training reduces hepatic fat content in NLFAD patients, and helps control hypertension (19, 20). Among the principal physiological benefits of physical training in hypertensive patients, there are reductions in heart rate (HR) and peripheral vascular resistance (21). These factors are related to the reduction of systolic and diastolic blood pressure (SBP and DBP) (22). In addition to the improvements in blood pressure, aerobic training is associated with a reduction of serum cholesterol, which often is above the reference values in hypertensive subjects (23). Generally, there are contradictory results about the effect of aerobic training on hypertension. Some studies indicate beneficial effects (24, 25) and others report no significant changes (26, 27). In addition, limited amount of studies have been conducted to show the effects of vitamin D supplementation on blood pressure in NAFLD. On the other hand, to our knowledge, there is no information in the literature comparing the separate and interactional effects of aerobic

training and vitamin D supplementation. Since the decrease in physical activity level and the high prevalence of vitamin D deficiency and NAFLD are more significant in elderly women, the present study intends to investigate the effects of aerobic training and vitamin D supplementation on hypertension in elderly women with NAFLD and vitamin D deficiency.

Materials & Methods

This is a semi-experimental research with pre-test and post-test design. The target population of this study was elderly women (60 to 65 years) of Kermanshah with NAFLD and vitamin D deficiency. Forty Subjects were divided into four groups including: Aerobic training (AT, n=10); Vitamin D supplementation (Vit D, n=10); Aerobic training with vitamin D supplements (AT+Vit D, n=10); and Control (C, n=10). Inclusion criteria includes: ultrasound (US) confirmation of being in second or third grade of NAFLD and certified by an internist, 25-OHD levels between 10 and 20 ng/ml, not having an specific diet and regular exercise program in the past year. Exclusion criteria includes: significant consumption of alcohol (more than 20 gr per day) (28, 29), presence of other liver diseases (B and C hepatitis), other disorders such as autoimmune hepatitis, joint disease, celiac and Wilson disease, CVD, kidney failure, hypothyroidism, also surgical treatment of obesity or severe weight loss, and the presence of any other chronic disease or skeletal disorders (29).

Three days prior to the start of the study, explanations about research conduction were provided for subjects and consent forms were signed by all 40 subjects. Also, PARQ, 3-day dietary questionnaire, nutritional questionnaires including food history, food frequency, habits and dietary behaviors, and Health History Questionnaire (HHQ) were completed. On the first day, height was measured with an accuracy of 0.5 cm. Body weight (BW, with an accuracy of 0.1 gr), BMI, body fat percentage (BFP), were obtained using the INBODY test in the fasting state. Also waist-hip ratio (WHR) was measured by dividing waist circumference by hip circumference according to WHO criteria.

Ethical considerations

The present study was approved by the Committee of Ethics in Research at Kermanshah University of Medical Sciences and registered in the Iranian Clinical Trial Registration Center under the code IR.KUMS.REC.1397.1059.

Before the study, informed written consent was obtained from all participants after a comprehensive oral and written explanation of the study was provided to them. All patients were allowed to leave the study at any point voluntarily. The demographic data of the participants were kept confidential. Also, all the clinical and paraclinical tests of the study were free of charge, with no costs to the patients.

Training program

The aerobic training program was executed 3 days/week for 8 weeks based on the ACSM's recommendation. Each session lasted 40-60 min consisting 10 minutes of warming-up, 20 to 40 minutes of aerobic exercise and 10 minutes of cooling-down. In the first week, subjects performed 20-min of aerobic exercise including: walking, jogging and running at 60% HR_{max}. The initial duration and intensity was gradually progressed to a maximum duration of 40 min and 75% HR_{max} respectively. The training program was performed with 60-75% of the individual HR_{max} (using Karvonen Formula) and 10-13 of RPE in 6-20 Borg scale (Table 1) (30).

caffeinated products and exercising, 30 min prior to the BP measurement. Throughout assessment, patients remained in a relaxed seated state for 10 min. The cuff was placed on the left arm for all patients and BP was recorded according to standard guidelines (31). A second recording was also taken after 2 min on the same arm. If the measurements had a difference of ≥ 5 mmHg of BP, further recording were obtained until there were 2 consecutive stable measurements. Final recording was considered as an average of the 2 stable measurements obtained. Measurement took place each day between 8.00 a.m. and 10.00 a.m. in accordance with the protocol of the American Heart Association (31) Mean arterial pressure (MAP) was calculated as $DBP + [0.333(SBP - DBP)]$.

Data Analysis

Data are presented as means \pm SD. Prior to analysis, data normality was checked with the Shapiro-Wilk test. In order to compare the mean SBP and DBP between and within groups, ANOVA and t-test were used, respectively. Tukey's post hoc test was used if significant

Table 1. Exercise protocol program

Variables	Weeks							
	First	Second	Third	Fourth	Fifth	Sixth	Seventh	Eight
Intensity (HR _{max})	60%	60%	65%	65%	70%	70%	75%	75%
Time (min)	20	25	25	30	30	35	35	40
Borg scale	10	10	11	11	11	12	12	13

Vitamin D supplementation

In this study, both AT+Vit D group and Vit D group received 50000 IU of vitamin D supplement once per week at the beginning of the week, made by the Zahravi Pharmaceutical Company in Iran. The C and AT groups also received a weekly placebo (a paraffin made by the Zahravi Pharmaceutical Company, Iran) with the same shape, color, smell and tastes to the vitamin D supplements pills, over a period of 8 weeks.

Blood pressure measurements

SBP and DBP were measured by standard manual sphygmomanometer. Participants were strictly prohibited from consuming any

differences were found. The SPSS software version 21 was used at a significance level of $P < 0.05$.

Results

The anthropometric indices of groups are presented in Table 2. There was no statistically significant difference between the four groups at the beginning of the study in weight, BMI, body fat and WHR.

Based on the results of t-test, there were significant differences in the mean of SBP, DBP, and HR between the pre-test and post-test conditions. After 8 weeks SBP, DBP and HR

significantly decreased in AT+Vit D, AT, Vit D groups while, in the control group a significant increase in SBP, DBP and HR was observed (Table 3). The results also

showed that after 8 weeks, the mean of MAP in AT + Vit D and AT groups decreased significantly ($P = 0.001$; $P = 0.018$, respectively) however, there was no

Table 2. Mean \pm SD and frequency distribution of the anthropometric indices before the intervention among the groups

Variables	AT+Vit D (n = 10)	AT (n = 10)	Vit D (n=10)	C (n = 10)	P Value
Age (years)	63.10 \pm 2.37	62.60 \pm 1.89	61.30 \pm 1.41	62 \pm 1.88	0.16
Height (cm)	157.10 \pm 5.25	160 \pm 5.45	158.30 \pm 4.59	159 \pm 5.79	0.11
Weight (kg)	87.40 \pm 4.64	85.80 \pm 3.35	86.10 \pm 3.14	87.50 \pm 4.19	0.679
BMI (kg/h ²)	35.55 \pm 3.55	33.57 \pm 1.92	34.40 \pm 1.78	34.65 \pm 1.70	0.332
Body Fat (%)	44 \pm 3.36	43.10 \pm 3.41	41.20 \pm 3.15	42 \pm 3.59	0.284
WHR	94.50 \pm 2.54	95.20 \pm 3.79	95.80 \pm 2.78	95.70 \pm 3.16	0.778

AT+Vit D group, Aerobic training and vitamin D supplementation, AT group, Aerobic training; Vit D group, vitamin D supplementation; C group, that had neither aerobic training nor vitamin D supplementation. P values are calculated using one-way analysis of variance test followed by Tukey's Post hoc Test.

Table 3. Comparison of Mean \pm SD of cardiovascular variables within and between the groups under the study

Variables	AT + Vit D (n = 10)	AT (n = 10)	Vit D (n=10)	C (n = 10)	P Value ^a
SBP (mmHg)					
Before	129.60 \pm 2.50	130.30 \pm 1.56	128.90 \pm 1.79	129.40 \pm 1.77	0.454
After	125.40 \pm 1.34	127.40 \pm 1.64	127.80 \pm 1.75	131.10 \pm 1.37	0.001 ¥
P Value ^b	P=0.001 *	P=0.002 *	P=0.001 *	P=0.002 *	
DBP (mmHg)					
Before	84.40 \pm 1.34	83.40 \pm 1.42	84.20 \pm 1.03	84.10 \pm 1.28	0.343
After	82.90 \pm 1.37	82.40 \pm 1.40	83.30 \pm 1.41	85.30 \pm 0.94	0.001 ¥
P Value ^b	P=0.001 *	P=0.023 *	P=0.002 *	P=0.002 *	
MAP (mmHg)					
Before	99.45 \pm 1.25	99.01 \pm 1.03	99.08 \pm 0.94	99.18 \pm 1.10	0.821
After	98.55 \pm 1.01	98.38 \pm 1.15	99.01 \pm 0.83	99.35 \pm 1.07	0.155
P Value ^b	0.001 *	0.018 *	0.343	0.096	
HR (mmHg)					
Before	85 \pm 1.33	83.40 \pm 1.42	84.20 \pm 1.03	84.10 \pm 1.19	0.059
After	79.60 \pm 1.42	79.70 \pm 1.15	80.70 \pm 1.63	86.40 \pm 0.84	0.001 ¥
P Value ^b	P=0.001 *	P=0.001 *	P=0.001 *	P=0.001 *	

AT+Vit D group: Aerobic training and vitamin D supplementation; AT group: Aerobic training; Vit D group: vitamin D supplementation; C group: that had neither aerobic training nor vitamin D supplementation; SBP: systolic blood pressure; DBP: diastolic blood pressure; MAP: mean of arterial pressure; HR: heart rate.

P values with superscript "a" are calculated using one-way analysis of variance test followed by Tukey's post hoc test; superscript letter "b" indicates values were calculated using paired t-test

*Significantly different in comparison with pre and posttest within the groups; ¥ significantly different in comparison with pre and posttest between groups.




Figure 1 (a). Pre-test and post-test value of the SBP among groups




Figure 1 (b). Pre-test and post-test value of the DBP among groups




Figure 1 (c). Pre-test and post-test value of the MAP among groups




Figure 1 (d). Pre-test and post-test value of the HR among groups

AT+Vit D group: Aerobic training and vitamin D supplementation; AT group: Aerobic training; Vit D group: vitamin D supplementation; C group: that had neither aerobic training nor vitamin D supplementation; SBP: systolic blood pressure; DBP: diastolic blood pressure; MAP: mean of arterial pressure; HR: heart rate.

*: Significantly different in comparison with pre and posttest groups; ‡: Significant difference between AT+Vit D group compare to AT group in posttest; ¥: Significant difference between AT+Vit D group compared to Vit D group in posttest; ‡: Significant difference between AT+Vit D, AT, and Vit D groups compared to C group.

significant difference in the mean of MAP in Vit D and C groups ($P = 0.343$; $P = 0.096$).

The results of one-way ANOVA showed no significant difference in SBP, DBP, MAP and HR between the groups in the pre-test, however in the post-test there was a significant difference in the SBP, DBP and HR between the groups (Table 3). Although, there was no significant difference in the mean of MAP between groups in the post-test ($P = 0.155$).

The results of Tukey's post hoc test are presented in Figure 1 (a,b,c,d). The lowest levels of SBP, DBP, HR and MAP were in the AT+Vit D group. The results also showed that there was a significant difference in SBP levels between the AT+Vit D group with AT group, Vit D group and control group; However, there was no significant difference between the AT+Vit D group with AT group and Vit D group in the DBP, HR and MAP. The results also showed a significant difference in the SBP, DBP and HR levels between AT+Vit D group, the AT group, and the Vit D group compared to the control group.

Discussion

This study is the first to assess the effect of aerobic training and vitamin D supplementation on hypertension in elderly women with nonalcoholic fatty liver and vitamin D deficiency. Since increased SBP, DBP, HR and MAP are risk factors for the development of CVD in elderly women, thus identifying effective programs in reducing SBP, DBP, HR and MAP has always been a controversial issue. Our study observed high prevalence of hypertension in patients with NAFLD. Based on the results of the studies, after 8 weeks SBP, DBP and HR significantly decreased in AT+Vit D, AT, Vit D groups while, in the control group a significant increase in SBP, DBP and HR were observed.

There are limited consensus in the literature as to how aerobic training affects cardiovascular variables (SBP, DBP, MAP and HR) in elderly women with NAFLAD patients. Most studies indicate a reduction of BP values (32, 33) and HR (34) during rest after aerobic training in this population. However, other authors conducting similar studies did not observe significant

changes in these parameters (26, 27). Among the factors reported, the main contributors to this disagreement seem to be hypertension stage in addition to the intensity, duration, and frequency of the aerobic training program (35). Our results indicated that elderly women had significant reductions in SBP, DBP, MAP and HR after the completion of an 8-week aerobic training program. A similar result for BP was shown by Lima et al (2011) with significant reductions in both systolic (142.7 ± 6.3 vs. 130.8 ± 5.8 mmHg, $P < 0.001$) and diastolic BP (87.0 ± 4.5 vs. 81.9 ± 4.3 mmHg, $P = 0.002$) in just 30 days of aerobic training with three sessions per week of moderate intensity (50-70% HR reserve + HR rest) exercise (36). Hypertension is multifactorial, and several mechanisms may be involved in the decrease in BP and bradycardia induced by aerobic training. The related chronic adaptations include improvement of endothelial function, (37) the reduction of sympathetic nerve activity, increased baroreflex sensitivity, (22) improvement in the hyperinsulinism state, (38) improvement in lipoprotein profile, (38) weight loss, (39) reduced cardiac output, (39) and reduced peripheral vascular resistance (PVR) (40). In the elderly, the main mechanism involved in the reduction of resting blood pressure in response to aerobic physical training seems to be a reduction in PVR (41).

On the other hand, in the view of the bile acid-dependent uptake of vitamin D and its hepatic metabolism, it is reasonable to expect an association between vitamin D status and both cholestatic and non-cholestatic chronic liver disease. Recently, a significant correlation between lower 25(OH) vitamin D levels and an increasing stage of fibrosis and severity of necro-inflammatory activity was observed in a population with genotype 1 chronic hepatitis C (42). Interestingly, reduced 25(OH) vitamin D levels have also been noted in patients with NAFLD. A study by Targher et al (2007) found a strong association between vitamin D deficiency and NAFLD with a close association of vitamin D levels with the histological severity of hepatic steatosis, necro-inflammation, and fibrosis (8). 25(OH) vitamin D levels were inversely associated with NAFLD independent of hypertension, suggesting that inadequate vitamin D status might contribute to the development and progression of NAFLD (8).

Results of the present study demonstrate that the lowest levels of SBP, DBP, HR and MAP

after 2 months were in the AT+Vit D group. Limited studies have been conducted on the relationship between vitamin D intake and hypertension. Martins et al (2007) have shown that low Vitamin D levels are associated with a higher risk of having hypertension (43). Pfeifer et al. showed that 800 IU of Vitamin D3 plus 1200 mg of calcium significantly reduces blood pressure after 8 weeks compared to the group receiving 1200 mg of calcium alone (44) Vitamin D has also been shown to play a critical role in the regulation of renin-angiotensin system and thus has an influence on the regulation of blood pressure (45, 46). It has also been demonstrated that calcium regulating hormones and renin-angiotensin system coordinately mediate blood pressure regulation via altering cellular concentrations of sodium and calcium ions. Thus, it is clear that Vitamin D supplementation has a role in the regulation of blood pressure and that it should be supplemented with the antihypertensive drugs to the patients with hypertension.

One of the limitations of our study was that the diagnosis of NAFLD in our patients was based on the presence of hepatic steatosis on ultrasonography and exclusion of other causes of hepatic steatosis and raised ALT without a histological diagnosis. However, in the absence of liver biopsy, the presence of significant fibrosis and the severity of liver disease were evaluated by transient elastography in our patients, which is now an established modality in patients with NAFLD. Even though the number of patients with raised ALT was small in the study, we could demonstrate significant improvement in ALT in the group that was given vitamin D.

Conclusions

In conclusion, our study demonstrates that low physical activity and vitamin D deficiency is common in patients with NAFLD as it is in the general population. Aerobic training and vitamin D supplementation are a suitable intervention leading to an improvement in the cardiovascular variables. In general terms, our results reinforce the recommendations available in the literature about non-pharmacological treatment of hypertension with planned aerobic training programs in the elderly with NAFLD and vitamin D deficiency.

More studies are needed to verify the effectiveness of different physical training

programs and different doses of vitamin D on cardiovascular and metabolic parameters, specifically for the elderly hypertensive population with NAFLD.

Acknowledgments

This article was part of an MSc thesis at the Department of Exercise Physiology, Faculty of Sport Sciences, Razi University, Kermanshah, Iran. This study was supervised at the Ethics Committee of the Kermanshah University of medical science, and was approved with ID: IR.KUMS.REC.1397.1059, date: March 6, 2019. The authors would like to thank the subjects for their willing participation in this study.

Conflict of Interests

The authors declare no competing interests.

Reference

1. Braz NF, Carneiro MV, Oliveira-Ferreira F, Arrieiro AN, Amorim FT, Lima MM, et al. Influence of aerobic training on cardiovascular and metabolic parameters in elderly hypertensive women. *International journal of preventive medicine*. 2012;3(9):652.
2. Esteghamati A, Etemad K, Koohpayehzadeh J, Abbasi M, Meysamie A, Noshad S, et al. Trends in the prevalence of diabetes and impaired fasting glucose in association with obesity in Iran: 2005–2011. *Diabetes research and clinical practice*. 2014;103(2):319-27.
3. Hill NR, Fatoba ST, Oke JL, Hirst JA, O'Callaghan CA, Lasserson DS, et al. Global prevalence of chronic kidney disease—a systematic review and meta-analysis. *PloS one*. 2016;11(7):e0158765.
4. Vernon G, Baranova A, Younossi Z. Systematic review: the epidemiology and natural history of non-alcoholic fatty liver disease and non-alcoholic steatohepatitis in adults. *Alimentary pharmacology & therapeutics*. 2011;34(3):274-85.
5. Torres DM, Williams CD, Harrison SA. Features, diagnosis, and treatment of nonalcoholic fatty liver disease. *Clinical Gastroenterology and Hepatology*. 2012;10(8):837-58.
6. Rivera-Andrade A, Kroker-Lobos MF, Lazo M, Freedman ND, Smith JW, Torres O, et al. High prevalence of non-alcoholic fatty liver disease and metabolic risk factors in Guatemala: A population-based study. *Nutrition, Metabolism and Cardiovascular Diseases*. 2019;29(2):191-200.
7. Ryoo JH, Suh YJ, Shin HC, Cho YK, Choi JM, Park SK. Clinical association between non-alcoholic fatty liver disease and the development of hypertension. *Journal of gastroenterology and hepatology*. 2014;29(11):1926-31.
8. Targher G, Bertolini L, Scala L, Cigolini M, Zenari L, Falezza G, et al. Associations between serum 25-hydroxyvitamin D3 concentrations and liver histology in patients with non-alcoholic fatty liver disease. *Nutrition, Metabolism and Cardiovascular Diseases*. 2007;17(7):517-24.
9. Arteh J, Narra S, Nair S. Prevalence of vitamin D deficiency in chronic liver disease. *Digestive diseases and sciences*. 2010;55(9):2624-8.
10. Al Mheid I, Patel R, Murrow J, Morris A, Rahman A, Fike L, et al. Vitamin D status is associated with arterial stiffness and vascular dysfunction in healthy humans. *Journal of the American College of Cardiology*. 2011;58(2):186-92.
11. Gotsman I, Shauer A, Zwas DR, Hellman Y, Keren A, Lotan C, et al. Vitamin D deficiency is a predictor of reduced survival in patients with heart failure; vitamin D supplementation improves outcome. *European journal of heart failure*. 2012;14(4):357-66.
12. Costanzo S, De Curtis A, Di Castelnuovo A, Persichillo M, Bonaccio M, Pounis G, et al. Serum vitamin D deficiency and risk of hospitalization for heart failure: Prospective results from the Moli-sani study. *Nutrition, Metabolism and Cardiovascular Diseases*. 2018;28(3):298-307.
13. Chandler PD, Scott JB, Drake BF, Ng K, Manson JE, Rifai N, et al. Impact of vitamin D supplementation on inflammatory markers in African-Americans: results of a four-arm, randomized, placebo-controlled trial. *Cancer prevention research*. 2013;canprevres. 0338.2013.
14. Ahmad YK, El-Ghamry EM, Tawfik S, Atia WM, Keder MM, Abd-El Kader SA. Assessment of Vitamin D Status In Patients With Essential Hypertension. *The Egyptian Journal of Hospital Medicine (July 2018)*. 2018;72(5):4434-8.

15. Grecksch G, Rüttrich H, Höllt V, Becker A. Transient prenatal vitamin D deficiency is associated with changes of synaptic plasticity in the dentate gyrus in adult rats. *Psychoneuroendocrinology*. 2009;34:S258-S64.
16. Yuan W, Pan W, Kong J, Zheng W, Szeto FL, Wong KE, et al. 1, 25-dihydroxyvitamin D3 suppresses renin gene transcription by blocking the activity of the cyclic AMP response element in the renin gene promoter. *Journal of Biological Chemistry*. 2007;282(41):29821-30.
17. Grübler MR, März W, Pilz S, Grammer TB, Trummer C, Müllner C, et al. Vitamin-D concentrations, cardiovascular risk and events—a review of epidemiological evidence. *Reviews in Endocrine and Metabolic Disorders*. 2017;18(2):259-72.
18. Rodriguez AJ, Mousa A, Ebeling PR, Scott D, de Courten B. Effects of vitamin D supplementation on inflammatory markers in heart failure: A systematic review and meta-analysis of randomized controlled trials. *Scientific reports*. 2018;8(1):1169.
19. Conraads VM, Pattyn N, De Maeyer C, Beckers PJ, Coeckelberghs E, Cornelissen VA, et al. Aerobic interval training and continuous training equally improve aerobic exercise capacity in patients with coronary artery disease: the SAINTEX-CAD study. *International journal of cardiology*. 2015;179:203-10.
20. Punia S, Kulandaivelan S, Singh V, Punia V. Effect of aerobic exercise training on blood pressure in Indians: Systematic review. *International journal of chronic diseases*. 2016;2016.
21. Collier S, Kanaley J, Carhart Jr R, Frechette V, Tobin M, Hall A, et al. Effect of 4 weeks of aerobic or resistance exercise training on arterial stiffness, blood flow and blood pressure in pre- and stage-1 hypertensives. *Journal of human hypertension*. 2008;22(10):678.
22. dos Santos Júnior EA, Sales JA, Brasileiro-Santos S, Barbosa BT. High-Intensity Interval Aquatic Exercise Session Promotes Post-Exercise Hypotension in Hypertensive Elderly: A Randomized Controlled Trial. *Journal of Exercise Physiology Online*. 2018;21(1):149-62.
23. Khan K, Jovanovski E, Ho H, Marques A, Zurbau A, Mejia S, et al. The effect of viscous soluble fiber on blood pressure: A systematic review and meta-analysis of randomized controlled trials. *Nutrition, Metabolism and Cardiovascular Diseases*. 2018;28(1):3-13.
24. Freire YA, da Silva CA, de Sousa FJR, Browne RAV, Farias-Junior LF, Schwade D, et al. A single multi-joint high-intensity resistance exercise involving large muscle groups elicits post-exercise hypotension in normotensive-trained women: a crossover trial. *Sport Sciences for Health*. 2018;14(1):127-34.
25. Souza LR, Vicente JB, Melo GR, Moraes VC, Olher RR, Sousa IC, et al. Acute hypotension after moderate-intensity handgrip exercise in hypertensive elderly people. *The Journal of Strength & Conditioning Research*. 2018;32(10):2971-7.
26. Church TS, Earnest CP, Skinner JS, Blair SN. Effects of different doses of physical activity on cardiorespiratory fitness among sedentary, overweight or obese postmenopausal women with elevated blood pressure: a randomized controlled trial. *Jama*. 2007;297(19):2081-91.
27. Stewart KJ, Bacher AC, Turner KL, Fleg JL, Hees PS, Shapiro EP, et al. Effect of exercise on blood pressure in older persons: a randomized controlled trial. *Archives of internal medicine*. 2005;165(7):756-62.
28. Al-Jiffri O, Al-Sharif F, El-Kader S, Ashmawy E. Weight reduction improves markers of hepatic function and insulin resistance in type-2 diabetic patients with non-alcoholic fatty liver. *African health sciences*. 2013;13(3):667-72.
29. Johnson NA, Sachinwalla T, Walton DW, Smith K, Armstrong A, Thompson MW, et al. Aerobic exercise training reduces hepatic and visceral lipids in obese individuals without weight loss. *Hepatology*. 2009;50(4):1105-12.
30. Heil DP. ACSM's guidelines for exercise testing and prescription. *Medicine and Science in Sports and Exercise*. 2001;33(2):343.
31. Townsend RR, Wilkinson IB, Schiffrin EL, Avolio AP, Chirinos JA, Cockcroft JR, et al. Recommendations for improving and standardizing vascular research on arterial stiffness: a scientific statement from the American Heart Association. *Hypertension*. 2015;66(3):698-722.
32. Moraes M, Bacurau R, Simões H, Campbell C, Pudo M, Wasinski F, et al. Effect of 12 weeks of resistance exercise on post-exercise hypotension in stage 1 hypertensive individuals. *Journal of human hypertension*. 2012;26(9):533.

33. Miller 3rd ER, Erlinger TP, Young DR, Jehn M, Charleston J, Rhodes D, et al. Results of the Diet, Exercise, and Weight Loss Intervention Trial (DEW-IT). *Hypertension*. 2002;40(5):612-8.
34. Farinatti P, Monteiro WD, Oliveira RB. Long term home-based exercise is effective to reduce blood pressure in low income Brazilian hypertensive patients: a controlled trial. *High Blood Pressure & Cardiovascular Prevention*. 2016;23(4):395-404.
35. Sisson SB, Katzmarzyk PT, Earnest CP, Bouchard C, Blair SN, Church TS. Volume of exercise and fitness non-response in sedentary, post-menopausal women. *Medicine and science in sports and exercise*. 2009;41(3):539.
36. Lima MMO, Britto RR, Baião EA, Alves GdS, Abreu CDG, Parreira VF. Aerobic exercises on the control of post menopause hypertension. *Fisioterapia em Movimento*. 2011;24(1):23-31.
37. Scapini KB, Bohlke M, Moraes OA, Rodrigues CG, Inácio JF, Sbruzzi G, et al. Combined training is the most effective training modality to improve aerobic capacity and blood pressure control in people requiring haemodialysis for end-stage renal disease: systematic review and network meta-analysis. *Journal of physiotherapy*. 2019;65(1):4-15.
38. Widlansky ME, Gokce N, Keaney JF, Vita JA. The clinical implications of endothelial dysfunction. *Journal of the American College of Cardiology*. 2003;42(7):1149-60.
39. Fahlman MM, Boardley D, Lambert CP, Flynn MG. Effects of endurance training and resistance training on plasma lipoprotein profiles in elderly women. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*. 2002;57(2):B54-B60.
40. Pedersen BK. Anti-inflammatory effects of exercise: role in diabetes and cardiovascular disease. *European journal of clinical investigation*. 2017;47(8):600-11.
41. Cardoso Jr CG, Gomides RS, Queiroz ACC, Pinto LG, Lobo FdS, Tinucci T, et al. Acute and chronic effects of aerobic and resistance exercise on ambulatory blood pressure. *Clinics*. 2010;65(3):317-25.
42. Petta S, Cammà C, Scazzone C, Tripodo C, Di Marco V, Bono A, et al. Low vitamin D serum level is related to severe fibrosis and low responsiveness to interferon-based therapy in genotype 1 chronic hepatitis C. *Hepatology*. 2010;51(4):1158-67.
43. Martins D, Wolf M, Pan D, Zadshir A, Tareen N, Thadhani R, et al. Prevalence of cardiovascular risk factors and the serum levels of 25-hydroxyvitamin D in the United States: data from the Third National Health and Nutrition Examination Survey. *Archives of internal medicine*. 2007;167(11):1159-65.
44. Pfeifer M, Begerow B, Minne HW, Nachtigall D, Hansen C. Effects of a short-term vitamin D3 and calcium supplementation on blood pressure and parathyroid hormone levels in elderly women. *The Journal of Clinical Endocrinology & Metabolism*. 2001;86(4):1633-7.
45. Jorge AJL, Cordeiro JR, Rosa MLG, Bianchi DBC. Vitamin D Deficiency and Cardiovascular Diseases. *International Journal of Cardiovascular Sciences*. 2018;31(4):422-32.
46. Cremer A, Tambosco C, Corcuff J-B, Boulestreau R, Gaillard P, Lainé M, et al. Investigating the association of vitamin D with blood pressure and the renin-angiotensin-aldosterone system in hypertensive subjects: a cross-sectional prospective study. *Journal of human hypertension*. 2018;32(2):114.

مقاله پژوهشی

تأثیر تمرینات هوازی و مکمل‌دهی ویتامین D بر فشار خون در زنان سالمند مبتلا به کبد چرب غیرالکلی و کمبود ویتامین D

زهرا حسینی، ناصر بهپور*، رستگار حسینی

گروه فیزیولوژی ورزشی، دانشگاه رازی، کرمانشاه، ایران

تاریخ پذیرش مقاله: ۱۳۹۸/۰۳/۲۰

تاریخ دریافت مقاله: ۱۳۹۸/۰۲/۲۱

چکیده

زمینه و هدف: این مطالعه به منظور تعیین تأثیر تمرینات هوازی و مکمل‌دهی ویتامین D بر متغیرهای قلبی-عروقی در زنان سالمند مبتلا به بیماری کبد چرب غیر الکلی (NAFLD) و کمبود ویتامین D انجام گردید. مواد و روش‌ها: در مجموع ۴۰ نفر از زنان مبتلا به NAFLD (مرحله ۲ و ۳) و کمبود ویتامین D برای مطالعه انتخاب شدند. شرکت کنندگان به صورت تصادفی به ۴ گروه تمرین هوازی+ مکمل‌دهی ویتامین D (AT + Vit D)، تمرین هوازی (AT)، مکمل‌دهی ویتامین D (Vit D) و کنترل (C) تقسیم شدند، همچنین دریافت ویتامین D (حدود ۵۰۰۰ IU در هفته به مدت هشت هفته) و یا دارونما مشخص گردید. پروتکل تمرین هوازی شامل ۴۰ تا ۶۰ دقیقه تمرین هوازی با شدت ۶۰ تا ۷۵ درصد ضربان قلب بیشینه، ۳ بار در هفته به مدت هشت هفته بود. شاخص‌های تن سنجی همراه با فشار خون سیستولیک (SBP)، فشار خون دیاستولیک (DBP) و ضربان قلب (HR) اندازه‌گیری شد و میانگین فشار شریانی (MAP) محاسبه شد. نتایج: گروه‌های AT + Vit D، Vit D و AT کاهش معنی‌داری در متغیرهای SBP، DBP و HR داشتند ($P < 0.05$)، در حالی که متغیرهای ذکر شده در گروه C به طور معنی‌داری افزایش یافت ($P < 0.05$). همچنین پس از ۸ هفته در گروه‌های AT + Vit D و AT کاهش معنی‌داری در میانگین MAP ($P = 0.01$ ؛ $P = 0.01$) مشاهده شد. در همه گروه‌ها در مقایسه با گروه کنترل اختلاف معنی‌داری در متغیرهای SBP، DBP و HR مشاهده شد. میانگین SBP در گروه AT + Vit D به طور معنی‌داری کمتر از گروه‌های AT، Vit D و C بود. همچنین بین گروه‌های AT و Vit D اختلاف معنی‌داری مشاهده نشد. نتیجه‌گیری: تمرینات هوازی در زنان سالمند مبتلا به NAFLD و کمبود ویتامین D باعث بهبود معنی‌داری در متغیرهای قلبی-عروقی شامل SBP، DBP، MAP و HR شد. علاوه بر این، تمرینات هوازی در ترکیب با ویتامین D باعث کاهش بیشتری در SBP شد و می‌تواند به منظور بهبود فشار خون در برنامه‌ی کنترل بیماران مبتلا به NAFLD گنجانده شود.

کلمات کلیدی: فعالیت ورزشی، ویتامین D، بیماری کبد چرب غیر الکلی، فشارخون

*نویسنده مسئول: ناصر بهپور، گروه فیزیولوژی ورزشی، دانشگاه رازی، کرمانشاه، ایران

Email: n_behpoor@yahoo.com

<https://orcid.org/0000-0003-4987-9872>