

Nano Silica Phosphoric Acid: An Efficient Catalyst for the One-Pot Synthesis of 1, 2, 4, 5-Tetrasubstituted Imidazoles

A. Bamoniri^{*a}, B. F. Mirjalili^b, S. Nazemian^a

^a*Department of Organic Chemistry, Faculty of Chemistry, University of Kashan, Kashan, I. R. Iran.*

^b*Department of Chemistry, College of Science, Yazd University, I. R. Iran .*

Article history:

Received 2/ 2/2012

Accepted/14/5/2012

Published online 1/6/2012

Keywords:

Tetrasubstitutedimidazoles

Multi-component reactions

Nano silica

Phosphoric acid

Sonication condition

**Corresponding author:*

E-mail address:

bamoniri@kashanu.ac.ir

Phone: +98-3615912305

Fax: +98 21 88006076

Abstract

Two simple protocols for the synthesis of 1, 2, 4, 5-tetrasubstitutedimidazoles using nano-SPA as a reusable, eco-friendly, inexpensive, and efficient catalyst are reported. Short reaction times, high yields, scale-up and easy workup are the advantages of these protocols.

2012 JNS All rights reserved

1. Introduction

Tetrasubstituted imidazole core present in many biological systems such as Losartan, Olmesartan, Trifenagrel and Eprosartane [1]. The presence of animidazole ring in natural products and pharmacologically active [2] compounds has instituted a diverse array of synthetic approaches to these heterocycles [3]. However, despite intensive

efforts, only a handful of general methods exist for the construction of tetrasubstitutedimidazoles. Recently, the synthesis of 1, 2, 4, 5-tetrasubstitutedimidazoles has been catalyzed by silica gel or Zeolite HY [4], silica gel/NaHSO₄ [5], molecular iodine [6], K₅CoW₁₂O₄₀.3H₂O [7], heteropolyacids [8], HClO₄-SiO₂ [9], InCl₃.3H₂O [10], ZrCl₄ [11] and BF₃.SiO₂ [12]. Some of the

reported protocols have disadvantages such as harsh reaction conditions, poor yields, prolonged time period, use of hazardous and often expensive acid catalysts. Moreover, the synthesis of these heterocycles has been usually carried out in polar solvents such as ethanol, methanol, acetic acid and DMSO leading to complex isolation and recovery procedures. These processes also generate waste containing catalyst and solvent, which have to be recovered, treated and disposed off. During the course of our studies towards the development of new protocols to the synthesis of organic compounds, we wish to report a simple and efficient method for the synthesis of 1, 2, 4, 5-tetraaryl substituted imidazoles in the presence of nano-silica phosphoric acid. Nano silica phosphoric acid (nano-SPA) [13] as an efficient and reusable catalyst was prepared by reaction of nano-silica chloride [14] with dry phosphoric acid. Nano silica chloride was prepared by reaction of commercial nanosilicagel with thionyl chloride in reflux condition. The particle size of nano-silica gel and nano-SPA was measured by SEM photography (Fig. 1).

The acidic capacity of nano-SPA is 10.32 mmol.g⁻¹ and was determined via titration of 0.2 g of catalyst with standard solution of NaOH.

The FT-IR (ATR) spectra of silica chloride, nano-SPA and H₃PO₄.SiO₂ were shown in Fig 2. In all ATR spectrum, the Si-O-H and Si-O-Si stretching bonds are appeared in 900 until 1100 cm⁻¹. In silica chloride spectrum, the Si-Cl stretching bond is appeared in 700 cm⁻¹. In ATR spectra of nano-SPA and H₃PO₄.SiO₂, the P-O-H, P=O, P-O stretching bonds are appeared 910-1040, 1637 and 2400-2800 cm⁻¹ respectively. According to above data, we suggested one structure for nano-SPA with PO₃H₂ on silica gel (scheme 1).




Fig. 1. SEM photograph of a) nano silica-gel and b) nano-SPA.





Fig. 2. ATR of a) silica chloride, b) nano-SPA and c) H₃PO₄.SiO₂



Scheme 1

The X-ray diffraction (XRD) patterns of nano-SiO₂ and nano-SPA are shown in figure 3. Nano-SiO₂ XRD pattern has a strong peak in 2θ value of 21.8024° and FWHM equal to 1771 and nano- SPA XRD pattern has a strong broad peak in 21.718 and FWHM equal to 2.3616.




Fig. 3. X-ray diffraction (XRD) pattern of a) nano SiO₂ and b) nano-SPA

2. Experimental procedure

2.1. General. The materials were purchased from Merck Company and were used without any additional purification. Products were

characterized by FT-IR, ¹H-NMR and comparison of their physical properties with those reported in the literature. FT-IR (ATR) spectra were run on a Bruker, Equinox 55 spectrometer. A Bruker (DRX-400 Avanes) NMR was used to record the ¹H-NMR spectra. The SEM of nano particles determined with VEGA/TESCAN scanning electron microscope. A BANDELIN Sonopulse HD 3200 ultrasonic apparatus was used for sonication. The X-ray diffraction (XRD) patterns of materials were recorded by employing a Philips XpertMPD diffractometer equipped with a Cu K α anode ($\lambda = 1.54 \text{ \AA}$) in the 2 θ range from 5 to 80°.

2.2. General procedure for the synthesis of 1, 2, 4, 5-tetrasubstitutedimidazoles:


Benzil (4 mmol), amine (4 mmol), aldehyde (4 mmol), ammonium acetate (4 mmol) and nano-SPA (0.08 g) was heated with stirring at 140 °C for 2 h (Table 2). The progress of the reaction was followed by TLC. After completion of the reaction, the mixture was cooled to room temperature. Chloroform was added to the mixture which was filtered to remove the catalyst. After evaporation of the solvent, an oily residue or an impure solid was obtained. By adding ethanol and water to the residue, a milky to yellow solid was obtained. The solid was then crystallized from ethanol. All the products are known and were identified by comparison of their physical and spectral data with those of authentic samples.

3. Results and discussion

In continuation of our investigation on application of solid acids in organic synthesis [15-19], we have used nano-SPA for the synthesis of 1, 2, 4, 5-tetrasubstitutedimidazoles. We have synthesized 1, 2, 4, 5-tetrasubstitutedimidazoles in the presence of SPA under various conditions (table 1). Reaction at

different temperatures and various molar ratios of substrates in the presence of SPA revealed that the best conditions were solvent-free at 140 °C and a ratio of aldehyde (mmol) / amine (mmol) / benzyl (mmol) / ammoniumacetate (mmol) / SPA (g) equal to 1:1:1:0.06. The ultrasonic assisted synthesis of the present reaction was studied by using SPA or nano-SPA in ethyl acetate under reflux conditions (Table 1, entries 15). It was found that under sonication conditions, the reaction time decreased to 40 minutes.

The reusability of the SPA catalyst was also examined. After each run, the product was filtered, the solvent was evaporated. The catalyst residue was washed with CHCl_3 and reused. Treatment with CHCl_3 removes tars more efficiently from the catalyst surface (Table 1, entries 13 and 14). The catalyst was reusable, although a gradual decline in activity was observed. Therefore, various amine and various aldehydes were used as substrates for the synthesis of 1, 2, 4, 5-tetrasubstituted Imidazoles derivatives at 140 °C under solvent-free condition (Scheme 2 and Table 2).



Scheme 2

4. Conclusion

In conclusion, we have demonstrated a simple method for the synthesis of 1, 2, 4, 5-tetrasubstituted Imidazoles using Nano-SPA as a reusable, eco-friendly, inexpensive, and efficient catalyst. Short reaction times, High yield, simplicity of operation and easy work-up are some advantages of this method.

Table 1. Synthesis of 1-benzyl-4, 5-diphenyl-2-*p*-tolyl-1*H*-imidazole under different conditions.

Ent.	Catalyst (g)	Solvent	Cond.	Time (h)/ Yield (%) ^{ref}
1	SPA (0.05)	-	140 °C	2/58
2	SPA (0.06)	-	140 °C	2/94
3	SPA (0.06)	-	r.t.	15/60
4	SPA (0.06)	CH_2Cl_2	140 °C	15/86
5	SPA (0.06)	H_2O	140 °C	15/80
6	SPA (0.06)	DMSO	110 °C	3/90
7	SPA (0.06)	n-Hexan	140 °C	15/60
8	SPA (0.06)	EtOH	140 °C	15/70
9	SPA (0.06)	HOAc	140 °C	15/75
10	SPA (0.06)	MeOH	140 °C	15/86
11	SPA (0.06)	Acetone	140 °C	15/87
12	nano-SPA (0.02)	-	140 °C	2/90
13	nano-SPA (0.02), 2nd run	-	140 °C	2/65
14	nano-SPA (0.02), 3rd run	-	140 °C	2/50
15	Nano-SPA (0.02)	EtOAc	Ultra.	0.6/80
16	Zeolite HY (1 g)	-	MW	0.1/85 ⁴
17	$\text{SiO}_2/\text{NaHSO}_4$ (0.4 g)	-	140 °C	2/92 ⁵
18	$\text{SiO}_2/\text{NaHSO}_4$ (0.4 g)	-	MW	0.25/96 ⁵
19	I_2 (10 mol%)	EtOH	45 °C	0.6/98 ⁶
20	$\text{K}_5\text{CoW}_{12}\text{O}_{40}\cdot 3\text{H}_2\text{O}$ (0.1 mol%)	-	140 °C	2/95 ⁷
21	$\text{K}_5\text{CoW}_{12}\text{O}_{40}\cdot 3\text{H}_2\text{O}$ (0.1 mol%)	-	MW	0.05/95 ⁷
22	$\text{H}_4[\text{PMo}_1\text{VO}_{40}]$ (1 mol%)	EtOH	78 °C	0.25/90 ⁸
23	$\text{SiO}_2/\text{HClO}_4$ (1 mol%)	-	140 °C	0.1/90 ⁹
24	$\text{InCl}_3\cdot 3\text{H}_2\text{O}$	MeOH	r.t.	6/83 ¹⁰
25	ZrCl_4	CH_3CN	r.t.	1/86 ¹¹
26	37% $\text{BF}_3\cdot \text{SiO}_2$	-	140 °C	2/80 ¹²

Acknowledgment

The authors are grateful to University of Kashan for supporting this work by Grant No (159189-90/12/14).

Table 2: Nano-SPA (0.01 g) catalyzed synthesis of 1, 2, 4, 5-tetrasubstituted Imidazoles at 140 °C for 2 hours.

Ent.	Ar	R	Yield (%) heating ^a / sonication ^b	M.P./ °C ^{ref}
1	C ₆ H ₅	C ₆ H ₅	79/70	216 ⁹
2	C ₆ H ₅	C ₆ H ₅ CH ₂	87/82	163 ⁷
3	C ₆ H ₅	Cyclohexyl	56/50	167 ⁴
4	C ₆ H ₅	CH ₃ CH ₂	60/55	115 ⁴
5	4-ClC ₆ H ₄	C ₆ H ₅	78/70	149 ⁶
6	4-ClC ₆ H ₄	C ₆ H ₅ CH ₂	79/65	162 ⁷
7	2-ClC ₆ H ₄	C ₆ H ₅ CH ₂	80/84	140 ⁷
8	4-OHC ₆ H ₄	C ₆ H ₅ CH ₂	62/60	134 ⁷
9	4-CH ₃ C ₆ H ₄	C ₆ H ₅	75/70	185 ⁸
10	4-CH ₃ C ₆ H ₄	C ₆ H ₅ CH ₂	90/80	165 ⁴
11	4-CH ₃ C ₆ H ₄	Cyclohexyl	70/60	162 ⁴
12	3-OMeC ₆ H ₄	C ₆ H ₅ CH ₂	90/80	128 ⁹
13	2-NO ₂ C ₆ H ₄	C ₆ H ₅ CH ₂	89/80	152 ⁹
14	(CH ₃) ₂ CH	C ₆ H ₅ CH ₂	55/50	129 ⁹

^aMolar ratio of benzil:aldehyde :amine :ammonium acetate: is 1:1: 1:1: heated in 140 °C for 120 minutes.

^b Molar ratio of benzil:aldehyde :amine :ammonium acetate: is 1:1: 1:1: in ethyl acetate was sonicated under power 100 for 45 minutes.

References

- [1] S.E. Wolkenberg, D.D. Wisnoski, W.H. Leister, Y. Wang, Z. Zhao and C.W. Lindsley, *Org. Lett.* 6 (2004) 1453-1456.
- [2] L. Nagarapu, A. Satyendra, B. Rajashekar, K. Srinivas, P. Rupa rani, K. Radhika and G. Subhashini, *Bioorg. Med. Chem. Lett.* 18 (2008) 1167-1171.
- [3] (a) J. Sisko, *J. Org. Chem.* 63 (1998) 4529-4531 (b) S.C. Shilcrat, M.K. Moukhallati, J.M.D. Fortunak, L.N. Pridgen, *J. Org. Chem.* 62 (1997) 8449-8454.
- [4] S. Balalaie and A. Arabian. *Green Chem.* 2 (2000) 274-276.
- [5] A.R. Karimi, Z. Alimohammadi, J. Azizian, A.A. Mohammadi and M.R. Mohammadzadeh, *Catal. Commun.* 7 (2006) 728-732.
- [6] M. Kidwai, P. Mothsra V. Bansal, R. K. Somvanshi, A. S. Ethayathulla; S. Dey, T. P Singh., *J. Mol. Catal. A: Chem.*, 265 (2006) 177-182.
- [7] L. Nagarapu, S. Apuri and S. Kantevari. *J. Mol. Catal. A: Chem.*, 266 (2006) 104-108.
- [8] M.M. Heravi; F. Derikvand, F.F. Bamoharram., *J. Mol. Catal. A: Chem.*, 263 (2007) 112-114.
- [9] S. Kantevari, S.V.N. Vuppapapati, D.O. Biradar and L. Nagarapu. *J. Mol. Catal. A: Chem.*, 266 (2006) 109-113.
- [10] S.D. Sharma, P. Hazarika and D. Konwar. *Tetrahedron Lett.*, 49 (2008) 2216-2220.
- [11] G. V. M. Sharma, Y. Jyothi, and P. S. Lakshmi *Synth. Commun.* 36 (2006) 2991-3000.
- [12] J. Sadeghi, B. F. Mirjalili, M. M. Hashemi, *Tetrahedron. Lett.* 49 (2008) 2575-2577.
- [13] M.A. Zolfigol, F. Shirini, K. Zamani, E. Ghofrani and S. Ebrahimi. *Phosphorus Sulfur Silicon Relat. Elem.*, 179 (2004) 2177-2182.
- [14] M. A. Zolfigol, F. Shirini, A. GhorbaniChoghamarani, AzamShiri, H. Keypour, S. Salehzadeh; *Asian J. Chem.* 13 (2001) 849-853.
- [15] B.F. Mirjalili, A. Bamoniri, L. Zamani, *Scientia Iranica.* 19 (2012) 565-568.
- [16] B.F. Mirjalili, A. Bamoniri, A. Akbari, *Tetrahedron Lett.* 49 (2008) 6454-6456.
- [17] B.F. Mirjalili, A. Bamoniri, A. Akbari, and N. Taghavinia, *J. Iran Chem. Soc.* 8 (2011) S129-S134.
- [18] B.F. Mirjalili, A. Bamoniri, A. Akbari, *J. Iran Chem. Soc.* 8 (2011) S135-S140.
- [19] B.F. Mirjalili, A. Bamoniri, A. Akbari, *Chin. Chem. Lett.* 22 (2011) 45-48.