

RESEARCH PAPER

SnCl₄/Nano-Sawdust as an Efficient Bio-based Catalyst for the Synthesis of 2-Substituted Benzimidazoles and Benzothiazoles

Bi Bi Fatemeh Mirjalili ^{1*}, Abdolhamid Bamoniri ², Sedighe Nazemian ³ and Reza Zare Reshquiyea ¹

¹ Department of Chemistry, College of Science, Yazd University, Yazd, Iran

² Department of Organic Chemistry, Faculty of Chemistry, University of Kashan, Kashan, Iran

³ Farhangian University, Yazd, Iran

ARTICLE INFO

Article History:

Received 04 October 2018

Accepted 13 December 2018

Published 01 January 2019

Keywords:

Benzimidazoles

Benzothiazoles

Carbohydrate-based catalyst

SnCl₄/nano-sawdust

Nano-sawdust

ABSTRACT

SnCl₄/nano-sawdust was prepared as a carbohydrate-based catalyst containing of tin bearing cellulose units. The catalyst was characterized by X-ray diffraction (XRD), fourier transform infrared spectroscopy (FT-IR), field emission scanning electron microscopy (FESEM) and energy dispersive X-ray spectroscopy (EDS). The catalyst was applied successfully as a readily available, inexpensive, biodegradable and environmentally benign heterogeneous bio-based solid acid for the one pot synthesis of 2-substituted benzimidazoles and benzothiazoles.

How to cite this article

Mirjalili B, Bamoniri A, Nazemian S, Zare Reshquiyea R. SnCl₄/Nano-Sawdust as an Efficient Bio-based Catalyst for the Synthesis of 2-Substituted Benzimidazoles and Benzothiazoles. *J Nanostruct*, 2019; 9(1):183-189. DOI: 10.22052/JNS.2019.01.020

INTRODUCTION

Benzimidazoles and benzothiazoles are among the most important N-containing five-membered heterocyclic compounds. They exhibit a wide range of biological activities [1] and have also been applied in dyes [2], liquid crystals [3], chemosensing [4], radioactive amyloid imaging agents [5], ligands for asymmetric transformations [6] and corrosion science [7]. Due to the wide range application of these compounds, their synthesis is an area of interest in synthetic chemistry. Therefore, a number of synthetic protocols have been developed for the synthesis of these class compounds [8-10]. The most common method for their synthesis is condensation of 1,2-phenylenediamine [11,12] or 2-aminothiophenols [13,14] with aldehydes in the presence of acid catalyst (Fig. 1).

Development of new solid acids with numerous advantages such as cost-effectiveness,

environmentally benign, easy workup and good stability for one-pot synthesis of these heterocyclic compounds is still required. In this regard, our aim is developing cheap and bio-based solid acid catalysts for these condensation.

Sawdust is a natural, cheap, renewable and readily available source of cellulose with OH groups. In this work, we have investigated the synthesis of bio-based catalyst by bonding Lewis acids to OH groups of D-glucose units in sawdust. Pectin, lignin, proteins, minerals and tannin in sawdust which caused leaching in organic mediums must be removed. For this purpose, the pine sawdust was treated with NaOH, NaOCl, and H₂O₂, respectively. For preparation of nano-sawdust, the sawdust has been treated with concentrated H₂SO₄ for partial hydrolysis of its cellulose [15]. Then, in continuation of our previous efforts about supported catalysts [15-20] and BF₃/nano-sawdust

* Corresponding Author Email: fmirjalili@yazd.ac.ir




Fig. 1. The most common method for synthesis of 2-substituted benzimidazoles (Eq. 1) and 2-substituted benzothiazoles (Eq. 2)

[21], the nano-sawdust was used to synthesis of $\text{SnCl}_4/\text{nano-sawdust}$ as a new, bio-based and green catalyst.

We have examined its catalytic behaviour for the synthesis of benzimidazoles and benzothiazoles under various conditions.

MATERIALS AND METHODS

All chemicals and solvents were buy from the Merck and Fluka chemical companies in high purity. FT-IR spectra were obtained on an attenuated total reflectance fourier transform infrared (ATR-FTIR) spectrophotometer (Bruker, Eqinox 55). ¹H NMR and ¹³C NMR spectra were recorded at 400 MHz and 100 MHz, respectively, on a Bruker DXR-400 spectrometer. Melting points were determined with a Buchi melting point B-540 B.V.CHI apparatus. X-ray diffraction (XRD) pattern was obtained in the 2θ range from 10 to 80° by a Philips Xpert MPD diffractometer equipped with a Cu Kα anode (k = 1.54 Å). Field emission scanning electron microscopy (FESEM) image was obtained on a Mira 3-XMU. Quantitative elemental information (EDS) of $\text{SnCl}_4/\text{nano-sawdust}$ was obtained by SEM/EDS instrument, Phenom pro X.

Preparation of nano-sawdust

4 g of pine sawdust was treated with a solution of 17.5% w/v sodium hydroxide and refluxed for 12 hours, filtered and washed with water. The resulted fibres was then bleached with 100 ml of 1:1 diluted 5% w/v sodium hypochlorite and refluxed for 8 hours, filtered and washed with water. Then, the fibres were treated with 10 ml of 20% v/v hydrogen peroxide at 50 °C for 2 hours, filtered and washed with water. A mixture of the obtained fibres and sulphuric acid (65% H₂SO₄ with fibre to liquor ratio of 1:20) was mixed strongly for 2 hours at 60 °C. The resulted mixture was cooled to room temperature and diluted by distilled water. Then, the suspension was repeatedly centrifuged

at 12000 rpm for 8 minutes using a refrigerated centrifuge (Appendorf Centrifuge 5417R). After each run, the nano-sawdust, as white powder, was washed with distilled water and centrifuged repeatedly till neutralization of the product.

Preparation of $\text{SnCl}_4/\text{nano-sawdust}$

In a ventilated room, by dropping funnel, SnCl_4 (1 mL) was charged drop wise in to a suspension of 1 g nano-sawdust and 10 mL of chloroform. The mixture was stirred for one hour at room temperature. The resulted mixture was filtered and the obtained white solid was washed with chloroform and dried at room temperature.

Typical procedure for synthesis of 2-arylbenzimidazoles or 2-arylbenzothiazoles

To a mixture of 1,2-phenylenediamine or 2-aminobenzothiazole (1mmol) and aldehyde (1.1mmol) in ethanol (10 ml), $\text{SnCl}_4/\text{nano-sawdust}$ (0.08 g) was added. The reaction mixture was refluxed and the progress of the reaction was monitored by TLC (n-hexane/EtOAc, 7:3). After completion of reaction, the reaction mixture was filtered and washed with acetone. By addition of water to the concentrated filtrated, the solid product was appeared. The product was re-crystallized in ethanol to obtain pure product as a white solid.

RESULTS AND DISCUSSION

For investigation of catalyst structure, we have studied the FT-IR (ATR) spectra of pine sawdust and $\text{SnCl}_4/\text{nano-sawdust}$ (Fig. 2). In sawdust FT-IR spectrum, three strong bands at 3323, 2889 and 1025 cm⁻¹ were observed. In $\text{SnCl}_4/\text{nano-sawdust}$, in addition to the sawdust FTIR bands, one band also appeared at 1108 which verify the C-O-Sn group on the catalyst. The our proposed structure for $\text{SnCl}_4/\text{nano-sawdust}$ is similar to the reported structure for cellulose triphosphate gels

(Fig. 3) [22].

The FESEM image of nano-sawdust and SnCl₄/nano-sawdust are shown in Fig. 4. According to the FESEM data, the particle size of catalyst is below 50 nm. Quantitative elemental information (EDS) of SnCl₄/nano-sawdust was measured by SEM/EDS instrument (Fig. 5). According to this data, the weight percentage of Sn, Cl and C are 45.4, 41.0 and 13.5 respectively.

The X-ray diffraction (XRD) pattern of SnCl₄/nano-sawdust is shown in Fig. 6. The values of 2θ and FWHM are shown in table 1. According to XRD pattern, the three signals in 2θ equal to 15.13, 16.74 and 22.93 with FWHM equal to 0.472, 0.944, and 0.472 respectively, show the existence of sawdust. Other signals in 2θ equal to 20.55, 28.56, 34.63 and 41.84 prove the bonding of Sn to

sawdust backbone.

In this study, we have investigated the catalytic efficiency of SnCl₄/nano-sawdust for the synthesis of 2-substituted benzimidazoles and benzothiazoles via reaction of 1,2-phenylenediamine or 2-aminothiophenol with aromatic aldehydes. Synthesis of 2-phenyl-1H-benzimidazole (I) was examined under various condition for finding suitable condition for this preparation. As shown in table 2, entry 6, the most yield of reaction in condensation of benzaldehyde (1.1 mmol) and 1, 2-phenylenediamine (1 mmol) was achieved in the presence of 0.08 g SnCl₄/nano-sawdust and ethanol after 25 minutes reflux. Reusability of catalyst was investigated for three cycles (Table 2, entries 13–15). For this purpose after each run the reaction mixture was diluted with acetone and




Fig. 2. FT-IR (ATR) spectra of (a) pine sawdust and (b) SnCl₄/nano-sawdust.




Fig. 3. Proposed structure for (a) SnCl₄/nano-sawdust and (b) cellulose triphosphate.




Fig. 4. The FESEM image of (a) nano-sawdust and (b) SnCl₄/nano-sawdust.




Fig. 5. EDS analysis diagram of SnCl₄/nano-sawdust.





Fig. 6. XRD pattern of SnCl₄/nano-sawdust

Table 1. *SnCl₄/nano-sawdust* reflexes in XRD diffract gram

No	1	2	3	4	5	6	7
Pos. [°2θ]	15.130	16.748	20.553	22.935	28.563	34.634	41.846
FWHM [°2θ]	0.472	0.944	0.472	0.472	0.708	0.295	1.728

Table 2. Optimization of reaction conditions for the synthesis of 2-phenyl-1*H*-benzimidazole^a




Entry	Cat (g)	Solvent	Temp (°C)/ Time (min)	Yield ^b (%)
1	-	EtOH	RT/60	trace
2	<i>SnCl₄/nano-sawdust</i> (0.02)	EtOH	RT/60	25
3	<i>SnCl₄/nano-sawdust</i> (0.02)	EtOH	Reflux/60	48
4	<i>SnCl₄/nano-sawdust</i> (0.05)	EtOH	Reflux/60	65
5	<i>SnCl₄/nano-sawdust</i> (0.08)	EtOH	Reflux/60	92
6	<i>SnCl₄/nano-sawdust</i> (0.08)	EtOH	Reflux/25	92
7	<i>SnCl₄/nano-sawdust</i> (0.08)	n-Hexan	Reflux/25	25
8	<i>SnCl₄/nano-sawdust</i> (0.08)	MeOH	Reflux/25	80
9	<i>SnCl₄/nano-sawdust</i> (0.08)	THF	Reflux/25	33
10	<i>SnCl₄/nano-sawdust</i> (0.08)	EtOAc	Reflux/25	40
11	<i>SnCl₄/nano-sawdust</i> (0.08)	DMF	80/25	65
12	<i>SnCl₄/nano-sawdust</i> (0.08)	S. F	90/25	N. R
13	<i>SnCl₄/nano-sawdust</i> (0.08), 2 nd run	EtOH	Reflux/25	88
14	<i>SnCl₄/nano-sawdust</i> (0.08), 3 rd run	EtOH	Reflux/25	84
15	<i>SnCl₄/nano-sawdust</i> (0.08), 4 th run	EtOH	Reflux/25	82

^aThe mmol amount of benzaldehyde and 1,2-phenylenediamine is 1:1:1

^b Isolated yield after re-crystallized in ethanol.

Table 3. Optimization of reaction conditions for the synthesis of 2-phenylbenzothiazole^a




Entry	Solvent	Temp (°C)	Time (min)	Yield ^b
1	EtOH	Reflux	25	65%
2	EtOH	Reflux	60	75%
3	EtOH	Reflux	90	75%
4	MeOH	Reflux	90	68%
5	THF	Reflux	90	35%
6	DMF	90	90	79%
7	S. F	90	90	75%
8	S. F	110	90	94%

^aThe mmol amount of benzaldehyde and 2-aminothiophenol is 1:1:1

^b Isolated yield.

Table 4. SnCl₄/nano-sawdust catalyzed synthesis of 2-arylbenzimidazoles and 2-arylbenzothiazoles



Condition X=NH₂ Ethanol, Reflux
X=SH Solvent free, 110 °C

Ent.	X	Y	Ar	Time (min)	Yield ^a	m.p. (°C)	Ref.
1	NH	H	C ₆ H ₅	25	92	292–294 ⁹	[9]
2	NH	H	4-ClC ₆ H ₄	25	88	289–291 ⁹	[9]
3	NH	NO ₂	2, 4-Cl ₂ C ₆ H ₃	25	90	214–215 ¹⁷	[23]
4	NH	H	4-MeC ₆ H ₄	30	90	275–277 ⁹	[9]
5	NH	H	4-NO ₂ C ₆ H ₄	35	85	259–260 ⁹	[9]
6	NH	H	4-OHC ₆ H ₄	30	87	240–242 ¹⁷	[23]
7	NH	H	2,3-(OH) ₂ C ₆ H ₃	40	86	201–202 ¹⁷	[23]
8	NH	H	CH ₂ CH ₂ Ph	30	88	246–247 ¹⁷	[23]
9	NH	CH ₃	4-NO ₂ C ₆ H ₄	25	92	234–236 ¹⁷	[23]
10	NH	NO ₂	4-NO ₂ C ₆ H ₄	25	88	255–256 ¹⁷	[23]
11	NH	NO ₂	2,3-(OH) ₂ C ₆ H ₃	35	86	206–207 ¹⁷	[23]
12	NH	H	4-BrC ₆ H ₄	27	89	250–252 ¹⁷	[23]
13	S	H	C ₆ H ₅	90	94	98–100 ¹⁰	[10]
14	S	H	4-ClC ₆ H ₄	90	95	110–112 ¹⁰	[10]
15	S	H	2, 4-Cl ₂ C ₆ H ₃	90	92	150–152 ¹³	[13]
16	S	H	4-NO ₂ C ₆ H ₄	110	85	226–228 ¹⁴	[14]
17	S	H	3-NO ₂ C ₆ H ₄	110	87	180–182 ¹³	[13]
18	S	H	2-OH C ₆ H ₄	100	92	130–131 ¹⁴	[14]
19	S	H	2, 3-(OH) ₂ C ₆ H ₃	110	85	186–188 ¹³	[13]
20	S	H	3, 4-(OH) ₂ C ₆ H ₃	110	82	190–192 ¹⁴	[14]
21	S	H	4-((CH ₃) ₂ CH) C ₆ H ₄	90	94	115–117 ¹⁴	[14]
22	S	H	4-N(CH ₃) ₂ C ₆ H ₄	120	83	173–175 ¹⁸	[24]
23	S	H	H	120	89	80–81 ¹⁸	[24]

^aIsolated yield

subsequently centrifuged to get the catalyst. The obtained catalyst was then washed with acetone followed by drying in oven at 100 °C for 4 h. It was found that the reactivity of the recovered catalyst decreases slightly for the next run (approx 4%).

According to above mentioned protocol, the best condition for synthesis of 2-phenylbenzothiazole was achieved using 0.08 g catalyst under solvent free condition at 110 °C after 90 min.

We have applied the previous method for the modification of 2-phenylbenzothiazole synthesis. We have found that using 0.08 g SnCl₄/nano-sawdust under solvent free condition at 110 °C and 90 min is the best condition (Table 3, Entry 8).

Finally, The above mentioned optimized conditions were used for synthesis of 2-aryl-1H-benzimidazoles and 2-arylbenzothiazoles (Table 4).

CONCLUSION

In summary, SnCl₄/nano-sawdust as a new carbohydrate based solid acid catalyst was prepared and characterized. This catalyst was successfully applied for synthesis of various 2-arylbenzimidazoles and 2-arylbenzothiazoles

via condensation of 1,2-phenyldiamine or 2-aminothiophenol with aldehydes. This protocol has many advantages including high conversions, eco-friendly, low-cost and easy workup which makes this method more attractive.

ACKNOWLEDGMENTS

The Research Council of Yazd University is gratefully acknowledged for financial support of this study.

CONFLICT OF INTEREST

The authors declare that there are no conflicts of interest regarding the publication of this manuscript.

REFERENCES

- Mavrova AT, Anichina KK, Vuchev DI, Tsenov JA, Denkova PS, Kondeva MS, et al. Anthelmintic activity of some newly synthesized 5(6)-(un)substituted-1H-benzimidazol-2-ylthioacetyl piperazine derivatives. *European Journal of Medicinal Chemistry*. 2006;41(12):1412–20.
- Asensio JA, Gómez-Romero P. Recent Developments on Proton Conduc-ting Poly(2,5-benzimidazole) (ABPBI) Membranes for High Temperature Poly-mer Electrolyte

- Membrane Fuel Cells. *Fuel Cells*. 2005;5(3):336-43.
- Lion CJ, Matthews CS, Wells G, Bradshaw TD, Stevens MFG, Westwell AD. Antitumour properties of fluorinated benzothiazole-substituted hydroxycyclohexa-2,5-dienones ('quinols'). *Bioorganic & Medicinal Chemistry Letters*. 2006;16(19):5005-8.
 - Singh N, Jang DO. Benzimidazole-Based Tripodal Receptor: Highly Selective Fluorescent Chemosensor for Iodide in Aqueous Solution. *Organic Letters*. 2007;9(10):1991-4.
 - Synthesis of novel benzothiazole compounds with an extended conjugated system. *Arkivoc*. 2007;2007(16):113.
 - Figge A, Altenbach HJ, Brauer DJ, Tielmann P. Synthesis and resolution of 2-(2-diphenylphosphinyl-naphthalen-1-yl)-1-isopropyl-1H-benzimidazole; a new atropisomeric P,N-chelating ligand for asymmetric catalysis. *Tetrahedron: Asymmetry*. 2002;13(2):137-44.
 - Chaudhuri P, Ganguly B, Bhattacharya S. An Experimental and Computational Analysis on the Differential Role of the Positional Isomers of Symmetric Bis-2-(pyridyl)-1H-benzimidazoles as DNA Binding Agents. *The Journal of Organic Chemistry*. 2007;72(6):1912-23.
 - Takeuchi K, Bastian JA, Gifford-Moore DS, Harper RW, Miller SC, Mullaney JT, et al. 1,2-disubstituted indole, azaindole and benzimidazole derivatives possessing amine moiety: a novel series of thrombin inhibitors. *Bioorganic & Medicinal Chemistry Letters*. 2000;10(20):2347-51.
 - Kim Y, Kumar MR, Park N, Heo Y, Lee S. Copper-Catalyzed, One-Pot, Three-Component Synthesis of Benzimidazoles by Condensation and C–N Bond Formation. *The Journal of Organic Chemistry*. 2011;76(23):9577-83.
 - Yang Z, Chen X, Wang S, Liu J, Xie K, Wang A, et al. Synthesis of 2-Aryl Benzothiazoles via K₂S₂O₈-mediated Oxidative Condensation of Benzothiazoles with Aryl Aldehydes. *The Journal of Organic Chemistry*. 2012;77(16):7086-91.
 - Bahrami K, Khodaei MM, Naali F. Mild and Highly Efficient Method for the Synthesis of 2-Arylbenzimidazoles and 2-Arylbenzothiazoles. *The Journal of Organic Chemistry*. 2008;73(17):6835-7.
 - Inamdar SM, More VK, Mandal SK. CuO nano-particles supported on silica, a new catalyst for facile synthesis of benzimidazoles, benzothiazoles and benzoxazoles. *Tetrahedron Letters*. 2013;54(6):579-83.
 - Chandrachood P S, Garud D R, Gadakari T V, Torane R C, Deshpande N R, Kashalkar R V. A Cobalt Nitrate/Hydrogen Peroxide System as an Efficient Reagent for the Synthesis of 2-Aryl Benzimidazoles and Benzothiazoles. *Acta Chim. Slov*. 2011; 58(2): 367-371.
 - Bandyopadhyay P, Sathe M, Prasad GK, Sharma P, Kaushik MP. Mesoporous mixed metal oxide nanocrystals: Efficient and recyclable heterogeneous catalysts for the synthesis of 1,2-disubstituted benzimidazoles and 2-substituted benzothiazoles. *Journal of Molecular Catalysis A: Chemical*. 2011;341(1-2):77-82.
 - Salavati-Niasari M, Elzami MR, Mansournia MR, Hydarzadeh S. Alumina-supported vanadyl complexes as catalysts for the CH bond activation of cyclohexene with tert-butylhydroperoxide. *Journal of Molecular Catalysis A: Chemical*. 2004;221(1-2):169-75.
 - Salavati-Niasari M, Hydarzadeh S, Amiri A, Salavati S. Manganese(III) bis(2-hydroxyaniil)acetylacetonato complex as effective catalyst for acylation of alcohols, amines and phenols with acetic anhydride. *Journal of Molecular Catalysis A: Chemical*. 2005;231(1-2):191-5.
 - Salavati-Niasari M, Hasanalian J, Najafian H. Alumina-supported FeCl₃, MnCl₂, CoCl₂, NiCl₂, CuCl₂, and ZnCl₂ as catalysts for the benzylation of benzene by benzyl chloride. *Journal of Molecular Catalysis A: Chemical*. 2004;209(1-2):209-14.
 - Azad S, Mirjalili BBF. Nano-TiCl₂/cellulose: an eco-friendly bio-based catalyst for one-pot synthesis of 4H-pyrimido[2,1-b]benzothiazole derivatives. *Research on Chemical Intermediates*. 2016;43(3):1723-34.
 - Mirjalili BBF, Bamoniri A, Azad S. Synthesis of 2,3-dihydroquinazolin-4(1H)-ones catalyzed by nano-Fe₃O₄/TiCl₂/cellulose as a bio-based magnetic catalyst. *Journal of the Iranian Chemical Society*. 2016;14(1):47-55.
 - Azad S, Fatameh Mirjalili BB. Fe₃O₄@nano-cellulose/TiCl: a bio-based and magnetically recoverable nano-catalyst for the synthesis of pyrimido[2,1-b]benzothiazole derivatives. *RSC Advances*. 2016;6(99):96928-34.
 - Mirjalili BBF, Zare Reshquiyea R. BF₃/nano-sawdust as a green, biodegradable and inexpensive catalyst for the synthesis of highly substituted dihydro-2-oxopyrroles. *RSC Advances*. 2015;5(20):15566-71.
 - Granja PL, Pouységu L, Pétraud M, De Jéso B, Baquey C, Barbosa MA. Cellulose phosphates as biomaterials. I. Synthesis and characterization of highly phosphorylated cellulose gels. *Journal of Applied Polymer Science*. 2001;82(13):3341-53.
 - Mirjalili BF, Bamoniri A, Rahimi Kazerouni M. One-Pot Synthesis of Benzimidazoles in Water in the Presence of SiO₂-OPO₃H. *Chemistry of Heterocyclic Compounds*. 2014;50(1):35-40.
 - Mirjalili BF, Bamoniri A, Akbari A. One-pot synthesis of 3,4-Dihydropyrimidin-2(1H)-ones (thiones) promoted by nano-BF₃.SiO₂. *Journal of the Iranian Chemical Society*. 2011;8(S1):S135-S40.