

## On multiplicative Zagreb indices of graphs

A. IRANMANESH,<sup>1,\*</sup> M. A. HOSSEINZADEH<sup>1</sup> AND I. GUTMAN<sup>2</sup>

<sup>1</sup>Department of Mathematics, TarbiatModares University, P. O. Box 14115-137, Tehran, Iran

<sup>2</sup>Faculty of Science, University of Kragujevac, P. O. Box 60, 34000 Kragujevac, Serbia

(Received August 1, 2012)

### ABSTRACT

Todeschini et al. have recently suggested to consider multiplicative variants of additive graph invariants, which applied to the Zagreb indices would lead to the multiplicative Zagreb indices of a graph  $G$ , denoted by  $\Pi_1(G)$  and  $\Pi_2(G)$ , under the name first and second multiplicative Zagreb index, respectively. These are define as  $\Pi_1(G) = \prod_{v \in V(G)} d_G(v)^2$  and

$\Pi_2(G) = \prod_{uv \in E(G)} d_G(v)d_G(v)$ , where  $d_G(v)$  is the degree of the vertex  $v$ . In this paper we compute these indices for link and splice of graphs. In continuation, with use these graph operations, we compute the first and the second multiplicative Zagreb indices for a class of dendrimers.

**Keywords:** Multiplicative Zagreb indices, splice, link, chain graphs, dendrime

### 1. INTRODUCTION

In this article, we are concerned with simple graphs, that is finite and undirected graphs without loops and multiple edges. Let  $G$  be such a graph and  $V(G)$  and  $E(G)$  be its vertex set and edge set, respectively. An edge of  $G$ , connecting the vertices  $u$  and  $v$  will be denoted by  $uv$ . The degree  $d_G(v)$  of a vertex  $v \in V(G)$  is the number of vertices of  $G$  adjacent to  $v$ .

A molecular graph is a simple graph, such that its vertices correspond to the atoms and the edges to the bonds. Note that hydrogen atoms are often omitted. Molecular descriptors play a significant role in chemistry, pharmacology, etc. Among them, topological indices have a prominent place [10]. The vertex-degree-based graph invariants

\*Corresponding author. (Email: iranmana@modares.ac.ir)

$M_1(G) = \sum_{v \in V(G)} d_G(v)^2$  and  $M_2(G) = \sum_{uv \in E(G)} d_G(u)d_G(v)$  are known under the name first and second Zagreb index, respectively. These have been conceived in the 1970s and found considerable applications in chemistry [3,8,13]. The Zagreb indices were subject to a large number of mathematical studies, of which we mention only a few newest [4,7].

Todeschini et al. [9, 11] have recently proposed to consider multiplicative variants of additive graph invariants, which applied to the Zagreb indices would lead to

$$\Pi_1 = \Pi_1(G) = \prod_{v \in V(G)} d_G(v)^2$$

and

$$\Pi_2 = \Pi_2(G) = \prod_{uv \in E(G)} d_G(u)d_G(v).$$

One can easily see that

$$\Pi_2(G) = \prod_{u \in V(G)} d_G(u)^{d_G(u)}. \quad (1)$$

Our notation is standard and mainly taken from standard books of chemical graph theory [10,12]. For background materials, see reference [2,5,6].

## 2. MAIN RESULTS AND DISCUSSION

The goal of this section is obtaining formulas for some graph operations of multiplicative Zagreb indices. In the next section we will applied these graph operations to get some applications in chemistry.

### SPLICE AND LINK

Suppose that  $G$  and  $H$  are graphs with disjoint vertex sets. Following Došlić [1], for given vertices  $u \in V(G)$  and  $v \in V(H)$  a splice of  $G$  and  $H$  by vertices  $u$  and  $v$ , denoted by  $(G \bullet H)(u, v)$ , is obtained by identifying the vertices  $u$  and  $v$  in the union of  $G$  and  $H$ . Similarly, a link of  $G$  and  $H$  by vertices  $u$  and  $v$  is defined as the graph  $(G \square H)(u, v)$  obtained by joining  $u$  and  $v$  by an edge in the union of these graphs.

Let  $G_1$  and  $G_2$  be graphs as specified above. Denote the vertex obtained by identifying  $v_1$  and  $v_2$  by  $v_{12}$ . Then directly from the definition of their splice and link we obtain:

**Lemma 1.**

$$(i) d_{(G_1 \bullet G_2)(v_1, v_2)}(u) = \begin{cases} d_{G_i}(u) & u \in V(G_i) \text{ and } u \neq v_i \\ d_{G_1}(v_1) + d_{G_2}(v_2) & u = v_{12} \end{cases},$$

$$(ii) d_{(G_1 \square G_2)(v_1, v_2)}(u) = \begin{cases} d_{G_i}(u) & u \in V(G_i) \text{ (} i=1,2 \text{) and } u \neq v_i \\ d_{G_i}(u) + 1 & u = v_i ; i=1,2 \end{cases}.$$

**Theorem 2.** The first multiplicative Zagreb index of the splice and link of  $G_1$  and  $G_2$  satisfies the relations

$$\Pi_1(G_1 \bullet G_2)(v_1, v_2) = \left( \frac{d_{G_1}(v_1) + d_{G_2}(v_2)}{d_{G_1}(v_1) d_{G_2}(v_2)} \right)^2 \Pi_1(G_1) \Pi_1(G_2)$$

$$\Pi_1(G_1 \square G_2)(v_1, v_2) = \left( \frac{(d_{G_1}(v_1) + 1)(d_{G_2}(v_2) + 1)}{d_{G_1}(v_1) d_{G_2}(v_2)} \right)^2 \Pi_1(G_1) \Pi_1(G_2)$$

**Proof.**

For the splice of  $G_1$  and  $G_2$  we have

$$V(G_1 \bullet G_2) = [V(G_1) \setminus \{v_1\}] \cup [V(G_2) \setminus \{v_2\}] \cup \{v_{12}\}$$

which implies

$$\begin{aligned} \Pi_1(G_1 \bullet G_2)(v_1, v_2) &= \left[ \prod_{x \in V(G_1) \setminus \{v_1\}} d_{G_1}(x)^2 \right] \left[ \prod_{y \in V(G_2) \setminus \{v_2\}} d_{G_2}(y)^2 \right] [d_{G_1 \bullet G_2}(v_{12})^2] \\ &= \left[ \frac{1}{d_{G_1}(v_1)^2} \prod_{x \in V(G_1)} d_{G_1}(x)^2 \right] \left[ \frac{1}{d_{G_2}(v_2)^2} \prod_{y \in V(G_2)} d_{G_2}(y)^2 \right] [d_{G_1}(v_1) + d_{G_2}(v_2)]^2 \\ &= \left[ \frac{1}{d_{G_1}(v_1)^2} \Pi_1(G_1) \right] \left[ \frac{1}{d_{G_2}(v_2)^2} \Pi_1(G_2) \right] [d_{G_1}(v_1) + d_{G_2}(v_2)]^2 \end{aligned}$$

from which the first formula follows. Further, by part (ii) of Lemma 1,

$$\begin{aligned}
\Pi_1(G_1 \square G_2)(v_1, v_2) &= \left[ \prod_{x \in V(G_1) \setminus \{v_1\}} d_{G_1}(x)^2 \right] \left[ \prod_{y \in V(G_2) \setminus \{v_2\}} d_{G_2}(y)^2 \right] [d_{G_1 \square G_2}(v_1)^2 d_{G_1 \square G_2}(v_2)^2] \\
&= \left[ \frac{1}{d_{G_1}(v_1)^2} \prod_{x \in V(G_1)} d_{G_1}(x)^2 \right] \left[ \frac{1}{d_{G_2}(v_2)^2} \prod_{y \in V(G_2)} d_{G_2}(y)^2 \right] [[d_{G_1}(v_1)+1]^2 [d_{G_2}(v_2)+1]^2] \\
&= \left[ \frac{1}{d_{G_1}(v_1)^2} \Pi_1(G_1) \right] \left[ \frac{1}{d_{G_2}(v_2)^2} \Pi_1(G_2) \right] [[d_{G_1}(v_1)+1][d_{G_2}(v_2)+1]]^2
\end{aligned}$$

from which the second formula follows. ■


In a fully analogous manner, bearing in mind Eq. (1), we can prove:

**Theorem 3.** The second multiplicative Zagreb index of the splice and link of  $G_1$  and  $G_2$  satisfies the relations

$$\begin{aligned}
\Pi_2(G_1 \bullet G_2)(v_1, v_2) &= \frac{[d_{G_1}(v_1) + d_{G_2}(v_2)]^{d_{G_1}(v_1) + d_{G_2}(v_2)}}{d_{G_1}(v_1)^{d_{G_1}(v_1)} d_{G_2}(v_2)^{d_{G_2}(v_2)}} \Pi_2(G_1) \Pi_2(G_2) \\
\Pi_2(G_1 \square G_2)(v_1, v_2) &= \frac{(d_{G_1}(v_1) + 1)^{d_{G_1}(v_1) + 1} (d_{G_2}(v_2) + 1)^{d_{G_2}(v_2) + 1}}{d_{G_1}(v_1)^{d_{G_1}(v_1)} d_{G_2}(v_2)^{d_{G_2}(v_2)}} \Pi_2(G_1) \Pi_2(G_2).
\end{aligned}$$

## CHAIN GRAPHS

Let  $G_i$  ( $1 \leq i \leq n$ ) be some graphs and  $v_i \in V(G_i)$ . A chain graph, denoted by  $G = G(G_1, \dots, G_n, v_1, \dots, v_n)$ , is obtained from the union of the graphs  $G_i$ ,  $i = 1, 2, \dots, n$ , by adding the edges  $v_i v_{i+1}$  ( $1 \leq i \leq n-1$ ), see Fig. 1. Then  $|V(G)| = \sum_{i=1}^n |V(G_i)|$  and  $|E(G)| = (n-1) + \sum_{i=1}^n |E(G_i)|$ .



**Fig. 1.** The Chain Graph  $G = G(G_1, \dots, G_n, v_1, \dots, v_n)$ .

One can see that  $G(G_1, G_2, v_1, v_2) \cong (G_1 \square G_2)(v_1, v_2)$ .

It is worth noting that the above-specified class of chain graphs embraces, as special cases, all trees and all unicyclic graphs. In addition, the molecular graphs of many polymers and dendrimers are chain graphs.

**Lemma 4.** Suppose that  $G = G(G_1, G_2, \dots, G_n, v_1, v_2, \dots, v_n)$  is a chain graph. Then:

$$d_G(u) = \begin{cases} d_{G_i}(u) & u \in V(G_i) \text{ and } u \neq v_i \\ d_{G_i}(u) + 1 & u = v_i, \quad i = 1, n \\ d_{G_i}(u) + 2 & u = v_i, \quad 2 \leq i \leq n - 1 \end{cases}.$$

**Theorem 5.** If  $n \geq 3$  and  $v_1, \dots, v_n \neq u_1, \dots, u_k$ , then for  $G = G(G_1, G_2, \dots, G_n, v_1, v_2, \dots, v_n)$  the following holds:

$$\begin{aligned} \Pi_1^{(u_1, \dots, u_k)}(G) &= \prod_{i=1}^n \Pi_1^{(v_i, u_1, \dots, u_k)}(G_i) \prod_{j=2}^{n-1} (d_{G_j}(v_j) + 2)^2 \prod_{k=1, n} (d_{G_k}(v_k) + 1)^2, \\ \Pi_2^{(u_1, \dots, u_k)}(G) &= \prod_{i=1}^n \Pi_2^{(v_i, u_1, \dots, u_k)}(G_i) \prod_{j=2}^{n-1} (d_{G_j}(v_j) + 2)^{d_{G_j}(v_j) + 2} \prod_{k=1, n} (d_{G_k}(v_k) + 1)^{d_{G_k}(v_k) + 1} \end{aligned}$$

where we use the truncated versions of the two multiplicative Zagreb indices, namely,

$$\Pi_1^{(u_1, u_2, \dots, u_k)}(G) = \prod_{\substack{v \in V(G) \\ v \neq u_1, u_2, \dots, u_k}} d_G(v)^2 \text{ and } \Pi_2^{(u_1, u_2, \dots, u_k)}(G) = \prod_{\substack{v \in V(G) \\ v \neq u_1, u_2, \dots, u_k}} d_G(v)^{d_G(v)}.$$

**Proof.**

$$\Pi_1^{(u_1, u_2, \dots, u_k)}(G) = \prod_{\substack{v \in V(G) \\ v \neq u_1, u_2, \dots, u_k}} d_G(v)^2$$

$$\begin{aligned}
&= \prod_{i=1}^n \left( \prod_{\substack{v \in V(G_i) \\ v \neq v_j, u_1, u_2, \dots, u_k}} d_{G_i}(v)^2 \right) (d_{G_1}(v_1)+1)^2 (d_{G_2}(v_2)+2)^2 \cdots (d_{G_{n-1}}(v_{n-1})+2)^2 (d_{G_n}(v_n)+1)^2 \\
&= \prod_{i=1}^n \Pi_1^{(v_i, u_1, \dots, u_k)}(G_i) \prod_{j=2}^{n-1} (d_{G_j}(v_j)+2)^2 \prod_{k=1, n} (d_{G_k}(v_k)+1)^2,
\end{aligned}$$

and similarly

$$\begin{aligned}
\Pi_2^{(u_1, u_2, \dots, u_k)}(G) &= \prod_{\substack{v \in V(G) \\ v \neq u_1, u_2, \dots, u_k}} d_G(v)^{d_G(v)} \\
&= \prod_{i=1}^n \left( \prod_{\substack{v \in V(G_i) \\ v \neq v_j, u_1, u_2, \dots, u_k}} d_{G_i}(v)^{d_{G_i}(v)} \right) \prod_{j=2}^{n-1} (d_{G_j}(v_j)+2)^{d_{G_j}(v_j)+2} \prod_{k=1, n} (d_{G_k}(v_k)+1)^{d_{G_k}(v_k)+1} \\
&= \prod_{i=1}^n \Pi_2^{(v_i, u_1, \dots, u_k)}(G_i) \prod_{j=2}^{n-1} (d_{G_j}(v_j)+2)^{d_{G_j}(v_j)+2} \prod_{k=1, n} (d_{G_k}(v_k)+1)^{d_{G_k}(v_k)+1}.
\end{aligned}$$

### 3. APPLICATION

The goal of this section is computing the truncated Zagreb indices of an infinite class of dendrimers. To do this, we use Theorems 2 and 3. The truncated Zagreb indices for other classes of dendrimers, can be computed similarly. Consider the graph  $G_1$  shown in Fig. 2. It is easy to see that

$$\Pi_1(G_1) = 2^{30}3^8 \text{ and } \Pi_2(G_1) = 2^{30}3^{12},$$


$$\Pi_1^{(v_1)}(G_1) = \Pi_1^{(v_2)}(G_1) = \Pi_1^{(v_3)}(G_1) = \Pi_1^{(v)}(G_1) = 2^{28}3^8$$

and

$$\Pi_2^{(v_1)}(G_1) = \Pi_2^{(v_2)}(G_1) = \Pi_2^{(v_3)}(G_1) = \Pi_2^{(v)}(G_1) = 2^{28}3^{12},$$

and so, for  $1 \leq i, j \leq 3, i \neq j$ ,

$$\Pi_1^{(v, v)}(G_1) = \Pi_1^{(v_i, v_j)}(G_1) = 2^{26}3^8 \text{ and } \Pi_2^{(v, v)}(G_1) = \Pi_2^{(v_i, v_j)}(G_1) = 2^{26}3^{12}.$$



**Fig. 2.**The Graph of Dendrimer  $G_n$  for  $n=1$ .

Consider now the graph  $G_n = (G_{n-1} \square H_1)(v_1, u_1)$ , shown in Fig. 2 (for  $n=1$ ) and Fig. 3, respectively. It is easy to see that  $H_i \cong G_1$  ( $1 \leq i \leq n-1$ ) and

$$\begin{aligned} G_n &= (G_{n-1} \square H_1)(v_1, u_1) \\ G_{n-1} &= (G_{n-2} \square H_2)(v_2, u_2) \\ &\vdots \\ G_{n-i} &= (G_{n-i-1} \square H_{i+1})(v_{i+1}, u_{i+1}) \\ &\vdots \\ G_2 &= (G_1 \square H_{n-1})(v_{n-1}, u_{n-1}). \end{aligned}$$

Then by using Theorem 2, we have the following relations:

$$\begin{aligned} \Pi_1(G_n) &= \Pi_1^{(v_1)}(G_{n-1})\Pi_1^{(u_1)}(H_1) \times 3^4 \\ \Pi_1^{(v_1)}(G_{n-1}) &= \Pi_1^{(v_2)}(G_{n-2})\Pi_1^{(v_1, u_2)}(H_2) \times 3^4 \\ &\vdots \\ \Pi_1^{(v_i)}(G_{n-i}) &= \Pi_1^{(v_{i+1})}(G_{n-i-1})\Pi_1^{(v_i, u_{i+1})}(H_{i+1}) \times 3^4 \\ &\vdots \\ \Pi_1^{(v_{n-2})}(G_2) &= \Pi_1^{(v_{n-1})}(G_1)\Pi_1^{(v_{n-2}, u_{n-1})}(H_{n-1}) \times 3^4. \end{aligned}$$

Multiplication of these relations yields

$$\Pi_1(G_n) = \Pi_1^{(v_{n-1})}(G_1)\Pi_1^{(u_1)}(H_1) \prod_{i=2}^{n-1} \Pi_1^{(v_{i-1}, u_i)}(H_i) \times 3^{4(n-1)},$$

and it is easy to obtain

$$\Pi_1(G_n) = \left(\Pi_1^{(v_1)}(G_1)\right)^2 \left(\Pi_1^{(v_1, v_2)}(G_1)\right)^{n-2} \times 3^{4(n-1)}$$

$$= 2^{26n+4} 3^{12n-4}.$$

In other words we arrived at the following:

**Theorem 6.** Consider the graph  $G_n = (G_{n-1} \square H_1)(v_1, u_1)$  ( $n \geq 2$ ), shown in Fig. 3. Then,

$$\Pi_1(G_n) = 2^{26n+4} 3^{12n-4}.$$

**Corollary 7.** Consider the dendrimer  $D$ , shown in Fig. 4. Then,

$$\Pi_1(D) = 2^{26n+4} 3^{12n-4},$$

Where  $n$  is the number of repetition of the fragment  $G_1$ .

Now for the second truncated Zagreb index, by using Theorem 3, the following relations are hold:

$$\begin{aligned} \Pi_2(G_n) &= \Pi_2^{(v_1)}(G_{n-1}) \Pi_2^{(u_1)}(H_1) \times 3^6 \\ \Pi_2^{(v_1)}(G_{n-1}) &= \Pi_2^{(v_2)}(G_{n-2}) \Pi_2^{(v_1, u_2)}(H_2) \times 3^6 \\ &\vdots \\ \Pi_2^{(v_i)}(G_{n-i}) &= \Pi_2^{(v_{i+1})}(G_{n-i-1}) \Pi_2^{(v_i, u_{i+1})}(H_{i+1}) \times 3^6 \\ &\vdots \\ \Pi_2^{(v_{n-2})}(G_2) &= \Pi_2^{(v_{n-1})}(G_1) \Pi_2^{(v_{n-2}, u_{n-1})}(H_{n-1}) \times 3^6. \end{aligned}$$

So by production of two sides of these relations, we have

$$\Pi_2(G_n) = \Pi_2^{(v_{n-1})}(G_1) \Pi_2^{(u_1)}(H_1) \prod_{i=2}^{n-1} \Pi_2^{(v_{i-1}, u_i)}(H_i) \times 3^{6(n-1)},$$

and it is easy to see that

$$\begin{aligned} \Pi_2(G_n) &= \left( \Pi_2^{(v_1)}(G_1) \right)^2 \left( \Pi_2^{(v_1, v_2)}(G_1) \right)^{n-2} \times 3^{6(n-1)} \\ &= 2^{26n+4} 3^{18n-6}. \end{aligned}$$

**Theorem 8.** Consider the graph  $G_n = (G_{n-1} \square H_1)(v_1, u_1)$  ( $n \geq 2$ ), shown in Fig. 3. Then,

$$\Pi_2(G_n) = 2^{26n+4} 3^{18n-6}.$$

**Corollary 9.** Consider the dendrimer  $D$ , shown in Fig. 4. Then,

$$\Pi_2(D) = 2^{26n+4} 3^{18n-6},$$

Where  $n$  is the number of repetition of the fragment  $G_1$ .






Fig. 3. The graph  $G_n$  and the labeling of its vertices.




Fig. 4: The Graph of the Dendrimer  $D$ .

**REFERENCES**

1. T. Došlić, Splices, links and their degree-weighted Wiener polynomials, *Graph Theory Notes New York*, **48** (2005), 47-55.
2. T. Došlić, M. Ghorbani and M. A. Hosseinzadeh, Eccentric connectivity polynomial of some graph operations, *Util. Math.*, **84** (2011), 297-309.
3. M. Ghorbani and M. A. Hosseinzadeh, A note of Zagreb indices of nanostar dendrimers, *Optoelectron. Adv. Mater.*, **4** (2010), 1877-1880.
4. M. Ghorbani and M. A. Hosseinzadeh, A new version of Zagreb indices, *Filomat*, **26** (2012), 93-100.
5. M. Ghorbani, M. A. Hosseinzadeh and M. V. Diudea, On Omega polynomial of Cartesian product graph, *Util. Math.*, **84** (2011), 165-172.
6. M. Ghorbani, M. A. Hosseinzadeh and I. Gutman, The truncated Randić-type indices, *Kragujevac J. Sci.*, **32** (2010), 47-56.
7. S. Li and M. Zhang, Sharp upper bounds for Zagreb indices of bipartite graphs with given diameter, *Appl. Math. Lett.*, **24** (2011), 131-137.
8. S. Nikolić, G. Kovačević, A. Miličević and N. Trinajstić, The Zagreb indices 30 years after, *Croat. Chem. Acta*, **76** (2003), 113-124.
9. R. Todeschini, D. Ballabio and V. Consonni, Novel molecular descriptors based on functions of new vertex degrees. in: I. Gutman and B. Furtula (Eds.), *Novel Molecular Structure Descriptors - Theory and Applications I*, Univ. Kragujevac, Kragujevac, 2010, pp. 73-100.
10. R. Todeschini and V. Consonni, *Handbook of Molecular Descriptors*, Wiley-VCH, Weinheim, 2000.
11. R. Todeschini and V. Consonni, New local vertex invariants and molecular descriptors based on functions of the vertex degrees, *MATCH Commun. Math. Comput. Chem.*, **64** (2010), 359-372.
12. N. Trinajstić, *Chemical Graph Theory*, CRC Press, Boca Raton, 1992.
13. N. Trinajstić, S. Nikolić, A. Miličević and I. Gutman, On Zagreb indices, *Kem. Ind.*, **59** (2010), 577-589.