

## The Topological Indices of some Dendrimer Graphs

M. R. DARAFSHEH<sup>a</sup>, M. NAMDARI<sup>b</sup> AND S. SHOKROLAHI<sup>b,\*</sup>

<sup>a</sup> School of Mathematics, Statistics and Computer Science, College of Science, University of Tehran, Tehran, Iran

<sup>b</sup> Department of Mathematics, Shahid Chamran University of Ahvaz, Ahvaz, Iran

---

### ARTICLE INFO

---

#### Article History:

Received 6 January 2015

Accepted 11 January 2016

Published online 5 September 2016

Academic Editor: Hassan Yousefi–Azari

---

#### Keywords:

Topological index

Dendrimer

Wiener index

Hyper Wiener index

---

---

### ABSTRACT

---

In this paper, the Wiener and hyper Wiener indices of two kinds of dendrimer graphs are computed. Using the Wiener index formula, the Szeged, Schultz, PI and Gutman indices of these graphs are also determined.

© 2017 University of Kashan Press. All rights reserved

---

## 1. INTRODUCTION

Let  $G = (V, E)$  be a simple connected graph with vertex set  $V$  and edge set  $E$ . A topological index of a simple connected graph  $G$  is a graph invariant which is related to the structure of the graph. The Wiener index is one of the best known topological index of a simple connected graph which is studied in both mathematical and chemical literature and its definition is in terms of distances between arbitrary pairs of vertices, see for example [1, 2, 3, 4]. The Wiener index of  $G$  is denoted by  $W(G)$  and it is defined by:

$$W(G) = \sum_{\{u,v\} \subseteq V(G)} d(u,v) = \frac{1}{2} \sum_{u \in V} d(u),$$

where  $d(u, v)$  is the distance between vertices  $u$  and  $v$  and  $d(u) = \sum_{v \in V} d(u, v)$ .

---

\*Corresponding Author: (Email address: [shokrolahisara@yahoo.com](mailto:shokrolahisara@yahoo.com))

DOI: 10.22052/ijmc.2017.15413

The Szeged index [5, 6] is another invariant of a graph which is based on the distribution of the vertices and introduced by Ivan Gutman and it is the same with the Wiener index in the case that  $G$  is a tree. The set of vertices of graph  $G$  which are closer to  $u$  (resp.  $v$ ) than  $v$  (resp.  $u$ ) is denoted by  $N_u(e|G)$  (resp.  $N_v(e|G)$ ). This index is defined as the summation of  $(n_u(e|G) n_v(e|G))$  where  $n_u(e|G)$  (resp.  $n_v(e|G)$ ), is the number of vertices of graph  $G$  closer to  $u$  (resp.  $v$ ) than  $v$  (resp.  $u$ ), over all edges  $e = uv$  of graph. Now, the Szeged index of  $G$  which is denoted by  $Sz(G)$  is defined as:

$$Sz(G) = \sum_{e=uv \in E} (n_u(e|G) \cdot n_v(e|G)).$$

The Padmaker-Ivan ( $PI$ ) index [7, 8] is another topological index of a simple connected graph that takes into account the distribution of edges so is closely related to Szeged index. The  $PI$  index of  $G$  is defined by

$$PI(G) = \sum_{e=uv \in E} (n_{eu}(e|G) + n_{ev}(e|G)),$$

where  $(n_{eu}(e|G))$  (resp.  $(n_{ev}(e|G))$ ) is the number of edges of the subgraph of  $G$  which has the vertex set  $N_u(e|G)$  (resp.  $N_v(e|G)$ ).

The molecular topological index (Schultz index) was introduced by Schultz and Schultz [9, 10]. In addition to the chemical applications, the Schultz index attracted some attention that in the case of a tree it is related to the Wiener index [11]. It is denoted by  $S(G)$  and defined as follows:

$$S(G) = \sum_{\{u,v\} \subseteq V} (\rho(u) + \rho(v)),$$

where  $\rho(u)$  (resp.  $\rho(v)$ ) is the degree of vertex  $u$  (resp.  $v$ ).


The Gutman index which attracts more attention recently is defined by Klavžar and Gutman in [11, 12]. This index is also known as the Schultz index of the second kind but in this paper the first name is used. Gutman [11] has proved that if  $G$  is a tree then there is a relation between Wiener and Gutman indices of  $G$  that we will mention this in Section 2. The Gutman index of  $G$  is denoted by  $Gut(G)$  and is defined as follows:

$$Gut(G) = \sum_{\{u,v\} \subseteq V} (\rho(u)\rho(v))$$

The hyper-Wiener index is one of the graph invariants, used as a structure descriptor related to physicochemical properties of compounds. This index was introduced by Randić in 1993 as extension of Wiener index [13] and it has come to be known as the hyper-Wiener index by Klein [14]. The hyper-Wiener index of  $G$  is denoted by  $WW(G)$  and is defined as follows:

$$WW(G) = \frac{1}{2} (W(G) + \sum_{\{u,v\} \subseteq V} d^2(u,v)).$$

Here we mainly try to determine the Wiener, hyper Wiener and *PI* indices of two kinds of dendrimer graphs (explained in Section 2), then the Schultz, Szeged and Gutman indices are obtained as results of the relation between the Wiener index with both the Schultz and Gutman indices.



**Figure 1.** The first dendrimer graph  $G_n$ .

## 2. CALCULATING THE WIENER, HYPER-WIENER AND PI INDICES OF THE FIRST DENDRIMER GRAPH $G_n$

Let  $G = (V, E)$  be the graph with vertex set  $V$  and edge set  $E$  as in Figure 1. This graph begins with one vertex  $u_0$  which connects to two other vertices such that each one of these two vertices connects to two other vertices and so on. The vertices which have the same distance from  $u_0$  are located on a branch. Let  $G$  have  $(n + 1)$  branches so there are  $2^i$  vertices in the  $i$ '-th branch ( $0 \leq i \leq n$ ). We denote this graph by  $G_n$ .

**Proposition 2.1.** Let  $G_n = (V, E)$  be the dendrimer graph in Figure 1, then:

$$W(G_n) = 4^{(n+1)} (n - 2) + 2^{(n+1)} (n + 4).$$

**Proof.** From definitions we have:

$$W(G_n) = \sum_{\{u,v\} \subseteq V(G)} d(u, v) = \frac{1}{2} \sum_{u \in V} d(u).$$

This graph has  $n + 1$  branches and there are  $2^i$  vertices in the  $i$ '-th branch, so we denote the vertex set of this branch by  $V_i$ , hence we have:  $V = \bigcup_{i=0}^n V_i$ . Because of the symmetric structure of the graph  $G_n$  (Figure 1), for every vertex  $u$  in the  $n$ 'th branch,  $d(u)$  is constant and doesn't depend on  $u$ . So we choose  $u_i$  as representative of the  $i$ '-th branch ( $0 \leq i \leq n$ ).

$$d(u_n) = \sum_{v \in V_n} d(u_n, v) + \sum_{v \in V - V_n} d(u_n, v). \tag{1}$$

$2^{n-1}$  vertices which are in lower branch of Figure 1, are of the same distance from  $u_n$  and this value equals to:

$$2d(u_n, u_0) = 2n.$$

Also  $2^{n-2}$  vertices are of the same distance from  $u_n$  and this value equals to:

$$2d(u_n, u_1) = 2(n-1).$$

Finally continuing in this way the distance between  $u_n$  to the last vertex in the  $n'$ -th branch is equals to:

$$2d(u_n, u_{n-1}) = 2.$$

So we have:

$$\begin{aligned} \sum_{v \in V_n} d(u_n, v) &= 2^{n-1} \times 2n + 2^{n-2} \times 2(n-1) + \dots + 2^{(1-1)} 2 \\ &= \sum_{v \in V_n} d(u_n, v) = n2^n \times (n-1)2^{(n-1)} + \dots + 1.2 \\ &= \sum_{i=1}^n i2^i = 2(1 + (n-1)2^n). \end{aligned} \tag{2}$$

For computing the second part of the summation in (1), note that because the graph  $G_n$  is a tree, for every vertex  $v \in \bigcup_{i=0}^n V_i$  we have:

$$d(u_n, v) = 1 + d(u_{n-1}, v)$$

$$\sum_{v \in \bigcup_{i=0}^{n-1} V_i} d(u_n, v) - \sum_{v \in \bigcup_{i=0}^{n-1} V_i} d(u_{n-1}, v) = \sum_{i=0}^{n-1} 2^i. \tag{3}$$

Considering (2) and (3):

$$d(u_n) - d(u_{n-1}) = \sum_{i=0}^{n-1} 2^i + 2(1 + (n-1)2^n) = 2n2^n - 2^n + 1.$$

Because,  $d(u_0) = 0$ . Hence :

$$d(u_n) = \sum_{i=1}^n (d(u_i) - d(u_{i-1})) = \sum_{i=1}^n 2i2^i - 2^i + 1 = (2n-3)2^{n+1} + n + 6. \tag{4}$$

By multiplying  $2^n$  in  $d(u_n)$  the distance between vertices in the  $n'$ -th branch is considered twice, so if the Wiener index of  $G_n$  with  $n$  (resp.  $n+1$ ) branch is denoted by  $W(n+1)$  (resp.  $W(n)$ ) we have:

$$\begin{aligned} W(n) - W(n-1) &= 2^n ((2n-3)2^{n+1} + (n+6)) - \sum_{\{u,v\} \subseteq V_n} d(u, v) \\ &= 2^n (2n-3)2^{(n+1)} + 2^n (n+6) - 2^n (1 + (n-1)2^n) \\ &= (3n-5)2^{2n} + (n+5)2^n. \end{aligned}$$

So,

$$W(n) = \sum_{i=1}^n (3k-5)2^{2k} + (k+5)2^k = 4^{(n+1)}(n-2) + 2^{(n+1)}(n+4). \quad \blacksquare$$

**Corollary 2.2.**  $Sz(G_n) = 4^{(n+1)}(n-2) + 2^{(n+1)}(n+4)$ .

Proof. The graph  $G_n$  is a tree, so by [11] the result is obtained. ■

**Corollary 2.3.**  $S(G_n) = 4^{(n+1)}(4n-9) + 2^{(n+1)}(4n+19) - 2$ .

Proof. Because  $G_n$  is a tree by [11] we have:  $S(G_n) = 4W(G_n) - n(n-1)$ , where  $n$  is the number of vertices of  $G_n$ . Now by replacing the closed form of  $W(G_n)$  which was obtained from proposition 2.1, the proof is completed. ■

**Corollary 2.4.**  $Gut(G_n) = 4^{(n+1)}(4n-10) + 2^{(n+1)}(4n+19) + 10$ .

**Proof.** Because  $G_n$  is a tree, by [11] we have,  $Gut(G_n) = 4W(G_n) - (2n-1)(n-1)$  where  $n$  is the number of vertices of  $G_n$  and by proposition 2.1 it is done. ■

**Corollary 2.5.**  $PI(G_n) = (2^{(n+1)} - 3)(2^{(n+1)} - 2)$ .

**Proof.** Because  $G_n$  is a tree so for every edge  $e = uv$  of  $G_n$  we have:

$$n_u(e | G_n) + n_v(e | G_n) = |V| = 2^{n+1} - 1.$$

Subgraphs of  $G_n$  with vertex sets  $N_u(e | G_n)$  and  $N_v(e | G_n)$  both are trees and whose number of edges are  $n_u(e | G_n) - 1$  and  $n_v(e | G_n) - 1$  respectively. Then we have:

$$\begin{aligned} n_{eu}(e | G_n) + n_{ev}(e | G_n) &= n_u(e | G_n) + n_v(e | G_n) - 2 = 2^{n+1} - 3 \\ |E|(2^{(n+1)} - 3) &= (2^{(n+1)} - 2)(2^{(n+1)} - 3) \\ PI(G_n) &= |E|(2^{(n+1)} - 3) = (2^{(n+1)} - 2)(2^{(n+1)} - 3) \end{aligned}$$

**Proposition 2.6.** The hyper-Wiener index of  $G_n$  in Figure 1 is:

$$WW(G_n) = 4^n(4n^2 - 14n + 24) + 2^n(n^2 - 3n - 31) - 1.$$

Proof. By definition we have:

$$WW(G) = \frac{1}{2}(W(G) + \sum_{\{u,v\} \subseteq V} d^2(u,v)). \quad (5)$$

Because of the symmetric structure of the graph  $G_n$  in Figure 1,  $d(u)$  for every vertex  $u$  in the  $n'$ -th branch is constant and doesn't depend on  $u$ , so we choose  $u_i$  as representative of the  $i'$ -th branch ( $0 \leq i \leq 1$ ).

$$d^2(u_n) = \sum_{v \in V} d^2(u_n, v) = \sum_{v \in \bigcup_{i=0}^{n-1} V_i} d^2(u_n, v) + \sum_{v \in V_n} d^2(u_n, v). \quad (6)$$

The graph  $G_n$  is a tree, so, for every vertex,  $v \in \bigcup_{i=0}^{n-1} V_i$ :

$$d(u_n, v) = 1 + d(u_{n-1}, v).$$

Now by (6) we have:

$$\begin{aligned} d^2(u_n) &= \sum_{v \in \bigcup_{i=0}^{n-1} V_i} (d(u_{n-1}, v) + 1)^2 + \sum_{v \in V_n} d^2(u_n, v) \\ &= \sum_{v \in \bigcup_{i=0}^{n-1} V_i} d^2(u_{n-1}, v) + 2 \sum_{v \in \bigcup_{i=0}^{n-1} V_i} d(u_{n-1}, v) + \sum_{v \in V_n} d^2(u_n, v) + 2^n - 1 \\ &= d^2(u_{n-1}) + 2d(u_{n-1}) + \sum_{v \in V_n} d^2(u_n, v) + 2^n - 1. \end{aligned} \quad (7)$$

$2^n$  vertices are in the  $n'$ -th branch and by symmetric structure of the graph  $G_n$  we have :

$$\begin{aligned} \sum_{v \in V_n} d^2(u_n, v) &= 2^{n-1}(2n)^2 + 2^{n-2}(2n-2)^2 + \dots + 2^0(2)^2 \\ &= \sum_{i=1}^n 2^{i+1} i^2 = 2^{n+2}(n^2 - 2n + 3) - 12. \end{aligned} \quad (8)$$

By (4) in the proof of the proposition 2.1, and considering (7), (8):

$$\begin{aligned} d^2(u_n) &= \sum_{i=1}^n d^2(u_i) - d^2(u_{i-1}) \\ &= \sum_{i=1}^n (4i^2 - 4i + 3)2^i + 2i - 3 = 2^{n+1}(4n^2 - 12n + 19) + n^2 - 2n - 38 \end{aligned}$$

Therefore:

$$\begin{aligned} \sum_{\{u,v\} \subseteq V} d^2(u, v) - \sum_{\{u,v\} \subseteq \bigcup_{i=0}^{n-1} V_i} d^2(u, v) &= 2^n d^2(u_n) - \sum_{\{u,v\} \subseteq V_n} d^2(u, v) \\ &= 2^n d^2(u_n) - 2^{n-1} \sum_{v \in V_n} d^2(u_n, v) \\ &= 2^{2n+1}(3n^2 - 10n + 16) + 2n(n^2 - 2n - 32). \end{aligned}$$

Now let,

$$F(i) = \sum_{\{u,v\} \subseteq \bigcup_{j=0}^i V_j} d^2(u, v).$$

So,

$$\begin{aligned} \sum_{\{u,v\} \subseteq V} d^2(u, v) &= \sum_{i=1}^n (F(i) - F(i-1)) = \sum_{i=1}^n 2^{2i+1}(3i^2 - 10i + 16) + 2^i(i^2 - 2i - 32) \\ &= 4^{n+1}.2(n^2 - 4n + 7) + 2^{n+1}(n^2 - 4n - 27) - 2. \end{aligned} \quad (9)$$

Now considering (9) and the formula of  $W(G_n)$  which was computed in proposition 2.1, and replacing those in (5), the proof is done. ■




Figure 2. The second dendrimer graph  $H_n$ .

### 3. CALCULATING THE WIENER, HYPER-WIENER AND PI INDICES OF THE SECOND DENDRIMER GRAPH $H_n$

Let  $G = (V, E)$  be the graph with vertex set  $V$  and edge set  $E$ , that begins with one vertex  $u_0$  in Figure 2 that connects to three vertices which form the first branch and each one of these three vertices connects to two other vertices in second branch and so on. It means that any vertex but  $u_0$  in the  $i$ '-th branch joins to the two vertices in the  $(i+1)$ '-th branch, so the vertices which have the same distance from  $u_0$  are located on one branch. Let  $G$  have  $n+1$  branches therefore, there are  $3 \times 2^{i-1}$  vertices in the  $i$ '-th branch ( $0 < i \leq n$ ). The graph  $G$  is another kind of dendrimer graph which have  $n+1$  branches, which is denoted by  $H_n$ .

**Proposition 3.1.** Let  $H_n = (V, E)$  be the dendrimer graph in Figure 2, then:

$$W(H_n) = 3(3n - 5)4^n + 18 \times 2^n - 3.$$

**Proof.** The graph  $H_n$  consists of a starting vertex  $u_0$  and  $n+1$  branches such that the vertex set of the  $i$ '-th branch ( $i > 0$ ), has  $3 \times 2^{i-1}$  vertices and is denoted by  $V_i$  and  $|V_0| = 1$ . So we have:

$$|V| = 1 + \left| \bigcup_{i=0}^n V_i \right| = 1 + 3 \sum_{i=0}^{n-1} 2^i = 3 \times 2^n - 2.$$

Because of the symmetric structure of the graph  $G$  in Figure 2,  $d(u)$  for every vertex  $u$  in the  $n'$ -th branch is constant and doesn't depend on  $u$ , so we choose  $u_i$  as representative of the  $i'$ -th branch ( $0 \leq i \leq n$ ).

$$d(u_n) = \sum_{v \in V_n} d(u_n, v) + \sum_{v \in V-V_n} d(u_n, v). \tag{10}$$

$2/3$  vertices in  $n'$ -th branch have the same distance from  $u_i$  which is:

$$2d(u_n, u_0) = 2n.$$

And the distance of  $1/2$  of the rest vertices in this branch from  $u_n$  is:

$$2d(u_n, u_1) = 2(n-1).$$

By continuing in this way we have:

$$\begin{aligned} \sum_{v \in V_n} d(u_n, v) &= \frac{2}{3}(3 \times 2^{n-1}) \times 2d(u_n, u_0) + \frac{1}{2} \times \frac{1}{3}(3 \times 2^{n-1}) \times 2d(u_n, u_1) \\ &+ \frac{1}{4} \times \frac{1}{3}(3 \times 2^{n-1}) \times 2d(u_n, u_2) + \dots + \frac{1}{2^{n-1}} \times \frac{1}{3}(3 \times 2^{n-1}) 2d(u_n, u_{n-1}) \\ &= 2n \times 2^n + 2(n-1) \times 2^{n-2} + 2(n-2) \times 2^{n-3} + \dots + 2 \times 2^0 \\ &= n \times 2^n + \sum_{i=1}^n i 2^i = 2 + (3n-2) \times 2^n. \end{aligned} \tag{11}$$

Now because  $H_n$  is a tree, the path between any two vertices is unique and for every vertex  $v \in \bigcup_{i=0}^n V_i$  we have:

$$d(u_n, v) = 1 + d(u_{n-1}, v).$$

So:

$$\sum_{v \in \bigcup_{i=0}^{n-1} V_i} d(u_n, v) - \sum_{v \in \bigcup_{i=0}^{n-1} V_i} d(u_{n-1}, v) = |\bigcup_{i=0}^{n-1} V_i| = 3 \times 2^{n-1} - 2. \tag{12}$$

By (10), (11) and (12) we have:

$$\begin{aligned} d(u_n) - d(u_{n-1}) &= \sum_{v \in V_n} d(u_n, v) - \sum_{v \in V-V_n} d(u_{n-1}, v) \\ 2 + (3n-2)2^n + (3 \times 2^{(n-1)}) - 2 &= (6n-1)2^{n-1} - 2^n + 1 \\ d(u_n) &= \sum_{i=1}^n d(u_i) - d(u_{i-1}) = 7 + (6n-7)2^n \end{aligned}$$

If the Wiener index of  $H_n$  with  $n+1$  branches is denoted by  $W(n)$ , we have:

$$\begin{aligned} W(n) - W(n-1) &= 3 \times 2^{(n-1)}(7 + (6n-7)2^n) - \frac{1}{2}(3 \times 2^{n-1})(2 + (3n-2)2^n) \\ &= 18 \times 2^{n-1}((3n-4).2^{n-2} + 1) \end{aligned}$$

Therefore,

$$W(n) = \sum_{i=0}^n 18 \times 2^{i-1}((3i-4).2^{i-2} + 1) = 3(3n-5)4^n + 18 \times 2^n - 3.$$

And the proof is completed. ■



**Corollary 3.2.**  $Sz(H_n) = 3(3n - 5)4^n + 18 \times 2^n - 3$ .

**Proof.** The graph  $H_n$  is a tree so, by [11] the result is obtained. ■

**Corollary 3.3.**  $S(H_n) = 4^n (36n - 69) + 87(2^n) - 18$ .

**Proof.** Because  $H_n$  is a tree by [11] we have,  $S(G_n) = 4W(G_n) - n(n - 1)$ , where  $n$  is the number of vertices of  $H_n$ . Now by replacing the closed form of  $W(H_n)$  which was obtained from proposition 3.1, the proof is completed. ■

**Corollary 3.4.**  $Gut(G_n) = 4^n (36n - 78) + 105(2^n) - 97$ .

**Proof.** Because  $H_n$  is a tree by [11] we have,  $Gut(G_n) = 4W(G_n) - (2n - 1)(n - 1)$  which  $n$  is the number of vertices of  $H_n$  and by proposition 3.1 it is done. ■

**Corollary 3.5.**  $PI(H_n) = (3 \times 2^n - 3)(3 \times 2^n - 4)$ .

**Proof.** Because  $H_n$  is a tree so for every edge  $e = uv$  of  $H_n$  we have:

$$n_u(e | H_n) + n_v(e | H_n) = |V| = 3 \times 2^n - 2.$$

Subgraphs of  $H_n$  with vertex sets  $N_u(e | H_n)$  and  $N_v(e | H_n)$  both are trees, so the number of edges of them are  $n_u(e | H_n) - 1$  and  $n_v(e | H_n) - 1$  respectively. Then we have:

$$\begin{aligned} n_{eu}(e | H_n) + n_{ev}(e | H_n) &= n_u(e | H_n) + n_v(e | H_n) - 2 = 3 \times 2^n - 2 \\ PI(H_n) = |E| (3 \times 2^n - 4) &= (3 \times 2^n - 3)(3 \times 2^{n+1} - 4) \end{aligned}$$

**Proposition 3.6.** The hyper-Wiener index of  $H_n$  is:

$$WW(H_n) = \frac{1}{2} \left( (18n^2 - 51n + 81)4^n - 87(2^n) + 6 \right).$$

**Proof.** By the definition we have:

$$WW(H_n) = \frac{1}{2} (W(H_n) + \sum_{\{u,v\} \subseteq V} d^2(u,v)). \quad (14)$$

Because of the symmetric structure of the graph  $H_n$  Figure 2,  $d(u)$  for every vertex  $u$  in the  $n'$ -th branch is constant and doesn't depend on  $u$ , so we choose  $u_i$  as representative of the  $i'$ -th branch ( $0 \leq i \leq 1$ ).

$$d^2(u_n) = \sum_{v \in V} d^2(u_n, v) = \sum_{v \in \bigcup_{i=0}^{n-1} V_i} d^2(u_n, v) + \sum_{v \in V_n} d^2(u_n, v). \quad (15)$$

The graph  $H_n$  is a tree so, for any vertex  $v \in \bigcup_{i=0}^{n-1} V_i$ :

$$d(u_n, v) = 1 + d(u_{n-1}, v)$$

Now by (15) we have:

$$\begin{aligned} d^2(u_n) &= \sum_{v \in \bigcup_{i=0}^{n-1} V_i} (d(u_{n-1}, v) + 1)^2 + \sum_{v \in V_n} d^2(u_n, v) \\ &= \sum_{v \in \bigcup_{i=0}^{n-1} V_i} d^2(u_{n-1}, v) + 2 \sum_{v \in \bigcup_{i=0}^{n-1} V_i} d(u_{n-1}, v) + \sum_{v \in V_n} d^2(u_n, v) + 3 \times 2^{n-1} - 2 \quad (16) \\ &= d^2(u_{n-1}) + 2d(u_{n-1}) + \sum_{v \in V_n} d^2(u_n, v) + 3 \times 2^{n-1} - 2. \end{aligned}$$

$2/3$  vertices of the  $n$ -th branch have the same distance from  $u_n$  which is:

$$2d(u_n, u_0) = 2n,$$

and the distance of  $1/2$  of the rest vertices in this branch from  $u_n$  is:

$$\begin{aligned} \sum_{v \in V_n} d^2(u_n, v) &= \frac{2}{3}(3 \times 2^{n-1})(2n)^2 + \frac{1}{2} \cdot \frac{1}{3}(3 \times 2^{n-1})(2n-2)^2 \\ &\quad + \frac{1}{4} \cdot \frac{1}{3}(3 \times 2^{n-1})(2n-4)^2 + \dots + 2^0(2)^2 \quad (17) \\ &= 2^{n+2}n^2 + 2^n(n-1)^2 + 2^{(n-1)}(n-2)^2 + \dots + 2^0(2)^2 \\ &= 2^{n+1}n^2 + \sum_{i=1}^n 2^{i+1}i^2 = 2^{n+1}(3n^2 - 4n + 6) - 12. \end{aligned}$$

By (13) in the proof of the proposition 3.1, and considering (16), (17):

$$\begin{aligned} d^2(u_n) &= \sum_{i=1}^n d^2(u_i) - d^2(u_{i-1}) \\ &= \sum_{i=1}^n 2^{i-1}(12i^2 - 4i + 1) = (12n^2 - 28n + 41)2^n - 41 \end{aligned}$$

Therefore:

$$\begin{aligned} \sum_{\{u,v\} \subseteq V} d^2(u, v) - \sum_{\{u,v\} \subseteq \bigcup_{i=0}^{n-1} V_i} d^2(u, v) &= (3 \times 2^n - 2)d^2(u_n) - \sum_{\{u,v\} \subseteq V_n} d^2(u, v) \\ &= (3 \times 2^n - 2)d^2(u_n) - (3 \times 2^{n-2}) \sum_{v \in V_n} d^2(u_n, v). \end{aligned}$$

Now let,

$$F(i) = \sum_{\{u,v\} \subseteq \bigcup_{j=0}^i V_j} d^2(u, v).$$

So, we have:

$$\sum_{\{u,v\} \subseteq V} d^2(u, v) = \sum_{i=1}^n F(i) - F(i-1) = 6(3n^2 - 10n + 16)4^n - 105(2^n) + 9. \quad (18)$$

Now considering (18) and the formula of  $W(H_n)$  which was computed in Proposition 3.1, and replacing those in (14), the proof is done. ■

## REFERENCES

1. R. Entringer, Distance in graphs: Trees, *J. Combin. Math. Combin. Comput.* **24** (1997) 65–84.
2. I. Gutman, S. Klavžar, B. Mohar (Eds), Fifty years of the Wiener index, *MATCH Commun. Math. Comput. Chem.* **35** (1997) 1–259.
3. H. Hosoya, Topological Index, A Newly Proposed Quantity Characterizing the Topological Nature of Structural Isomers of Saturated Hydrocarbons, *Bull. Chem. Soc. Jpn.* **44** (1971) 2332–2339.
4. H. Wiener, Structural Determination of Paraffin Boiling Points, *J. Am. Chem. Soc.* **69** (1947) 17–20.
5. I. Gutman, A formula for the Wiener number of trees and its extension to graphs containing cycles, *Graph Theory Notes N.Y.* **27** (1994) 9–15.
6. S. Klavžar, A. Rajapakse, I. Gutman, The Szeged and the Wiener index of graphs, *Appl. Math. Lett.* **9** (1996) 45–49.
7. P. V. Khadikar, On a Novel structural descriptor PI, *Nat. Acad. Sci. Lett.* **23** (2000) 113–118.
8. P. V. Khadikar. S. Karmakar and V. K. Agrawal, Relationship and relative correction potential of the Wiener, Szeged and PI Indices, *Nat. Acad. Sci. Lett.* **23** (2000) 165–170.
9. H. P. Schultz, T. P. Schultz, Topological organic chemistry. 3. Graph theory, Binary and Decimal Adjacency Matrices, and Topological indices of Alkanes, *J. Chem. Inf. Comput. Sci.* **31** (1991) 144–147.
10. H. P. Schultz, T. P. Schultz, Topological organic chemistry. 6. Theory and topological indices of cycloalkanes, *J. Chem. Inf. Comput. Sci.* **33** (1993) 240–244.
11. I. Gutman, Selected properties of the Schultz molecular topological index, *J. Chem. Inf. Comput. Sci.* **34** (1994) 1087–1089.
12. I. Gutman, Y. N. Yeh, S. L. Lee, Y. L. Luo, Some recent results in the theory of the Wiener number, *Indian J. Chem.* **32** (1993) 651–661.
13. M. Randić, Novel molecular descriptor for structure–property studies, *Chem. Phys. Lett.* **211** (1993) 478–483.
14. D. J. Klein, I. Lukovits, I. Gutman, On the definition of the hyper–Wiener index for cycle-containing structures, *J. Chem. Inf. Comput. Sci.* **35** (1995) 50–52.

## The Topological Indices of some Dendrimer Graphs

M. R. DARAFSHEH<sup>a</sup>, M. NAMDARI<sup>b</sup> AND S. SHOKROLAHI<sup>b</sup>

<sup>a</sup> School of Mathematics, Statistics and Computer Science, College of Science, University of Tehran, Tehran, Iran

<sup>b</sup> Department of Mathematics, Shahid Chamran University of Ahvaz, Ahvaz, Iran

### اندیس های توپولوژیکی برخی گراف های دندریمر

ادیتور رابط : مسن یوسفی آذری

#### چکیده

در این مقاله، اندیس های وینر و وینر بالایی دو نوع گراف دندریمر محاسبه شده است. با به کارگیری فرمول به دست آمده برای اندیس وینر، اندیس های سگد، شولز، پادماکار- ایوان و گوتمن نیز برای این گراف ها تعیین می شود.

لغات کلیدی: اندیس توپولوژیکی، دندریمر، اندیس وینر، اندیس وینر بالایی