

Some Relations between Kekulé Structure and Morgan–Voyce Polynomials

İNCI GÜLTEKİN^{1,*} AND BÜNYAMİN ŞAHİN²

¹Department of Mathematics, Faculty of Science, Atatürk University, 25240 Erzurum, Turkey

²Department of Elementary Mathematics Education, Faculty of Education, Bayburt University, 69000 Bayburt, Turkey

ARTICLE INFO

Article History:

Received 2 March 2016

Accepted 9 May 2016

Published online 11 April 2017

Academic Editor: Michel Marie Deza

Keywords:

Kekulé structure

Hosoya index

Morgan–Voyce polynomials

Caterpillar trees

ABSTRACT

In this paper, Kekulé structures of benzenoid chains are considered. It has been shown that the coefficients of a $B_n(x)$ Morgan–Voyce polynomial equal to the number of k -matchings ($m(G, k)$) of a path graph which has $N = 2n + 1$ points. Furthermore, two relations are obtained between regularly zig–zag non-branched catacondensed benzenoid chains and Morgan–Voyce polynomials and between regularly zig–zag non branched catacondensed benzenoid chains and their corresponding caterpillar trees.

© 2017 University of Kashan Press. All rights reserved

1. INTRODUCTION

A benzenoid system is obtained by using the regular hexagons consecutively so that two hexagons are either disjoint or have a common edge [1]. An example of benzenoid chain is illustrated in Figure 1.




Figure 1. A Benzenoid Chain.

In connection with the benzenoid chains the LA -sequence is defined as an ordered h -tuple ($h > 1$) of the symbols L and A . The i -th symbol is L if the i -th hexagon is of

*Corresponding Author (Email: shnbnymn25@gmail.com)

DOI: 10.22052/ijmc.2017.49481.1177

mode L_1 or L_2 . The i -th symbol is A if the i -th hexagon is of mode A . The definition of L_1 , L_2 and A modes of hexagons is clear from Figure 2.




Figure 2. Illustration of L_1 , L_2 and A modes of hexagons, respectively.

For instance, the LA -sequence of the benzenoid chain in Figure 1 is $LLLALLLLAALL$ or, in the abbreviated form $L^3AL^2AL^3A^2L^2$. Each perfect matching of a benzenoid system (if any exists) represents a Kekulé structure. The number of Kekulé structures of benzenoid chains is called its “ K number”. The K -number of a benzenoid chain is calculated by its LA -sequence.

Balaban and Tomescu coined the term isarithmicity for the benzenoid chains which their K numbers are same [2]. It is denoted by $\langle x_1, x_2, \dots, x_n \rangle$ the class of isarithmic benzenoid chains with the LA -sequence

$$L^{x_1}AL^{x_2}A \dots AL^{x_n}$$

where $n \geq 1$, and $x_1 \geq 1$, $x_n \geq 1$, $x_i \geq 0$ for $i = 2, 3, \dots, n - 1$. For example isarithmic class of the benzenoid chain which is depicted in Figure 1 is $\langle 3, 2, 3, 0, 2 \rangle$.

Every benzenoid chain can be represented in this form. It is denoted by $K_n \langle x_1, x_2, \dots, x_n \rangle$ the number of Kekulé structures of the chain $\langle x_1, x_2, \dots, x_n \rangle$. It is defined for the initial terms of the K numbers such that ([1]) $K_0 = 1, K_1 \langle x_1 \rangle = 1 + x_1$.

Theorem 1. If $n \geq 2$ then for arbitrary $x_1 \geq 1$, $x_n \geq 1$, $x_i \geq 0$, ($i = 2, 3, \dots, n - 1$), the following recurrence relation holds [1]

$$K_n \langle x_1, x_2, \dots, x_n \rangle = (x_n + 1)K_{n-1} \langle x_1, x_2, \dots, x_{n-1} \rangle + K_{n-2} \langle x_1, x_2, \dots, x_{n-2} \rangle.$$

2. THE HOSOYA INDEX AND MORGAN-VOYCE POLYNOMIALS

The Hosoya or Z -index was defined by Hosoya in 1971 [3] and the Hosoya index of a graph G is denoted by $Z(G)$. The $Z(G)$, is the total number of k -matchings which are the number of k choosing from a graph G such that the k lines are non-adjacent where N is the number of points.

Definition 1. The number of k -matchings is denoted by $m(G, k)$ and the $Z(G)$ is defined as $Z(G) = \sum_{k=0}^{\lfloor N/2 \rfloor} m(G, k)$ such that $m(G, 0) = 1$ for any graph G .

Theorem 2. The number of k -matchings of the path graph is calculated by the following equation [4]

$$m(G, k) = \binom{N-k}{k}, \text{ for } 0 \leq k \leq \lfloor N/2 \rfloor.$$

Relations between topological indices and some orthogonal polynomials for example Hermite, Laguerre and Chebyshev polynomials were found by Hosoya ([5]). Another relation between the sextet polynomial of a hexagonal chain and the matching polynomial of a caterpillar tree was discovered by Gutman [6]. As a result of this paper, it has been shown that the K -number of a hexagonal chain is equal to the Hosoya index of the corresponding caterpillar [7]. For instance, corresponding caterpillar tree of the hexagonal chain which is depicted in Figure 1 is on the below.




Figure 3. The hexagonal chain in Figure 1 has 14 hexagons and the corresponding caterpillar tree has 14 edges.

The caterpillar tree of the hexagonal chain in Figure 3 is $C_5(4, 3, 4, 1, 3)$.

Definition 2. The Morgan–Voyce polynomials $B_n(x)$ is defined by [8] as

$$B_n(x) = \sum_{i=0}^n \binom{n+i+1}{n-i} x^i$$

and the first five Morgan–Voyce polynomials are found from this equation like that

$$\begin{aligned} B_0(x) &= 1 \\ B_1(x) &= x + 2 \\ B_2(x) &= x^2 + 4x + 3 \\ B_3(x) &= x^3 + 6x^2 + 10x + 4 \\ B_4(x) &= x^4 + 8x^3 + 21x^2 + 20x + 5. \end{aligned}$$

3. REGULARLY ZIG–ZAG NON–BRANCHED CATA CONDENSED BENZENOIDS

The Kekulé number of regularly zig–zag non-branched cata condensed benzenoids was found by He, He and Xie [9] by Peak–Valley matrix.




Figure 4. Dualist graph of a general non-branched cata-condensed benzenoids.

In Figure 4, $a_i \in (i = 1, 2, \dots, s)$ and $b_i \in (i = 1, 2, \dots, s')$ where $s' = s$ for Figure 4(a) and $s' = s + 1$ for Figure 4(b). $a_i + 1$ and $b_i + 1$ represent the numbers of linearly condensed six-membered rings horizontally and diagonally, respectively. For the benzenoid shown in Figure 4(a) and 4(b), the Peak-Valley matrix is as follows.

$$A_n = \begin{bmatrix} t_1 & 1 & 0 & & & \\ 1 & t_2 & 1 & & 0 & \\ 0 & 1 & t_3 & & & \\ & & & \ddots & 1 & 0 \\ & 0 & & 1 & t_{-1} & 1 \\ & & & 0 & 1 & t \end{bmatrix}$$

where $t_i = \begin{cases} b_{k+1} + 2, & \text{if } i = \sum_{j=0}^k a_j + 1 \\ 2, & \text{if } i \neq \sum_{j=0}^k a_j + 1 \end{cases}$, $k = 1, 2, \dots, s$; $i = 1, 2, \dots, n$. Here n is the number of peaks (or valleys) in a graph G . The Kekulé number of a graph G is shown by $K_n(G)$ ($n = 1, \dots, n$).




Figure 5. Simple binary regularly cata-condensed benzenoids.

Lemma 1. From Figure 5, the K -number of the graph G is calculated by the following tri-diagonal determinantal expression[9]:

$$K_n(G) = \det A_n = \begin{vmatrix} b+2 & 1 & 0 & & & \\ 1 & b+2 & 1 & & 0 & \\ 0 & 1 & b+2 & & & \\ & & & \ddots & 1 & 0 \\ & & 0 & & 1 & b+2 & 1 \\ & & & & 0 & 1 & b+2 \end{vmatrix}.$$

The order of the above determinant is $s + 1$, where s is the repeat times of horizontal linear segments on the graph G .

4. CONTINUANTS AND CATERPILLAR TREES

Lemma 2. If H is a hexagonal chain whose LA -sequence is $L^{x_1}AL^{x_2}A \dots L^{x_{n-1}}AL^{x_n}$, then the number $K(H)$ of its Kekulé structures is equal to the Z -index of the caterpillar tree $C_n(x_1, x_2, \dots, x_n)$ [7].

If it is written $C(H)$ for caterpillar tree of a H hexagonal chain, Lemma 2 is equivalent to the equality $K(H) = Z(C(H))$.

Definition 3. The continuants (or continuant polynomials) are introduced by Euler [10] as $L_n(x_1, x_2, \dots, x_n) = x_n L_{n-1}(x_1, x_2, \dots, x_{n-1}) + L_{n-2}(x_1, x_2, \dots, x_{n-2})$ with initial conditions $L_0() = 1$, $L_1(x_1) = x_1$ and $L_2(x_1, x_2) = x_1 x_2 + 1$.

From this it is shown that the Z -index of the caterpillar trees coincides with Euler's continuant like the following lemma.

Lemma 3. $Z(C_n(x_1, x_2, \dots, x_n)) = L_n(x_1, x_2, \dots, x_n)[7]$.

5. MAIN RESULTS

Theorem 3. The coefficients of a $B_n(x)$ Morgan–Voyce polynomial are equal to the number of k -matchings ($m(G, k)$) of a path graph which has $N = 2n + 1$ points.

Proof. We denote the coefficients of Morgan–Voyce polynomials with

$$C(B_n(x)) = \binom{n+i+1}{n-i}$$

such that $0 \leq i \leq n$ and we take the point number of the path graph $N = 2n + 1$. The number of k -matchings of a path graph for $0 \leq k \leq \lfloor N/2 \rfloor$ is

$$m(G, k) = \binom{N-k}{k}$$

and $\lfloor N/2 \rfloor = \lfloor (2n + 1)/2 \rfloor = n$ by the definition of the Hosoya index. Now we demonstrate the coefficients of the Morgan–Voyce polynomials in combinatorial form with respectively for $0 \leq i \leq n$

$$C(B_n(x)) = \binom{n+1}{n}, \binom{n+2}{n-1}, \dots, \binom{2n}{1}, \binom{2n+1}{0}$$

and $m(G, k) = \binom{N-k}{k}$ for $0 \leq k \leq \lfloor N/2 \rfloor = n$ with respectively

$$m(G, k) = \binom{2n+1}{0}, \binom{2n}{1}, \dots, \binom{n+2}{n-1}, \binom{n+1}{n}.$$

It is clear that $C(B_n(x))$ and $m(G, k)$ are same in reverse order. From this we say for every n^{th} degree Morgan–Voyce polynomial there is a path graph (P_N) which has $N = 2n + 1$ points such that the coefficients of the Morgan–Voyce polynomials equal to the number of k -matchings of P_N .

Example 1. We show an application of the previous theorem for the first three Morgan–Voyce polynomials. For $B_0(x)$, $C(B_0(x)) = 1$ equals to $m(G, k)$ for $N = 2 \times 0 + 1 = 1$. For $B_1(x)$, $C(B_1(x)) = 1, 2$ equal to $m(G, k)$ for $N = 2 \times 1 + 1 = 3$. For $B_2(x)$, $C(B_2(x)) = 1, 4, 3$ equal to $m(G, k)$ for $N = 2 \times 2 + 1 = 5$.

Lemma 4. If $b_1 + 1 = b_2 + 1 = \dots = b_s + 1 = b + 1$ (numbers of the regular hexagons on diagonal wise are same) like in Figure 5 and we take x instead of b_i , then

(the right equation is used to express many properties of the Morgan–Voyce polynomials like in [8])

$$K_n(G) = \det A_n = B_n(x).$$

Proof.

$$\begin{aligned} K_1(G) &= \begin{vmatrix} x+2 \end{vmatrix} &= x+2 &= B_1(x) \\ K_2(G) &= \begin{vmatrix} x+2 & 1 \\ 1 & x+2 \end{vmatrix} &= (x+2)(x+2) - 1 &= x^2 + 4x + 3 = B_2(x) \\ K_3(G) &= \begin{vmatrix} x+2 & 1 & 0 \\ 1 & x+2 & 1 \\ 0 & 1 & x+2 \end{vmatrix} &= x^3 + 6x^2 + 10x + 4 &= B_3(x) \end{aligned}$$

and by the determinant of the tri-diagonal matrix in Lemma 1,

$$K_n(G) = B_n(x) = (x + 2)B_{n-1}(x) - B_{n-2}(x).$$

In Lemma 1, the (n) indice on the notatin K_n is the number of the repetition of the diagonal hexagons. We also take the number of the hexagons $b_i + 1$ on diagonal wise like the previous lemma. For Figure 5, $b_1 + 1 = b_2 + 1 = \dots = b_s + 1 = b + 1$ and its corresponding caterpillar tree is $C_{2n}(b + 1, 1, b, 1, \dots, b, 1)$.

There is a relation between the K -number of the hexagonal chain in Figure 5 and Z -index of its corresponding caterpillar tree as noted in the next theorem.

Theorem 4. $K_n(G) = Z(C_{2n}(G))$.

Proof. Induct on n . For $n = 1$, $K_1(G) = Z(C_2(b + 1, 1)) = b + 2$, as desired. We assume that the equality is true for $n \leq k$ and we will show that it is true for $n = k + 1$. This means

$$K_{k+1}(G) = Z(C_{2k+2}(b + 1, 1, b, 1, \dots, b, 1)).$$

By assumption

$$K_k(G) = Z(C_{2k}(b + 1, 1, b, 1, \dots, b, 1))$$

and

$$K_{k-1}(G) = Z(C_{2k-2}(b + 1, 1, b, 1, \dots, b, 1)).$$

By Lemma 1,

$$\begin{aligned} K_{k+1}(G) &= (b + 2)K_k(G) - K_{k-1}(G) \\ &= (b + 2)Z(C_{2k}(G)) - Z(C_{2k-2}(G)) \\ &= bZ(C_{2k}(G)) + 2[Z(C_{2k-1}(G)) + Z(C_{2k-2}(G))] - Z(C_{2k-2}(G)) \\ &= bZ(C_{2k}(G)) + Z(C_{2k-1}(G)) + Z(C_{2k-1}(G)) + Z(C_{2k-2}(G)) \\ &= Z(C_{2k+1}(G)) + Z(C_{2k}(G)) = Z(C_{2k+2}(G)) \end{aligned}$$

This complete the proof.

Example 2. We calculate the Kekulé number of simple binary regularly catacondensed benzenoid in Figure 5 by two ways mentioned in the Theorem 4. The matrix form of K -number of the chain shown in Figure 5 is

$$K_3(G) = \begin{bmatrix} 4 & 1 & 0 \\ 1 & 4 & 1 \\ 0 & 1 & 4 \end{bmatrix}$$

and $K_3(G) = \det A = 56$. Now we use the corresponding caterpillar tree of the hexagonal chain as the follows:




Figure 6. The hexagonal chain in Figure 5 has 9 hexagons and the corresponding caterpillar tree has 9 edges.

This caterpillar tree is denoted by $C_6(3, 1, 2, 1, 2, 1)$ and $Z(C_6(3, 1, 2, 1, 2, 1)) = 56$. So that $K_3(G) = Z(C_6(3, 1, 2, 1, 2, 1))$.

ACKNOWLEDGEMENT

B.ŞAHİN thanks to the Scientific and Technological Research Council of Turkey (TUBITAK) for support.

REFERENCES

1. R. Tošić, I. Stojmenović, Chemical graphs, Kekulé structures and Fibonacci numbers, *Zb. Rad. Prirod.-Mat. Fak. Ser. Mat.* **25** (2) (1995) 179–195.
2. A. T. Balaban, I. Tomescu, Algebraic expressions for the number of Kekulé structure of isoarithmic cata-condensed benzenoid polycyclic hydrocarbons, *MATCH Commun. Math. Comput. Chem.* **14** (1983) 155–182.
3. H. Hosoya, Topological index. A newly proposed quantity characterizing the topological nature of structural isomers of saturated hydrocarbons, *Bull. Chem. Soc. Jpn.* **44** (1971) 2332–2339.
4. H. Hosoya, Topological index and Fibonacci numbers with relation to chemistry, *Fibonacci Quart.* **11** (1973) 255–269.
5. H. Hosoya, Graphical and combinatorial aspects of some orthogonal polynomials, *Natur. Sci. Rep. Ochanomizu Univ.* **32** (2) (1981) 127–138.

6. I. Gutman, Topological properties of benzenoid systems. An identity for the sextet polynomial, *Theor. Chim. Acta* **45** (1977) 309–315.
7. H. Hosoya, I. Gutman, Kekulé structures of hexagonal chains—some unusual connections, *J. Math. Chem.* **44** (2008) 559–568.
8. T. Koshy, *Fibonacci and Lucas numbers with applications*, Pure and Applied Mathematics (New York), Wiley–Interscience, New York, 2001.
9. W. J. He, W. C. He, S. L. Xie, Algebraic expressions for Kekulé structure counts of nonbranched cata–condensed benzenoid, *Discrete Appl. Math.* **35** (1992) 91–106.
10. R. L. Graham, D. E. Knuth, O. Patashnik, *Concrete Mathematics. A Foundation for Computer Science*, Addison–Wesley, Reading, 1989.

Some Relation between Kekule Structure and Morgan-Voyce Polynomials

INCI GULTEKIN¹ AND BUNYAMIN SAHIN²

¹Department of Mathematics, Faculty of Science, Ataturk University, 25240 Erzurum, Turkey

²Department of Elementary Mathematics Education, Faculty of Education, Bayburt University, 69000 Bayburt, Turkey

بررسی از روابط میان ساختار ککول و چندجمله‌ایهای مورگان-ووویس

ادیتور رابط: میشل ماری دزا

چکیده

در این مقاله، ساختارهای ککول از زنجیره‌های بنزوئیدی مورد توجه قرار گرفته‌اند. نشان داده شده است که ضرایب چندجمله‌ای مورگان-ووویس $B_n(x)$ ، با تعداد k -جفت‌های $(m(G, K))$ از یک گراف مسیر که $N=2n+1$ رأس دارد، برابر است. بعلاوه، دو رابطه بین زنجیره‌های به‌طور منظم بنزوئیدی کاملاً متراکم غیر شاخه‌ای مورب و چندجمله‌ایهای مورگان-ووویس، و زنجیره‌های به‌طور منظم بنزوئیدی کاملاً متراکم غیر شاخه‌ای مورب و درختان کاترپیلار متناظر به دست آمده است.

لغات کلیدی: ساختار ککول، شاخص هوسویا، چندجمله‌ایهای مورگان-ووویس، درختان کاترپیلار