

A Note on Revised Szeged Index of Graph Operations

NASRIN DEHGARDI*

Department of Mathematics and Computer Science, Sirjan University of Technology, Sirjan, I. R. Iran

ARTICLE INFO

Article History:

Received 30 July 2016

Accepted 8 June 2017

Published online 6 January 2018

Academic Editor: Hassan Yousefi –Azari

Keywords:

Topological index

Revised Szeged index

Graph operation

ABSTRACT

Let G be a finite and simple graph with edge set $E(G)$. The revised Szeged index is defined as

$$Sz^*(G) = \sum_{e=uv \in E(G)} \left(n_u(e|G) + \frac{n_G(e)}{2} \right) \left(n_v(e|G) + \frac{n_G(e)}{2} \right),$$

where $n_u(e|G)$ denotes the number of vertices in G lying closer to u than to v and $n_G(e)$ is the number of equidistant vertices of e in G . In this paper, we compute the revised Szeged index of the join and corona product of graphs.

© 2018 University of Kashan Press. All rights reserved

1 INTRODUCTION

Let G be a finite and simple graph with vertex set $V = V(G)$ and edge set $E = E(G)$. The integers $n = n(G) = |V(G)|$ and $m = m(G) = |E(G)|$ are the order and the size of the graph G , respectively. For a vertex $v \in V(G)$, the *open neighborhood* of v , denoted by $N_G(v) = N(v)$ is the set $\{u \in V(G) \mid uv \in E(G)\}$. The *degree* of $v \in V(G)$, denoted by $d_G(v)$, is defined by $d_G(v) = |N_G(v)|$. Let $u, v \in V(G)$, then the *distance* $d_G(u, v)$ between u and v is defined as the length of any shortest path in G connecting u and v . We consult [14] for notation and terminology which are not defined here.

The first and second Zagreb indices are defined as $M_1(G) = \sum_{u \in V(G)} d_G^2(u)$ and $M_2(G) = \sum_{uv \in E(G)} d_G(u)d_G(v)$, respectively. Furtula and Gutman [5] defined the

* Corresponding Author (Email address: n.dehgardi@sirjantech.ac.ir)
DOI: 10.22052/ijmc.2017.58647.1228

forgotten topological index as $F(G) = \sum_{uv \in E(G)} (d_G^2(u) + d_G^2(v))$. The interested readers are referred to [3,7] for more information on this topic.

A vertex $w \in V(G)$, is said to be equidistant from the edge $e = uv$ of G if $d_G(u, w) = d_G(v, w)$. The number of equidistant vertices of e is denoted by $n_G(e)$. Let uv be an edge of G . Define the sets $N(u, G) = \{x \in V(G) | d_G(u, x) < d_G(v, x)\}$ and $N(v, G) = \{x \in V(G) | d_G(v, x) < d_G(u, x)\}$ consisting, respectively, of vertices of G lying closer to u than to v , and lying closer to v than to u . The number of such vertices is then $n_u(e|G) = |N(u, G)|$ and $n_v(e|G) = |N(v, G)|$. Note that vertices equidistant to u and v are not included into either $N(u, G)$ or $N(v, G)$. It also worth noting that $u \in N(u, G)$ and $v \in N(v, G)$, which implies that $n_u(e|G) \geq 1$ and $n_v(e|G) \geq 1$. The Szeged index $Sz(G)$ was introduced by Gutman [6]. It is defined as $Sz(G) = \sum_{e=uv \in E(G)} n_u(e|G)n_v(e|G)$.

The Szeged index in graphs is well studied in the literature, see for example [9,10]. Randić [13] observed that the Szeged index does not take into account the contributions of the vertices at equal distances from the endpoints of an edge, and so he conceived a modified version of the Szeged index which is named as the revised Szeged index. The revised Szeged index of a connected graph G is defined as $Sz^*(G) = \sum_{e=uv \in E(G)} (n_u(e|G) + \frac{n_G(e)}{2})(n_v(e|G) + \frac{n_G(e)}{2})$. Nagarajan et al. [11] obtained the revised Szeged index of the Cartesian product of two connected graphs. In this paper we compute the revised Szeged index of the join and corona product of graphs. Readers interested in more information on computing topological indices of graph operations can be referred to [1,2,4,8,12].

2. MAIN RESULTS

In this section, we compute the revised Szeged index of the join and corona product of graphs. We let for every edge $e = uv \in E(G)$, $t_{uv}(G) = |N_G(u) \cap N_G(v)|$.

2.1. THE JOIN OF GRAPHS

The join $G = G_1 + G_2$ of graphs G_1 and G_2 with disjoint vertex sets V_1 and V_2 and edge sets E_1 and E_2 is the graph union $G_1 \cup G_2$ together with all the edges joining V_1 and V_2 . Obviously, $|V(G)| = |V_1| + |V_2|$ and $|E(G)| = |E_1| + |E_2| + |V_1||V_2|$.

Theorem 1. Let G_1 be a graph of order n_1 and of size m_1 and let G_2 be a graph of order n_2 and of size m_2 . If $G = G_1 + G_2$, then

$$Sz^*(G) = \frac{2M_2(G_1) + 2M_2(G_2) - n_2M_1(G_1) - n_1M_1(G_2) - F(G_1) - F(G_2)}{4} + \frac{4n_1^2n_2^2 + 8m_1m_2 + m_1(n_1^2 + 6n_1n_2 - 3n_2^2) + m_2(n_2^2 + 6n_1n_2 - 3n_1^2)}{4}.$$

Proof. By definition, $Sz^*(G) = \sum_{uv \in E(G)} (n_u(e|G) + \frac{n_G(e)}{2})(n_v(e|G) + \frac{n_G(e)}{2})$. We partition the edges of G in to three subset E_1, E_2 and E_3 , as $E_1 = \{e = uv \mid u, v \in V(G_1)\}$, $E_2 = \{e = uv \mid u, v \in V(G_2)\}$ and $E_3 = \{e = uv \mid u \in V(G_1), v \in V(G_2)\}$.

Let $e = uv \in E_1$. If $w \in V(G_2)$ or $w \in N_{G_1}(u) \cap N_{G_1}(v)$, then $d_G(u, w) = d_G(v, w) = 1$ and if $w \notin N_{G_1}(u) \cup N_{G_1}(v)$, then $d_G(u, w) = d_G(v, w) = 2$. Hence $n_u(e|G) = d_{G_1}(u) - t_{uv}(G_1) + 1, n_v(e|G) = d_{G_1}(v) - t_{uv}(G_1) + 1$ and $n_G(e) = n_1 + n_2 + 2t_{uv}(G_1) - (d_{G_1}(u) + d_{G_1}(v)) - 2$. Then for every edge $e = uv \in E_1$,

$$\begin{aligned} \left(n_u(e|G) + \frac{n_G(e)}{2}\right) \left(n_v(e|G) + \frac{n_G(e)}{2}\right) &= \left(\frac{n_1+n_2+d_{G_1}(u)-d_{G_1}(v)}{2}\right) \left(\frac{n_1+n_2+d_{G_1}(v)-d_{G_1}(u)}{2}\right) \\ &= \frac{(n_1+n_2)^2}{4} + \frac{d_{G_1}(u)d_{G_1}(v)}{2} - \frac{d_{G_1}^2(u)+d_{G_1}^2(v)}{4}. \end{aligned}$$

Therefore

$$\begin{aligned} \sum_{uv \in E_1} \left(n_u(e|G) + \frac{n_G(e)}{2}\right) \left(n_v(e|G) + \frac{n_G(e)}{2}\right) &= \sum_{uv \in E_1} \frac{(n_1+n_2)^2}{4} + \sum_{uv \in E_1} \frac{d_{G_1}(u)d_{G_1}(v)}{2} \\ &\quad - \sum_{uv \in E_1} \frac{d_{G_1}^2(u)+d_{G_1}^2(v)}{4} \\ &= \frac{(n_1+n_2)^2}{4} m_1 + \frac{M_2(G_1)}{2} - \frac{F(G_1)}{4}. \end{aligned} \tag{1}$$

Similarly,

$$\sum_{uv \in E_2} \left(n_u(e|G) + \frac{n_G(e)}{2}\right) \left(n_v(e|G) + \frac{n_G(e)}{2}\right) = \frac{(n_1+n_2)^2}{4} m_2 + \frac{M_2(G_2)}{2} - \frac{F(G_2)}{4}. \tag{2}$$

Let $e = uv \in E_3$ such that $u \in V(G_1)$ and $v \in V(G_2)$. If $w \in N_{G_1}(u) \cup N_{G_2}(v)$, then $d_G(u, w) = d_G(v, w) = 1$. Hence $n_u(e|G) = n_2 - d_{G_2}(v) + 1, n_v(e|G) = n_1 - d_{G_1}(u) + 1$ and $n_G(e) = d_{G_1}(u) + d_{G_2}(v) - 2$. Then for every edge $e = uv \in E_3$,

$$\begin{aligned} \left(n_u(e|G) + \frac{n_G(e)}{2}\right) \left(n_v(e|G) + \frac{n_G(e)}{2}\right) &= \left(\frac{2n_2+d_{G_1}(u)-d_{G_2}(v)}{2}\right) \left(\frac{2n_1+d_{G_2}(v)-d_{G_1}(u)}{2}\right) \\ &= n_1 n_2 + \frac{n_1-n_2}{2} d_{G_1}(u) + \frac{n_2-n_1}{2} d_{G_2}(v) \\ &\quad - \frac{d_{G_1}^2(u)}{4} - \frac{d_{G_2}^2(v)}{4} + \frac{d_{G_1}(u)d_{G_2}(v)}{2}. \end{aligned}$$

Set $Y = \sum_{uv \in E_3} \left(n_u(e|G) + \frac{n_G(e)}{2}\right) \left(n_v(e|G) + \frac{n_G(e)}{2}\right)$. Then,

$$\begin{aligned} Y &= \sum_{uv \in E_3} n_1 n_2 + \sum_{uv \in E_3} \frac{n_1-n_2}{2} d_{G_1}(u) + \sum_{uv \in E_3} \frac{n_2-n_1}{2} d_{G_2}(v) - \sum_{uv \in E_3} \frac{d_{G_1}^2(u)}{4} \\ &\quad - \sum_{uv \in E_3} \frac{d_{G_2}^2(v)}{4} + \sum_{uv \in E_3} \frac{d_{G_1}(u)d_{G_2}(v)}{2} \end{aligned}$$

$$= n_1^2 n_2^2 + m_1 n_2 (n_1 - n_2) + m_2 n_1 (n_2 - n_1) + 2m_1 m_2 - \frac{n_2 M_1(G_1)}{4} - \frac{n_1 M_1(G_2)}{4}. \quad (3)$$

By Equations (1), (2) and (3), we have:

$$\begin{aligned} Sz^*(G) &= \frac{(n_1+n_2)^2}{4} m_1 + \frac{M_2(G_1)}{2} - \frac{F(G_1)}{4} + \frac{(n_1+n_2)^2}{4} m_2 + \frac{M_2(G_2)}{2} - \frac{F(G_2)}{4} + n_1^2 n_2^2 + 2m_1 m_2 \\ &\quad - \frac{n_1 M_1(G_2)}{4} + m_1 n_2 (n_1 - n_2) + m_2 n_1 (n_2 - n_1) - \frac{n_2 M_1(G_1)}{4} \\ &= \frac{2M_2(G_1) + 2M_2(G_2) - n_2 M_1(G_1) - n_1 M_1(G_2) - F(G_1) - F(G_2)}{4} \\ &\quad + \frac{4n_1^2 n_2^2 + 8m_1 m_2 + m_1 (n_1^2 + 6n_1 n_2 - 3n_2^2) + m_2 (n_2^2 + 6n_1 n_2 - 3n_1^2)}{4}. \end{aligned}$$

Let P_n , $n \geq 2$ and C_n , $n \geq 3$ denote the path and the cycle on n vertices, respectively.

Corollary 2. The following equalities are hold:

1. $Sz^*(P_n + P_m) = \frac{4n^2 m^2 + n^3 + m^3 + 3nm^2 + 3mn^2 + 2n^2 + 2m^2 - 2n - 2m - 12nm + 4}{4}$.
2. $Sz^*(P_n + C_m) = \frac{4n^2 m^2 + n^3 + m^3 + 3nm^2 + 3mn^2 - n^2 + 3m^2 - 2m - 6nm - 2}{4}$.
3. $Sz^*(C_n + C_m) = \frac{4n^2 m^2 + n^3 + m^3 + 3nm^2 + 3mn^2}{4}$.

2.2. THE CORONA PRODUCT OF GRAPHS

The corona product $G = G_1 o G_2$ of graphs G_1 and G_2 with disjoint vertex sets V_1 and V_2 and edge sets E_1 and E_2 is as the graph obtained by taking one copy of G_1 and $|V_1|$ copies of G_2 and joining the i -th vertex of G_1 to every vertex in i -th copy of G_2 . Obviously, $|V(G)| = |V_1| + |V_1||V_2|$ and $|E(G)| = |E_1| + |V_1||E_2| + |V_1||V_2|$.

Theorem 3. Let G_1 be a graph of order n_1 and of size m_1 and let G_2 be a graph of order n_2 and of size m_2 . If $G = G_1 o G_2$, then

$$\begin{aligned} Sz^*(G) &= \frac{(n_1 n_2 + n_1)^2}{4} (m_1 + m_2) + n_1 n_2 (n_1 n_2 + n_1 - 1) \\ &\quad + n_1 m_2 (n_1 n_2 + n_1 - 2) - \frac{(n_2 + 1)^2}{4} \sum_{uv \in E_1} (n_u^2(e|G_1) + n_v^2(e|G_1)) \\ &\quad + \frac{2(n_2 + 1)^2 Sz(G_1) + 2M_2(G_2) - n_1 M_1(G_2) - F(G_2)}{4}. \end{aligned}$$

Proof. By definition, $Sz^*(G) = \sum_{uv \in E(G)} (n_u(e|G) + \frac{n_G(e)}{2})(n_v(e|G) + \frac{n_G(e)}{2})$. We partition the edges of G in to three subsets E_1 , E_2 and E_3 , as $E_1 = \{e = uv \mid u, v \in V(G_1)\}$, $E_2 = \{e = uv \mid u, v \in V(G_2)\}$ and $E_3 = \{e = uv \mid u \in V(G_1), v \in V(G_2)\}$. Let $e = uv \in E_1$. Then for each vertex w closer to u than v , the vertices of the copy of G_2 attached to w are

also closer to u than v . Since each copy of G_2 has exactly n_2 vertices, then $n_u(e|G) = (n_2 + 1)n_u(e|G_1)$. Similarly $n_v(e|G) = (n_2 + 1)n_v(e|G_1)$. Then $n_G(e) = n_1n_2 + n_1 - (n_2 + 1)n_u(e|G_1) - (n_2 + 1)n_v(e|G_1)$. Hence for every edge $e = uv \in E_1$,

$$\left(n_u(e|G) + \frac{n_G(e)}{2}\right)\left(n_v(e|G) + \frac{n_G(e)}{2}\right) = \frac{(n_1n_2+n_1)^2}{4} + \frac{(n_2+1)^2n_u(e|G_1)n_v(e|G_1)}{2} - \frac{(n_2+1)^2(n_u^2(e|G_1)+n_v^2(e|G_1))}{4}.$$

Define $Z = \sum_{uv \in E_1} \left(n_u(e|G) + \frac{n_G(e)}{2}\right)\left(n_v(e|G) + \frac{n_G(e)}{2}\right)$. Then,

$$\begin{aligned} Z &= \sum_{uv \in E_1} \frac{(n_1n_2+n_1)^2}{4} + \sum_{uv \in E_1} \frac{(n_2+1)^2n_u(e|G_1)n_v(e|G_1)}{2} - \sum_{uv \in E_1} \frac{(n_2+1)^2(n_u^2(e|G_1)+n_v^2(e|G_1))}{4} \\ &= \frac{(n_1n_2+n_1)^2}{4} m_1 + \frac{(n_2+1)^2 Sz(G_1)}{2} - \frac{(n_2+1)^2}{4} \sum_{e=uv \in E_1} (n_u^2(e|G_1) + n_v^2(e|G_1)). \end{aligned} \quad (4)$$

Let $e = uv \in E_2$. If $w \in V(G_2)$ and $w \in N_{G_2}(u) \cap N_{G_2}(v)$, then $d_G(u, w) = d_G(v, w) = 1$ and if $w \notin N_{G_2}(u) \cup N_{G_2}(v)$, then $d_G(u, w) = d_G(v, w) = 2$. Hence $n_u(e|G) = d_{G_2}(u) - t_{uv}(G_2) + 1$, $n_v(e|G) = d_{G_2}(v) - t_{uv}(G_2) + 1$ and $n_G(e) = n_1n_2 + n_1 + 2t_{uv}(G_2) - (d_{G_2}(u) + d_{G_2}(v)) - 2$. Hence for every edge $e = uv \in E_2$,

$$\begin{aligned} \left(n_u(e|G) + \frac{n_G(e)}{2}\right)\left(n_v(e|G) + \frac{n_G(e)}{2}\right) &= \left(\frac{n_1n_2+n_1+d_{G_2}(u)-d_{G_2}(v)}{2}\right)\left(\frac{n_1n_2+n_1+d_{G_2}(v)-d_{G_2}(u)}{2}\right) \\ &= \frac{(n_1n_2+n_1)^2}{4} + \frac{d_{G_2}(u)d_{G_2}(v)}{2} - \frac{d_{G_2}^2(u)+d_{G_2}^2(v)}{4}. \end{aligned}$$

Therefore

$$\begin{aligned} \sum_{uv \in E_2} \left(n_u(e|G) + \frac{n_G(e)}{2}\right)\left(n_v(e|G) + \frac{n_G(e)}{2}\right) &= \sum_{uv \in E_2} \frac{(n_1n_2+n_1)^2}{4} + \sum_{uv \in E_2} \frac{d_{G_2}(u)d_{G_2}(v)}{2} \\ &\quad - \sum_{uv \in E_2} \frac{d_{G_2}^2(u)+d_{G_2}^2(v)}{4} \\ &= \frac{(n_1n_2+n_1)^2}{4} m_2 + \frac{M_2(G_2)}{2} - \frac{F(G_2)}{4}. \end{aligned} \quad (5)$$

Let $e = uv \in E_3$ such that $u \in V(G_1)$ and $v \in V(G_2)$. Hence $n_u(e|G) = n_1n_2 + n_1 - d_{G_2}(v) - 1$. Since $v \in N(v, G)$, we have $n_v(e|G) = 1$ and so $n_G(e) = d_{G_2}(v)$. Hence

$$\begin{aligned} \left(n_u(e|G) + \frac{n_G(e)}{2}\right)\left(n_v(e|G) + \frac{n_G(e)}{2}\right) &= \left(\frac{2(n_1n_2+n_1-1)-d_{G_2}(v)}{2}\right)\left(\frac{2+d_{G_2}(v)}{2}\right) \\ &= (n_1n_2 + n_1 - 1) + \frac{n_1n_2+n_1-2}{2} d_{G_2}(v) - \frac{d_{G_2}^2(v)}{4}. \end{aligned}$$

Therefore,

$$\begin{aligned} \sum_{uv \in E_3} \left(n_u(e|G) + \frac{n_G(e)}{2}\right)\left(n_v(e|G) + \frac{n_G(e)}{2}\right) &= \sum_{uv \in E_3} (n_1n_2 + n_1 - 1) - \sum_{uv \in E_3} \frac{d_{G_2}^2(v)}{4} \\ &\quad + \sum_{e=uv \in E_3} \frac{n_1n_2+n_1-2}{2} d_{G_2}(v) \end{aligned}$$

$$= n_1 n_2 (n_1 n_2 + n_1 - 1) - \frac{n_1 M_1(G_2)}{4} + n_1 m_2 (n_1 n_2 + n_1 - 2). \quad (6)$$

By Equations (4), (5) and (6), we have:

$$\begin{aligned} Sz^*(G) &= \frac{(n_1 n_2 + n_1)^2}{4} m_1 + \frac{(n_2 + 1)^2 Sz(G_1)}{2} - \frac{(n_2 + 1)^2}{4} \sum_{uv \in E_1} (n_u^2(e|G_1) + n_v^2(e|G_1)) \\ &+ \frac{(n_1 n_2 + n_1)^2}{4} m_2 + \frac{M_2(G_2)}{2} - \frac{F(G_2)}{4} + n_1 n_2 (n_1 n_2 + n_1 - 1) \\ &+ n_1 m_2 (n_1 n_2 + n_1 - 2) - \frac{n_1 M_1(G_2)}{4} \\ &= \frac{(n_1 n_2 + n_1)^2}{4} (m_1 + m_2) + n_1 n_2 (n_1 n_2 + n_1 - 1) \\ &+ n_1 m_2 (n_1 n_2 + n_1 - 2) - \frac{(n_2 + 1)^2}{4} \sum_{uv \in E_1} (n_u^2(e|G_1) + n_v^2(e|G_1)) \\ &+ \frac{2(n_2 + 1)^2 Sz(G_1) + 2M_2(G_2) - n_1 M_1(G_2) - F(G_2)}{4}. \end{aligned}$$

Corollary 4. The following equalities are hold:

1. $Sz^*(P_n o P_m) = \frac{2n^3 m^2 + 3n^2 m^3 + 24n^2 m^2 + 4n^3 m - 2nm^2 + 2n^3 + 15mn^2 - 15n^2 + 40n - 52nm - 6}{4}$.
2. $Sz^*(P_n o C_m) = \frac{2n^3 m^2 + 2n^2 m^3 + 28n^2 m^2 + 4n^3 m - 2nm^2 + 2n^3 + 26mn^2 - 2n - 52nm}{4}$.
3. $Sz^*(C_n o P_m) = \frac{n^3 m^2 + n^2 m^3 + 9n^2 m^2 + 2n^3 m + n^3 + 3mn^2 - 5n^2 - 16nm - 2n - 2}{4}$.
4. $Sz^*(C_n o C_m) = \frac{n^3 m^2 + n^2 m^3 + 10n^2 m^2 + 2n^3 m + n^3 + 9mn^2 - 16nm}{4}$.

REFERENCES

1. A. R. Ashrafi, T. Došlić and A. Hamzeh, The Zagreb coindices of graph operations, *Discrete Appl. Math.* **158** (2010) 1571–1578.
2. M. Azari and A. Iranmanesh, Some inequalities for the multiplicative sum Zagreb index of graph operations, *J. Math. Inequal.* **9** (2015) 727–738.
3. K. C. Das and I. Gutman, The first Zagreb index 30 years after, *MATCH Commun. Math. Comput. Chem.* **50** (2004) 83–92.
4. N. De, S. M. Abu Nayeem and A. Pal, F-index of some graph operations, *Discrete Math. Algorithm. Appl.* **8** (2016) 1650025.
5. B. Furtula and I. Gutman, A forgotten topological index, *J. Math. Chem.* **53** (2015) 1184–1190.
6. I. Gutman, A formula for the Wiener number of trees and its extension to graphs containing cycles, *Graph Theory Notes New York* **27** (1994) 9–15.

7. I. Gutman and N. Trinajstić, Graph theory and molecular orbitals, Total π -electron energy of alternant hydrocarbons, *Chem. Phys. Lett.* **17** (1972) 535–538.
8. M. H. Khalifeh, H. Yousefi-Azari and A. R. Ashrafi, The first and second Zagreb indices of some graph operations, *Discrete Appl. Math.* **157** (2009) 804–811.
9. S. Klavžar, A. Rajapakse and I. Gutman, The Szeged and the Wiener index of graphs, *Appl. Math. Lett.* **9** (1996) 45–49.
10. T. Mansour and M. Schork, The vertex PI and Szeged index of bridge and chain graphs, *Discrete Appl. Math.* **157** (2009) 1600–1606.
11. S. Nagarajan, K. Pattabiraman and M. Chendra Sekharan, Revised Szeged index of product graphs, *Gen. Math. Notes* **23** (2014) 71–78.
12. Z. Yarahmadi and A. R. Ashrafi, The Szeged, vertex PI, first and second Zagreb indices of corona product of graphs, *Filomat* **26** (2012) 467–472.
13. M. Randić, On generalization of Wiener index for cyclic structures, *Acta Chim. Slov.* **49** (2002) 483–496.
14. D. B. West, *Introduction to Graph Theory*, Prentice–Hall, Inc., Upper Saddle River, New Jersey, 2000.

A Note on Revised Szeged Index of Graph Operations

NASRIN DEHGARDI

Department of Mathematics and Computer Science, Sirjan University of Technology,
Sirjan, I. R. Iran

یادداشتی در مورد شاخص سگد بازنگری شده از عملگرهای گراف

ادیتور رابطا : مسن یوسفی آذری

چکیده

فرض کنید G گرافی متناهی و ساده با مجموعه یال‌های $E(G)$ باشد. شاخص سگد بازنگری شده به صورت زیر تعریف می‌شود:

$$Sz^*(G) = \sum_{e=uv \in E(G)} \left(n_u(e|G) + \frac{n_G(e)}{2} \right) \left(n_v(e|G) + \frac{n_G(e)}{2} \right)$$

که $n_u(e|G)$ تعداد رئوس در G است که به u نزدیکتر از v هستند و $n_G(e)$ تعداد رئوس هم‌فاصله از e در G است. در این مقاله، شاخص سگد بازنگری شده را برای عملگرهای اتصال و ضرب کرونا در گراف‌ها بررسی می‌کنیم.

لغات کلیدی: شاخص توپولوژیک، شاخص سگد بازنگری شده، عملگر گراف