

International Journal of Molecular and Clinical Microbiology



In Vitro Activity of *Mentha longifolia* Leaves and *Pimpinella anisum* Seeds Against a Clinical Strain of *Trichomonas Vaginalis*

Maryam Niyyati^{1*}; Zaynab Joneidi²; Mohammad KamaliNejad³; Ali Haghighi²; Alireza Abadi⁴, Zahra Arab-Mazar¹, Soudabeh Heidari²

1. Infectious Diseases and Tropical Medicine Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

2. Department of Medical Parasitology and Mycology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Dep. Pharmacognosy, School of Pharmacy, Shahid Beheshti University of Medical Sciences, Tehran, Iran
Dept. of Health and Community Medicine, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

_____ABSTRACT

ARTICLE INFO Article history: Received 31 April 2015 Accepted 19 May 2015 Available online 1 June 2015 Keywords: Mentha longifolia, Pimpinella anisum, in vitro, Trichomonas vaginalis, STD

Trichomoniasis is one of the most common urinary-genital tract disease. The causal agent of trichomoniasis is flagellate protozoan called Trichomonas vaginalis. Metronidazole and tinidazole are the drug of choice for trichomoniasis treatment. However, carcinogenic and embryo-teratogenic effects of the mentioned drugs and resistance to metronidazole has been reported. According to cytotoxic effects of medicinal herbs including Mentha longifolia and Pimpinella anisum on microbial organisms and their traditional use in sexually transmitted diseases the in vitro activity of the Mentha longifolia and Pimpinella anisum against a clinical strain of Trichomonas vaginalis was investigated. Concentrations of aqueous extract of Mentha longifolia leave (100-250 µg/ml) and Pimpinella anisum seed (100-500 µg/ml) were obtained. Trichomonas vaginalis isolated from patient with urogenital complications was cultivated in TYI-S-33 medium. After treatment of T. vaginalis (1×10^6) with various concentrations of extracts, the minimum inhibitory concentration (MIC) was calculated and data were analyzed by repeated measures using SPSS software. Aqueous extract of Mentha longifolia decreased the average number of live parasites, in accordance of the time and concentration manner. The MIC of *Mentha longifolia* at 24 and 48 h were 250 and 200 µg/ml, respectively. The percent of growth inhibition under various concentration of Pimpinella anisum showed only 73.7% and 86.9% at 500 µg/ml concentration after 24 and 48 hours, respectively. The present study demonstrate the higher anti-Trichomonas activity of Mentha longifolia in comparison to Pimpinella anisum. There is a clear need to isolate effective fragments of Mentha longifolia in future and expand the data for in vivo evaluation including cell culture and animal models study.

1. Introduction

Trichomonas vaginalis is a parasitic protozoan with a predilection for human urogenital tract and causative agent for vaginitis, cervicitis and urethritis in females (Garcia, 2006). There are serious health consequences in trichomoniasis such as adverse pregnancy outcome, increased risk of infertility and

cervical carcinoma (Gutierrez, 2000). Interestingly, there are several studies indicating the association of trichomoniasis infection in men and benign hyperplasia and prostate cancer (Petrin et al., 1998; Rezaeian et al., 2009).

The prevalence of trichomoniasis in Iran has a range from 0.1 to 15% due to various methods of detection and different examined population (Rezaeian et al., 2009). It should be mentioned

^{*}Corresponding author: Dr.Maryam Niyyati

Tel: 0912-2407432

E-mail address: maryamniyati@sbmu.ac.ir, maryamniyati@yahoo.com

there are many unreported cases of trichomoniasis in the region, because of chronic pattern and mild symptoms in over 50% of women (Rezaeian et al., 2009). The drug of 5-nitroimidazole choice for therapy is including metronidazole compounds and tinidazole (Ryu and Min, 2006). In Iran the only 5-nitroimidazole available drug is metronidazole, however the main drawback to metronidazole is wide adverse effects including nausea, gastrointestinal disturbance, leukopenia and headache (Sobel and Sobel, 2015). Besides, there are evidences of emerging resistance to metronidazole as Center for Disease control (CDC) reported that 6% of Trichomonas vaginalis isolated from patients showed resistant to metronidazole (Sena et al., 2014).

There were reports regarding teratogenic and carcinogenic effects of metronidazole. To this end, natural products derived from medicinal plants are a priority objectives (Sobel et al., 1999). Accordingly, the world health organization (WHO) estimates that approximately 80% of the world population mainly use plant based remedies for various diseases. Indeed, the potential of natural products such as *Mentha longifolia* (Lamiaceae) and Pimpinella anisum in the treatment of diseases especially sexually transmitted disease is evident (Kose et al., 2015; Palmeira-de-Oliveira et al., 2013; Saeidi et al., 2014). Noteworthy, in Iran medicinal plants are an attractive approaches as an alternative options to the chemical drugs in various diseases (Bakht et al., Shaheen and Shafi, 2014: Kose et al., 2015: et al., 2014). Mentha longifolia Saeidi (Lamiaceae) and Pimpinella anisum has been used in medicine due to their cytotoxic effect in bacteria and fungi (Bakht et al., 2014; Eissa et al., 2014; Kose et al., 2015). Pimpinella anisum (anise) belonging to Umbelliferae family is an endemic plant which has been used in Iranian traditional medicine, anise showed antibacterial effects on various microorganisms (Ghoshegir et al., 2014). It should be mention that there were no previous reports of anti-trichomonas activity of Mentha longifolia and Pimpinella anisum in Iran and around the world.

Considering the adverse effect of metronidazole, the present study was aimed to determine the effect of *Mentha longifolia* and *Pimpinella anisum* on *Trichomonas vaginalis*

isolated from a female patient with urogenital complication.

2. Materials and Methods

2.1. Parasite and cultivation in TYI-S-33

Trichomonas vaginalis parasite was isolated from vaginal discharge of women with symptomatic trichomoniasis (Heidari et al. 2013). Confirmation regarding the presence of T. vaginalis was done using direct wet mount and culture. T. vaginalis were cultured axenically in the Tripticase (Sigma-Aldrich, Germany), Yeast extract (Merck, Germany), iron-serum (TYI-S-33) medium for 24-48 h according to previous studies (Diamond et al., 1995). The medium was supplemented with 10% heat inactivated bovine serum, vitamins (Merck, Germany) and streptomycin (Merck, Germany) and incubated at 37°C (Diamond et al., 1995). Organisms in the logarithmic phase of growth were harvested, centrifuged and suspended on fresh medium for experiments. The criteria used for viability was eosin tests. An initial inoculum of 1×10^6 trophozoites/ml was incubated in the presence of the extracts.

2.2. Preparation of plant extracts

Mentha longifolia leaves and *Pimpinella anisum* seeds were collected from the Ray area of Iran and identified and approved by a local herbalist (Mohammad Kamalinejad) from the School of Pharmacology, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Plants were chosen regarding their traditional application in the treatment of venereal disease (Furst and Zundorf, 2015). Aqueous extracts of two endemic plant including *Mentha longifolia* leaves and *Pimpinella anisum* seeds were obtained using maceration technique as previously described (Rogers et al., 2014; Shreaz et al., 2011). Briefly, plants were air dried, powdered and suspended in sterile distilled water for several hours. The extracts were centrifuged and the supernatants filtered through 0.22 filters. The concentration range of the extracts were from 100-250 µg/ml for *Mentha longifolia* and 100-500 for *Pimpinella anisum*.

2.3. Anti-Trichomonas activity of the plants

On thousands microliter of trichomonads with the same volume of extracts were thoroughly mixed and the micro tubes kept at 37° C for 24 and 48 hours. Eosin dye was used to determine the extract effectiveness. Same volume of eosin 0.1% and parasite suspensions was then mixed. Positive and negative controls were metronidazole (0.1 µg/ml) and culture medium, respectively. Experiments were performed in triplicate.

2.4. Determination of the Minimum Inhibition Concentration (MIC) of the plant extracts

Results were recorded as the percentage of living organisms using exclusion test by eosin (0.1%). The percent of growth reduction were calculated for each concentration. The minimum inhibitory concentration (MIC) was defined as the lowest concentration of plant extract in which no lived parasite was detected.

2.5. Statistical analysis

The data were analyzed using repeated measures of SPSS. The P value was set at 0.05 for significance level.

3. Results

The results of the present research are presented as the percentage of growth reduction of Trichomonas vaginalis by comparison of the number of parasites found in the tests to the number of parasites in control. The present research, showed the effectiveness of the extract Mentha longifolia by remarkable of trichomonocidal effect (Fig 1), since the aqueous extract obtained from Mentha longifolia leaves totally abolished the trophozoite viability. This extract was active in a dose dependent manner. The MIC of Mentha longifolia at 24 h and 48 h were 250 and 200 µg/ml, respectively (Fig 2, Table 1). However, it should be mention that the 100-250 concentrations µg/ml failed to completely destroy the parasites (Table 1). The concentration 150 g/ml, 8.4×10^4 viable T. vaginalis were detected at the end of the incubation period. Viable trophozoites within the control media did not show any relevant changes during the test. Average number of viable trophozoites revealed increase with the

decreasing concentrations of the aqueous extract of *Mentha longifolia*.

However, the percent of growth inhibition under various concentration of *Pimpinella anisum* showed only 73.7% and 86.9% at 500 μ g/ml concentration after 24 and 48 hours, respectively. The effect of *Pimpinella anisum* on the viability of *T. vaginalis* is shown in Table 2 and fig 3.

4. Discussion

Trichomoniasis is considered as a neglected disease mainly due to poor understanding of pathogenesis factors. According to WHO report, trichomoniasis affects approximately 174 million new cases annually (Arab-Mazar and Niyyati, 2015; Rogers et al., 2014; Sena et al., 2014). In Iran, the only available 5nitroimidazole drug is metronidazole, however there are several reports in this region regarding resistance. Due to adverse effect of chemical drugs such as metronidazole, use of plant- based compounds could be an alternative approach for treatment of Trichomonasiasis. In Iran traditional healers prescribe maceration of the leaves and seeds of lamiacea family specially Mentha longifolia and Pimpinella seeds to be taken orally for treatment of urogenital infectious organisms (Kose et al., 2015). The traditional use of the leaves of Mentha longifolia was mainly for recovery from several diseases including sexually transmitted diseases such as bacterial vaginitis. The present study revealed that *M. longifolia* could be an appropriate candidate for further phytochemical analysis as percent of growth inhibition was %100 at 200 and 250 µg/ml after 24 and 48 hours, respectively. This is important because it showed an efficient capacity in destroying the parasites at low concentrations. Previous researchers demonstrate 1,8 cineole as the most important component of M. longifolia (LaGow, 2004). Interestingly a recent research on Eucalyptus leaves revealed a high anti-Trichomonas activity. These researchers highlighted 1,8 cineole as a vital component of Eucalyptus for abolishing trophozoites viability. Indeed, 1,8 cineole is a common component between both plants and it may lead to high anti-trichomonas activity (Robles-Zepeda et al., 2011). It should be mention that there were no previous research regarding trichomonicidal effect of *M. longifolia* in Iran and worldwide. Other researchers reported the effectiveness of several plants on *Trichomonas* including *Stachys*, however it was not able to destroy *Trichomonas* after 72 hours (Youse et al., 2012). On the other hand, there are some promising reports regarding anti-*trichomonas* activity of herbal medicines (Gamberini et al., 2015; Kaur and Arora, 2009; Robles-Zepeda et al., 2011). Further analysis is highly recommended with shorter incubation periods and analysis of active component of *M. longifolia*.

Pimpinella anisum (Anis) is an endemic growing plant in Iran and it is widely used for gastrointestinal disturbance. However there are reports regarding its cytotoxic effect to microbial population including *Streptococcus* and *Bacillus* (Robles-Zepeda et al., 2011).

It is worthy to mention the main antimicrobial component is Trans-anethole. Indeed previous researchers reported that transanethole could lead to death of various microorganisms (Gamberini et al., 2015; Kaur and Arora, 2009). The present research revealed the decline of parasite average number by increasing the concentration, however the parasite decrease percent was not significant and thus the mentioned extract may not be as effective as *M. longifolia*. The highest percent of growth reduction of the mentioned extract was 86% at 500 μ l/ml concentration after 48 hours. It should be mention that aqueous extracts were examined, since traditional healers mainly prepare their remedies as infusion or maceration.

Altogether, there is a clear need for further analysis regarding isolation of active component of *M. longifolia* extract, cytotoxicity effect and in vivo tests should be in priority for further researchers.

Acknowledgments

Thanks are due to Dr. Nozhat Zebardast for her valuable comments. Dr. Maryam Niyyati has been supported by Infectious Diseases and Tropical Medicine Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran (No: 422/176).



Figure 1. Eosin stained smear showing lethal effect of Mentha longifolia on T. vaginali

	24h				48h	D Value		
Dosage of treatment µg/ml	Average Count of	percentage	Standard	Average Count of	percentage	Standard deviation	P-value	
	Parasite in 1 ml*	of growth reduction	deviation	Parasite in 1 ml*	of growth reduction		Concentration effect	Interaction of time- concentration
100	53.65		13.7	15.6		3.1	<0/0011	<0/0011
150	40.55	24.4	10.1	8.4	46.1	1.5	$< 0/001^{2}$	$< 0/001^{2}$
200	12.8	68.4	2.1	0	100	0	<0/001 ³	<0/0013
250	0	100	0	0	0	0		
Negative control	552.1		31.8	872.1		65.9		
Interaction of time-concentration								<0/001

 1 . Comparison of 100 and 150 μg / ml concentration through the analysis of repeated measures 2 . Comparison 150 and 200 μ g / ml concentration through the analysis of repeated measures

³ Comparison 200 and 250 μ g / ml concentration through the analysis of repeated measures * Data: $\times 10^4$



Figure 2. Percentage of growth inhibition of Trichomonas vaginalis at 24 and 48 h after exposure to different concentration of Mentha longifolia seed extract

	24h				48h	P-Value			
Dosage of treatment µg/ml	Average Count of	percentage	Standard	Average Count of	percentage	Standard deviation			
	Parasite in 1 ml*	of growth reduction	deviation	Parasite in 1 ml [*]	of growth reduction		Concentration effect	Interaction of time- concentration	
100	315.4		25.8	153.1		4.7	<0/0011	<0/0011	
200	183.6	41.7	7.3	135.4	11.5	3.2	<0/001 ²	<0/001 ²	
300	153.7	16.2	2.4	125.7	7.7	12.5	0.005^{3}	0.005^{3}	
400	150.9	1.8	3.3	118.3	5.8	4	0.011^4	0.011^{4}	
500	145	3.9	10	114	3.6	7.7			
Negative control	552.1		31.8	872.1		65.9			
Interaction of time-Density								<0/001	

Table 2. Effect of aqueous extract of Pimpinella anisum on the in vitro growth of T. vaginalis for Different incubation periods

1. Comparison of 100 and 200 μ g / ml concentration through the analysis of repeated measures

2. Comparison of 200 and 300 μ g / ml concentration through the analysis of repeated measures

3. Comparison of 300 and 400 μ g / ml concentration through the analysis of repeated measures

4. Comparison of 400 and 500 μ g / ml concentration through the analysis of repeated measures

* Data $\times 104$



Figure 3. Percentage of growth inhibition of *Trichomonas vaginalis* at 24 and 48 h after exposure to different concentration of *Pimpinella anisum* seed extract

Refereces

- Arab-Mazar, Z., Niyyati, M. 2015. Trichomonas vaginalis Pathogenesis: a Narrative Review. Novelty in Biomedicine. 3(3); 148-154.
- Bakht, J., Shaheen, S., Shafi, M. 2014. Antimicrobial potentials of Mentha longifolia by disc diffusion method. Pak J Pharm Sci. 27(4); 939-945.
- Diamond, L.S., Clark, C.G., Cunnick, C.C. 1995. YI-S, a casein-free medium for axenic cultivation of Entamoeba histolytica, related Entamoeba, Giardia intestinalis and Trichomonas vaginalis. J Eukaryot Microbiol. 42(3); 277-278.
- Eissa, T.F., Gonzalez-Burgos, E., Carretero, M.E., Gomez-Serranillos, M.P. 2014. Compositional analysis and in vitro protective activity against oxidative stress of essential oils from egyptian

plants used in traditional medicine. Nat Prod Commun. 9(9); 1377-1382.

- Furst, R., Zundorf, I. 2015. Evidence-Based Phytotherapy in Europe: Where Do We Stand? Planta Med. doi: 10.1055/s-0035-1545948
- Gamberini, M.T., Rodrigues, D.S., Rodrigues, D., Pontes, V.B. 2015. Effects of the aqueous extract of Pimpinella anisum L. seeds on exploratory activity and emotional behavior in rats using the open field and elevated plus maze tests. J Ethnopharmacol. doi: 10.1016/j.jep.2015.03.053
- Garcia, L.S., 2006. Diagnostic medical parasitology: American Society for Microbiology Press.
- Ghoshegir, S.A., Mazaheri, M., Ghannadi, A., Feizi, A., Babaeian, M., Tanhaee, M., Adibi, P. 2014. Pimpinella anisum in modifying the quality of life in patients with functional dyspepsia: A doubleblind randomized clinical trial. J Res Med Sci. 19(12); 1118-1123.
- Gutierrez, Y., 2000. Diagnostic pathology of parasitic infections with clinical correlations: Oxford University Press.
- Heidari, S., Bandehpour, M., Seyyed-Tabaei, SJ., Valadkhani, Z., Haghighi, A., Abadi., Kazemi, B., 2013. Ferredoxin Gene Mutation in Iranian Trichomonas vaginalis Isolates. <u>Ir</u>an J of Parasitol. 8(3); 402-7.
- Kaur, G.J., Arora, D.S., 2009. Antibacterial and phytochemical screening of Anethum graveolens, Foeniculum vulgare and Trachyspermum ammi. BMC Complement Altern Med. 9,30; doi: 10.1186/1472-6882-9-30
- Kose, L.S., Moteetee, A., Van Vuuren, S., 2015. Ethnobotanical survey of medicinal plants used in the Maseru district of Lesotho. J Ethnopharmacol. doi: 10.1016/j.jep.2015.04.047
- LaGow, B., 2004. PDR for herbal medicines: Thomson PDR.
- Nourollahpour Shiadeh, M., Niyyati, M., Fallahi, S., Rostami, A., 2015. Human parasitic protozoan infection to infertility: a systematic review. Parasitol Res. 2015 Nov 16.
- Palmeira-de-Oliveira, A., Silva, B.M., Palmeira-de-Oliveira, R., Martinez-de-Oliveira, J., Salgueiro, L. 2013. Are plant extracts a potential therapeutic approach for genital infections? Curr Med Chem, 20(23), 2914-2928.
- Petrin, D., Delgaty, K., Bhatt, R., Garber, G. 1998. Clinical and microbiological aspects of Trichomonas vaginalis. Clin Microbiol Rev. 11(2); 300-317.

- Rezaeian, M., Vatanshenassan, M., Rezaie, S., Mohebali, M., Niromand, N., Niyyati, M., Babaei, Z., 2009. Prevalence of Trichomonas vaginalis using parasitological methods in Tehran. Iranian ournal of Parasitology. 4(4); 43-47.
- Robles-Zepeda, R.E., Velazquez-Contreras, C.A., Garibay-Escobar, A., Galvez-Ruiz, J.C., Ruiz-Bustos, E., 2011. Antimicrobial activity of Northwestern Mexican plants against Helicobacter pylori. J Med Food. 14(10); 1280-1283.
- Rogers, S.M., Turner, C.F., Hobbs, M., Miller, W.C., Tan, S., Roman, A.M., Erbelding, E., 2014. Epidemiology of undiagnosed trichomoniasis in a probability sample of urban young adults. PLoS One. 9(3); e90548.
- Ryu, J.S., Min, D.Y., 2006. Trichomonas vaginalis and trichomoniasis in the Republic of Korea. Korean J Parasitol. 44(2); 101-116.
- Saeidi, S., Hassanpour, K., Ghamgosha, M., Heiat, M., Taheri, R.A., Mirhosseini, A., Farnoosh, G., 2014. Antibacterial activity of ethyl acetate and aqueous extracts of Mentha longifolia L. and hydroalcoholic extract of Zataria multiflora Boiss. plants against important human pathogens. Asian Pac J Trop Med, 7S1, S186-189. doi: 10.1016/s1995-7645(14)60229-7
- Sena, A. C., Bachmann, L. H., & Hobbs, M. M. (2014). Persistent and recurrent Trichomonas vaginalis infections: epidemiology, treatment and management considerations. Expert Rev Anti Infect Ther. 12(6); 673-685.
- Shreaz, S., Bhatia, R., Khan, N., Muralidhar, S., Basir, S.F., Manzoor, N., Khan, L.A., 2011. Exposure of Candida to p-anisaldehyde inhibits its growth and ergosterol biosynthesis. J Gen Appl Microbiol. 57(3); 129-136.
- Sobel, J.D., Nagappan, V., Nyirjesy, P. 1999. Metronidazole-resistant vaginal trichomoniasis an emerging problem. N Engl J Med, 341(4), 292-293.
- Sobel, R., Sobel, J.D., 2015. Metronidazole for the treatment of vaginal infections. Expert Opin Pharmacother. 16 (7); 1109-1115.
- Youse, H.A., Kazemian, A., Sereshti, M., Rahmanikhoh, E., Ahmadinia, E., Rafaian, M., Darani, H.Y. 2012. Effect of Echinophora platyloba, Stachys lavandulifolia, and Eucalyptus camaldulensis plants on Trichomonas vaginalis growth in vitro. Adv Biomed Res. 1, 79. doi: 10.4103/2277-9175.102987.