

Study on Some Chemical Compounds on the Vase Life of Two Cultivars of Cut Roses

S. Mohammadi Ostad Kalayeh

Graduated Student of Islamic Azad University.

Science and Research Branch,

Tehran

Iran

Y. Mostofi,

Associated Prof.

University of Tehran.

Karaj,

Iran

M. Basirat

Scientific member of Soil and Water Research Center

Tehran

Iran

Received: 15 February 2011 Accepted: 8 May 2011

*Corresponding author email: mohamadisona@gmail.com

Rose (*Rosa hybrida* L.) is one of the mostly cultivated cut flowers in Iran and around the world. The study was carried out to increase the vase life and quality of cut rose 'Grand Prix' and 'Avalanche' by using various floral preservative solutions. Floral preservative solutions were AgNO₃, 8-HQS, nano silver and distilled water. Treatments were compared with tap water as control. The experiment was arranged in a randomized complete block design. The effects of applied treatments on longevity and quality of cut rose flowers were evaluated by using the percentage of opening the flower, flower diameter, the vase life and relative fresh weight. Results showed that the flowers treated with 8-HQS (250 mg l⁻¹) + nano silver (2mg l⁻¹) had the highest flower diameter and the percentage of opening the flower among treatments. Maximum vase life (14.33 d) of cut rose obtained with (nano silver 2mg l⁻¹), and minimum (6.54 d) with AgNO₃. The vase life of control was 5.79 d.

Abstract

Keywords: Distilled water, Nano silver, Rose, Silver nitrate, 8-hydroxy quinoline sulphate.

INTRODUCTION

Rose (*Rosa sp.*) is belong to Rosaceae family which identified as the highest demand flower in world (Khalighi & Shfie, 1994). One problem in our country is postharvest waste of horticultural products, especially ornamental plants. Rose has short vase life. The most important way to effects of ethylene on horticultural products, especially cut flowers, is to use of chemical materials for blocking activity or synthesis of ethylene. This materials included that carbohydrates, bactericides, biocides, acidosis and ethylene inhibitors (Ebrahimzadeh and Seifi, 1999). Application of some compound such as sucrose or 8-hydroxy quinoline (8-HQ) is effective to increase vase life of cut roses (Michalcznk *et al.*, 1989).

Among the preservative solutions which are used to increase cut flowers longevity, 8-HQ is more effective to control microbial agents and mostly uses in many of cut flowers (Kader, 2002; Reddy *et al.*, 1995; Khalighi and Shafiei, 1994 and Hussein, 1994). Some part of suitable effect of 8-HQ had been attributed to the water balance and closing of stomata.

Also, 8-HQS and 8-HQC, cause to acidify of water and using them continuously, had been increased vase life of cut roses remarkably, and when it used with sucrose, because of facilitating in transfer water through vessel elements, cause to increase sugar in petals and to increase flower diameter (Ichimura and Sotu, 1999). It had been reported that using 8-HQS in cut *Lisianthus* improves flower vase life (Ichimura and Korenaga, 1998). In a research, Reddy *et al.*, (1995), used mixed compounds such as citric acid, 8-HQS and sucrose. 8-HQS and %4 sucrose caused to increase flower longevity by 16 days and improve water retention and water uptake by flowers.

Two common compounds containing silver which are used for cut flowers are AgNO_3 and STS (Azadi *et al.*, 2001 and Fahmy *et al.*, 2005). AgNO_3 is relatively unmobile in plant and acts as an antimicrobial and ethylene blocker factor in plants (Bartoli *et al.*, 1996). Effects of AgNO_3 was studied on vase life and bud opening in cut orchids (*Dendrobium pompadour*), results showed that AgNO_3 is better than STS in controlling of microbial growth and increasing in bud opening and vase life (Ketsa, 2000). Today, several chemical compounds were introduced for preservation of plant against pathogens and increasing of vase life, but some of these compounds have side effects for environment. Silver nano particles are good and safe antibacterial tool for ornamental and horticultural products.

Main aim of this study is research on effects of different chemical compounds such as AgNO_3 , 8- hydroxy quinoline sulphate (8-HQS), SNP and distilled water for increasing vases life of cut roses 'Grand prix' and Avalanche'.

MATERIALS AND METHODS

Cut roses 'Avalanche' and 'Grand prix' harvested from a greenhouse around Takestan in commercial stage and after pre-cooling in temperature 5 °C for 4 hours, transferred to laboratory. All of the flowers harvested in optimum physiological stage. The flowers recutted under the water and the height of all flower branches set on 55 cm.

The leaves and blades of lower part of branches eliminated and only 3 leaves over the stem protected. Then the concerned treatments performed on them (Table 1). Environmental conditions in vase life room was: temperature 19-21°C, photoperiod 9 hr, light intensity 15-20 $\mu\text{mol s}^{-1}\text{m}^{-2}$ using white fluorescent lamp and 60% relative humidity.

Experimental Design and Treatments

This experiment had been performed in RCBD on two cultivar of cut roses 'Avalanche' and 'Grand prix' with 11 different chemical treatments in three replications, each plot was include 4 cut flowers. Chemical treatment in this experiment is as follow (Table 1).

Measurement of Characteristics

Relative fresh weight, fresh weight to dry weight ratio, flower diameter, flower opening and vase life were measured. Vase life finished with some symptoms such as necrosis, wilting and abscission of petals, chlorosis and abscission of leaves, bent neck and outrolling of petals which cause to reduce flower attraction and marketing.

Every 4 days, flower diameter measured by caliper and record in each flower, separately.

Relative fresh weight measured every 2 days. For this purpose, flower branch was weighted and its relative fresh weight calculated by using the following formula (Bartoli *et al.*, 1996; Setyadjit *et al.*, 2004).

$$RFW = W_t / W_{t=0} \times 100$$

$$W_t = \text{F.W in } 2, 4, 6^{\text{th}}, \dots \text{day.}$$

$$W_{t=0} = \text{F.W in zero day.}$$

For measuring of dry weight to fresh weight ratio, fresh weight was measured every 2 days and after the end vase life, all of the cut flowers located in oven 60 °C for 72 hrs.

Data Analysis

Data were analysed with MSTATC software. Mean comparison carried out with DNMRT.

RESULTS AND DISCUSSIONS

In Avalanche cultivar, 2 and 1 mg l⁻¹ SNP (T₄, T₅) and distilled water (T₁₀) have the highest vase life, 18.17, 17, 5 and 16, 58 days, respectively. These treatments had significant difference with other treatments. Also the least vase life was showed in AgNO₃ and control. In Grand prix cultivar, T₆ and T₈ had highest vase life and the least average in this cultivar is related to control treatment (Table 2). Increasing vase life in T₄ and T₅ in Avalanch cultivar can attribute to the effect of SNP in various levels and also it can refer to in water quality in T₁₀ treatment (table 3). The existence of SNP inhibits growth of microorganisms. These results were agreed with Kim *et al.*, (2005).

In AgNO₃ treatment (T₁), wilting and bent neck observed in early days. Water stress symptoms (wilting) and bent neck are explainable in T₁, AgNO₃ banded to organic compounds and cannot moves upward in vessels (Nowak and Rudnicki, 1990). Also, susceptibility of vessels of herbaceous plants and toxicity of Ag⁺ as a heavy metal, are two important reasons for this effect (Alvarez *et al.*, 1994). It seems that Ag⁺ damages to vessels and inhibits water transfer in xylem and flowers will wilt. The results of this research about the effect of AgNO₃ is not agreed with Khondakar and Mazumdar (1985), because they had been obtained the best results with sucrose 3% + 8-HQC 3% + AgNO₃ %1, while the results of current research was similar to Alvarez *et al.*, (1994), which in both two research, undesirable effect of AgNO₃ on cut rose and tuberoses had been observed. Alvarez *et al.*, (1994) used 600 mg l⁻¹ AgNO₃, while Khondakar and Mazumdar (1985) applied less than 1% and observed that this concentration had negative effect. It seems that 8-HQ eliminated the negative effect of AgNO₃. Our results were agreed with the results of Jowkar and Salehi (2006) but it was not agreed with Lal (1993).

If distilled water has high quality cut flowers have more vase life. Halvey (1976) showed that distilled water is suitable for preservative solution. In the present research, distilled water was one of the best treatment to increasing vase life of cut Rose 'Avalanche'.

According to ANOVA, simple effect of time, cultivar and treatment on flower diameter and the interaction between treatment and were significant (P ≤ 0.01).

Treatments T₆, T₈ and T₉ had the highest average of flower diameter, but their difference with T₃, T₄, T₅ and T₇ was not significant. Lowest flower diameter was for control plants.

In rose 'Grand Prix' effect of interaction between chemical compounds and cultivars on flower diameter was significant and T₈ had the highest average of flower diameter. T₃, T₄, T₅, T₆,

T₇, T₈ and T₉ had not significant differences with each other. Also in cut roses 'Avalanche' distilled water with the average of 105.7 mm had the highest flower diameter which had not significant differences with T₆ treatment.

This results had agreed with the results of Hussein (1994), Fakhrae and Maidani (2004) and Kalatejari (2009).

Literature Cited

- Alvarez, V. N., Colinas, M. T. and Villanueva, V. C. 1994. Postharvest uses of chemicals compounds to increase vase life of cut tuberose flowers. Hort. Sci. 1: 15-20.
- Azadi, P., Edrisi, B., Banijamali, S. M., Bayat, H., Shafiei, M.R., Mirzakhani, A. and Hashemabadi, D. 2001. Carnation, (1st Special Bulletin of National Research Center of Ornamental Plants). Mahallat. Iran.
- Bartoli, C., Guimet, G. and Montaidi, S. 1996. Ethylene production and response to exogenous ethylene in senescing petals of *Chrysanthemum morifolium*. Plant Science, 124: 15-21.
- Ebrahimzadeh, A. and Seifi, Y. 1999. Storage and transportation of cut flowers, green ornamental plants. Akhtar publication.
- Fahmy, A. E. L. Rahman, and S. Hassan. 2005. Postharvest studies on some important flower crops. www. Lib. Uni-Corvinus. Hu/PhD/Sadeck-hassan.
- Fakhrayi, M. and Meidani, J. 2003. Effect of sucrose with 8-hydroxy quinoline citrate on the quality of cut Rose "First Red" on the vase life and postharvest in different situation. National Congress of cut flowers of Iran. Tehran.
- Halvey, A. H. 1976. Treatments to improve water balance of cut flowers. Acta Hort. 64: 223-236.
- Hussein, H.A.A. 1994. Varietal responses of cut flowers to different antimicrobial agents of bacterial contamination and keeping quality. Acta Hort. 368: 106-116.
- Ichimura, K. and Korenaga, M. 1998. Improvement of vase life and petal color expression in several cultivars of cut *Eustoma* flowers using sucrose with 8-hydroxy quinoline sulfate. Bull. Natl. Res. Veg. Ornam. Plant and Tea, Japan. 13: 31-39.
- Ichimura, K. and Sotu, K. 1999. Effects of the time of sucrose treatment of vase life, soluble carbohydrate concentrations and ethylene production in cut sweet pea flowers. Plant Growth Regul. 28: 117-123.
- Jowkar, M., and Salehi, M.H. 2006. Effect of preservative compounds on the vase life of cut. *Polianthus*.
- Kader, A. A. 2002. Postharvest technology of horticulture crops. Third edition. University of California. Publication 3311 .
- Kalateh jari, S. 2008. Increasing the qualificational characters and vase life with some chemical treatments. PhD thesis. Islamic Azad University, Tehran. Iran.
- Ketsa, S. 2000. The effect of AgNO₃ on the vase life and bud opening of cut flower of *Dendrobium "Pompadour"*. Hort. Rev., 18: 1-85.
- Khalighi, A., and Shafiei, M. 2000. Study on effect of chemical, treatments and stage stage of harvest on vase life and some qualificational characters of cut Carnation. Iran agricultural Sciences. 31: 119-125.
- Khondakar, S.R.K. and Mazumdar, B.C. 1985. Studies on prolonging the vase life of tuberose cut flowers. South Indian Hort. 33 (2): 145-147.
- Kim, J. H., Lee, A. K. and Suh, J. K. 2005. Effect of certain pre-treatment substances on vase life and physiological character in *Lilium* spp. Acta Hort , 373: 22-30.
- Lal, S. D. 1993. Effect of chemical substances on the vase life and quality of gladiolus 'Silver Horn'. Progressive Horticulture. 22(1-4): 63-68.
- Michalczuk, B.D. Goszczynska, M. Rudnicki, R.M. and Halevy, A. H. 1989. Calcium promotes longevity and bud opening in cut rose flowers. Israel Journal of Botany. 38: 209-215.
- Nowak, J. and Rundicki, R. M. 1990. Postharvest handling and storage of cut flowers, florist

- green, and pot plants. Timber Press INC. USA . P: 199.
- Reddy, B. S., Singh, K. and Singh, A. 1995. Effect of sucrose, citric acid and hydroxyquinoline sulfate on postharvest physiology of tuberose 'single'. *Advances in Agricultural Research in India*. 3: 161-167.
- Setyadjit, J.D.C., Irving, D.E. and Simon, D.H. 2004. Effects of 6- benzyl aminopurine treatments on the longevity of harvested grevillea 'Sylvia' inflorescences. *Plant Growth Regul.* 43: 9-14.

Archive of SID

Tables

Table 1. Chemical compounds

Treatment	Chemical compounds
T ₁	AgNO ₃ (300 mg l ⁻¹)
T ₂	8-HQS (250mg l ⁻¹)
T ₃	8-HQS (500mg l ⁻¹)
T ₄	SNP (1mg l ⁻¹)
T ₅	SNP (2mg l ⁻¹)
T ₆	8-HQS (250mg l ⁻¹) + SNP(1mg l ⁻¹)
T ₇	8-HQS (500mg l ⁻¹) + SNP(1mg l ⁻¹)
T ₈	8-HQS (250mg l ⁻¹) + SNP(2mg l ⁻¹)
T ₉	8-HQS (500mg l ⁻¹) + SNP(2mg l ⁻¹)
T ₁₀	Distilled Water
T ₁₁	Control

Table 2. Mean comparison of traits in 'Grand prix'

Treatment	Flower opening (%)	FW/DW	Flower diameter (mm)	RFW	Vaseliflife (days)
T ₁	48.75 f	4.506 ghi	81.38 e	69.92 i	7.250 hij
T ₂	84.17 ab	5.352 cde	94.05 d	100.6 a	11.17 bcde
T ₃	90.42 a	4.499 hi	97.31 bcd	90.01 f	8.833 fgh
T ₄	82.92 abc	4.179 ij	95.91 bcd	84.40 g	9.917 efg
T ₅	84.17 ab	4.447 hi	95.22 cd	91.41 f	10.50 def
T ₆	84.58 ab	5.116 def	98.76 bcd	97.87 abcd	12.92 b
T ₇	86.25 ab	5.006 efg	96.96 bcd	98.24 abcd	12.17 bcd
T ₈	86.25 ab	5.098 def	100.7 abc	99.31 abc	12.67 bc
T ₉	83.75 ab	4.805 fgh	100.3 abcd	93.41 ef	10.92 cde
T ₁₀	61.25 e	4.009 ijk	81.21 e	68.23 ij	7.667 hi
T ₁₁	47.50 f	3.544 k	64.85 f	62.51 k	5.500 j

Means followed by some letter within each column don't differ significantly according to Duncan multiple range test at P≤0.01

Table 3. Mean comparison of traits in 'Avalanche'

Treatment	Flower opening (%)	FW/DW	Flower diameter (mm)	RFW	Vaseliflife (days)
T ₁	30.00 g	4.930 efgh	86.23 e	64.59 k	5.833 j
T ₂	72.50 d	5.933 ab	95.39 bcd	95.86 cde	11.00 cde
T ₃	64.17 e	5.906 ab	98.00 bcd	79.37 h	8.667 gh
T ₄	85.83 ab	5.879 ab	100.6 abc	97.89 abcd	16.58 a
T ₅	85.83 ab	5.977 ab	98.19 bcd	97.84 abcd	18.17 a
T ₆	85.00 ab	5.684 abc	101.8 ab	97.24 abcd	11.42 bcde
T ₇	78.75 bcd	5.780 abc	96.58 bcd	96.57 bcde	10.75 de
T ₈	83.33 ab	6.042 a	100.6 abc	99.97 ab	11.83 bcd
T ₉	75.42 cd	5.796 abc	101.1 abc	96.56 bcde	10.00 efg
T ₁₀	82.92 abc	5.524 bcd	105.7 a	95.79 de	17.50 a
T ₁₁	30.00 g	3.822 jk	65.79 f	65.41 jk	6.083 ij

Means followed by some letter within each column don't differ significantly according to Duncan multiple range test at P≤0.01