

## Effect of Silicon on Growth and Ornamental Traits of Salt-stressed *Calendula officinalis* L.)

Hassan Bayat<sup>1\*</sup>, Morteza Alirezaie<sup>1</sup>, Hossein Neamati<sup>1</sup> and Ali Abdollahi Saadabad<sup>2</sup>

<sup>1</sup> Department of Horticultural Science, Faculty of Agriculture, Ferdowsi University of Mashhad, Mashhad, Iran.

<sup>2</sup> Department of Horticultural Science, Faculty of Agriculture, University of Tabriz, Tabriz, Iran.

Received: 17 December 2012

Accepted: 14 October 2013

\*Corresponding author's email: [hassanbayat55@gmail.com](mailto:hassanbayat55@gmail.com)

A pot experiment was carried out to determine the effect of foliage spraying of silicon (Si) on growth and ornamental characteristics of calendula grown under salt stress and greenhouse conditions. A factorial experiment based on completely randomized design was conducted with 3 levels of Si (0, 50 and 100 mg/l) and 3 levels of NaCl (0, 100 and 200 mM) with 4 replications. At flowering stage, Si was applied with spraying two times in two week intervals. NaCl was also applied as drench (200 ml per pot) in two days interval. The results showed that salinity decreased the growth, SPAD values, flower number per plant and flower diameter. However, foliar applications of Si resulted in greater root, shoot and total dry weight, plant height and leaf area of calendula plants under salt stress. The highest SPAD values were obtained under 100 mg/l Si application in all NaCl treatments. Salinity decreased number of flower per plant and flower diameter as ornamental characteristics; however Si increased them under salinity stress. Plants treated with 100 mg/l Si had the highest flower diameter at 100 mM of NaCl. Electrolyte leakage increased by salinity, however foliar application of Si significantly reduced electrolyte leakage under salt stress. These results suggest that the negative effects of salinity on the growth and ornamental characteristics of calendula plants can ameliorate by foliar application of Si treatments.

Abstract

**Keywords:** Electrolyte leakage, Foliar application, Landscape, Salinity, SPAD value.

## INTRODUCTION

One of the most important environmental factors limiting plant growth and productivity is salinity (Kaya *et al.*, 2003). In most arid and semiarid areas, this problem is accentuated by competition for high quality water among agriculture, industry and landscape users has promoted the use of alternative water sources for irrigation. Thus, marginal quality water, somewhat saline, will become important in these areas (Chartzoulakis *et al.*, 2002) and could be used for the irrigation of ornamental plants (Carter *et al.*, 2005). However, the use of low quality water for irrigation affects plants in different ways, depending on the degree of salt tolerance of the species (Alarcon *et al.*, 1994) and even within a given species (Sanchez-Blanco *et al.*, 2003). Salt stress can affect plant survival, biomass, plant height and plant morphology and affect the capacity of a plant to collect water and nutrients (Parida and Das, 2005). Salinity can cause hyper ionic and hyper osmotic effects on plants leading to membrane disorganization, increase in reactive oxygen species (ROS) levels and metabolic toxicity (Jaleel *et al.*, 2007a). The great effect of salinity is the inhibition of crop growth by the reduced hormone delivery from root to leaves (Jaleel *et al.*, 2007b).

Calendula, mostly known as the pot marigold, is planted widely in gardens and landscapes. It is popular for the lush color and aromatic scent. It grows in sun or partial shade and is easy to grow requiring little cultivation (Dole and Wilkins, 2004).

Silicon (Si) is the second most abundant element on the surface of the earth, yet its role in plant biology has been poorly understood. Silicon has not been considered an essential element for the growth of higher plants; however, soluble Si has enhanced the growth, development and yield of several plant species including rice (*Oryza sativa* L.), sugarcane (*Saccharum officinarum* L.) and most other cereals and several dicotyledons (Jones and Handreck, 1967; Elawad and Green, 1979; Takahashi *et al.*, 1990; Belanger *et al.*, 1995; Savant *et al.*, 1999), especially under biotic and abiotic stress conditions (Epstien, 1994). Silicon concentration in the soil solution is controlled by silicate minerals and ranges from 0.01 to 1.99 mM (Karathanasis, 2002). Silicon plays a significant role in imparting biotic and abiotic stress resistance and enhancing crop productivity (Okuda and Takahashi, 1965; Ma *et al.*, 1989; Epstein, 1994; Liang *et al.*, 1994). Improvement of salt tolerance has been reported by addition of Si in wheat (Ahmad *et al.*, 1992), barley (Liang *et al.*, 2003), rice (Yeo *et al.*, 1999), maize (Wang *et al.*, 2004), cucumber (Zhu *et al.*, 2004), rose (Savvas *et al.*, 2007) and zinnia (*Zinnia elegans*) (Kamenidou *et al.*, 2009). Savvas *et al.* (2002) reported that quality and peduncle stem thickness in gerbera flower increased when potassium silicate was included in the hydroponic nutrient solution. Hwang *et al.* (2005) also reported that applications of potassium silicate improved the growth and quality of cut miniature rose 'Pinocchio' in the rock wool culture system.

Water deficiency is a serious problem in arid and semi-arid regions of Iran that characterized by little rainfall, high solar radiation and high temperatures in the summer. In recent years, the normal seasonal droughts that have occurred in Iran have caused local and state government to enact water conservation ordinances. Urbanization and increases in population, however, are seriously threatening sustainable natural resources. At present, non-renewable groundwater resources are being depleted to an alarming extent. As high-quality water supply becomes limited, the use of saline water with high salt levels for landscape irrigation is being encouraged. While crop tolerance to salinity has been given considerable attention; fewer studies have dealt specifically with ornamental plants. Therefore, the aim of this research was to investigate the effects of Si on growth and quality flower of calendula grown under normal and salinity stress conditions.

## MATERIALS AND METHODS

### Plant material and treatments application

For this experiment, the seeds of *Calendula officinalis* L. 'Nana Bon Bon Naranja' were selected and germinated in cocopeat in plastic germination trays under greenhouse conditions at

Ferdowsi University of Mashhad (36°17'44" N and 59°36'42" E), Iran. Uniform size seedlings (40-days old) were transplanted to plastic pots (15 cm top diameter) and filled with mixture of loam soil:sand:compost (1:1:1,v:v). The plants were grown in a naturally illuminated greenhouse with night/day set temperatures of 18/24 °C. A factorial experiment based on completely randomized design was conducted with 3 levels of Si (0, 50 and 100 mg/l) and 3 levels of NaCl (0, 100 and 200 mM) with 4 replications. Potassium silicate (K<sub>2</sub>SiO<sub>3</sub>) was used for silicon treatments. Thirty days after transplanting, Si was applied on the foliage of calendula plants with a hand sprayer. The volume of the spray was 30 ml per pot. NaCl was also applied as drench (200 ml per pot) in two days interval. A control group of plants was grown without NaCl and sprayed with deionized water. All plants were harvested 60 days after planting (30 days after treatments) and separated into leaves, stem, and root.

### Measurements and data collection

Dry weights of separated roots and shoots were recorded on four randomly selected plants per treatment. Total leaf area was determined with a Delta-T Image Analysis System (Delta-T, LTD, Cambridge, UK). Plant height, flower diameter and number of flower per plant were also measured. Electrolyte leakage which is used to assess membrane permeability was determined according to Lutts *et al.* (1996). Leaf greenness or chlorophyll reading values (measured as the optical density, SPAD value) was recorded at the end of the experiment on three leaves per plant at similar middle positions of shoots for all plants in each treatment using a portable SPAD chlorophyll meter (SPAD 502, Minolta, Japan).

### Statistical analysis

Analysis of variance (ANOVA) for all the variables was carried out using the JMP8 software. Treatment means were compared using the protected Least Significant Difference (LSD) test at  $p < 0.05$  level.

## RESULTS

### Plant growth

Shoot, root and total dry weight were recorded as influenced by different levels of Si and NaCl. As shown in Fig. 1, shoot, root and total dry weight of calendula plants were lower at salt stress treatment as compared to non-saline conditions. However, foliar application of Si increased dry matter accumulation in all parts of calendula plants under salt stress. Application of Si (100 mg/l) under salt stress (100 mM NaCl) gave the higher values for these parameters than the other treatments. It indicated that foliar application of Si alleviated the growth inhibition induced by added NaCl. Exogenous application of Si also enhanced shoot, root and total plant dry weight under no salt stress (Fig. 1).




Fig. 1. The dry weight of shoot (A), root (B) and total plant (C) of calendula in response to foliar Si applications under salt stress. Different letters on top of bars indicate significant differences according to LSD test ( $p < 0.05$ ) at each salt level. Vertical bars indicate the mean  $\pm$  SE.




Fig. 2. Plant height (A) and leaf area (B) of calendula plant in response to foliar Si applications under salt stress. Different letters on top of bars indicate significant differences according to LSD test ( $p < 0.05$ ) at each salt level. Vertical bars indicate the mean  $\pm$  SE.

Salt stress significantly decreased plant height; however, foliar application of Si improved plant height under salt stress (Fig. 2A). Treatment 100 mg/l Si + 100 mM NaCl had the highest plant height as compared to the other treatments under salt stress. Foliar application of Si (100 mg/l) increased plant height by 27% as compared to control in 100 mM NaCl (Fig. 2A).

Salt stress decreased leaf area as compared to the non-saline conditions. NaCl (200 mM) decreased leaf area by 58%, but the reduction was 41% when 100 mg/l Si treatment was applied to NaCl-treated plants, as compared to control. Under non-saline conditions, foliar application of Si (100 mg/l) significantly increased leaf area (1.6 fold) as compared to control (Fig. 2B).

### Ornamental characteristics

Salt treatment (200 mM) significantly decreased number of flower per plant by 70% as compared to control, but the reduction was only 40% when Si (100 mg/l) was applied to NaCl-treated plants. Under non-salt stress condition, Si increased flower number by 35% as compared to control (Fig. 3A).

Flower diameter of calendula plants decreased dramatically with the increasing NaCl concentration. All Si treatments increased the flower diameter compared to non-treated plants both in absence and presence of salinity. Plants treated with 100 mg/l Si had the highest flower diameter at 100 mM of NaCl (Fig. 3B).




Fig. 3. Flower number per plant (A) and flower diameter (cm) (B) of calendula plant in response to foliar Si applications under salt stress. Different letters on top of bars indicate significant differences according to LSD test ( $p < 0.05$ ) at each salt level. Vertical bars indicate the mean  $\pm$  SE.




Fig. 4. SPAD value (A) and electrolyte leakage (B) of calendula plant in response to foliar Si applications under salt stress. Different letters on top of bars indicate significantly differences according to LSD test ( $p < 0.05$ ) at each salt level. Vertical bars indicate the mean  $\pm$  SE.

### SPAD value and electrolyte leakage

SPAD values were significantly decreased with the increasing salinity stress. However, foliar Si applications were caused to the elevated SPAD values. Under saline conditions, the highest SPAD values were obtained from 50 mg/l Si application in all NaCl treatments (Fig. 4A).

Electrolyte leakage increased by salt treatment; however, Si significantly decreased electrolyte leakage under salt stress. The lowest electrolyte leakage was obtained by 100 mg/l Si in 100 mM NaCl concentration under salt stress condition. Under no salt stress, foliar application of Si only slightly decreased electrolyte leakage compared to control (Fig. 4B).

### DISCUSSION

Under saline conditions, plant growth of calendula decreased significantly. Growth reduction under saline stress has been reported in various plants by many researchers (Senaratna *et al.*, 2000; Alpaslan and Gunes, 2001; Kaya *et al.*, 2003; Sivritepe *et al.*, 2003). As stated by Munns (2002), suppression of plant growth under saline conditions may either be due to decreased availability of water or to the toxicity of sodium chloride. Si treatments alleviated the deleterious effects of salinity on plant growth. Similar results were reported in rice (Matoh *et al.*, 1986; Yeo *et al.*, 1999), maize (Wang *et al.*, 2004), cucumber (Zhu *et al.*, 2004), rose (Savvas *et al.*, 2007) and zinnia (*Zinnia elegans*) (Kamenidou *et al.*, 2009) who observed exogenous Si treatments ameliorated the negative effects of salt stress on plant growth. It has been observed that GA1 and its precursor GA20 enhanced with N and Si application in rice cultivars (Hwang *et al.*, 2008). Gibberellins affect cell enlargement and division which leads to internode elongation in stems and increases stem height. Under saline growth conditions, it has been reported that the benefits of Si are due to the reduction of Na content in the shoots of rice, *P. juliflora*, and barley (Matoh *et al.*, 1986; Bradbury and Ahmad, 1990; Yeo *et al.*, 1999; Liang *et al.*, 2003), enhanced K uptake of barley (Liang *et al.*, 1996) and improved photosynthesis rate in barley and tomato (Liang, 1998; Al-Aghabary *et al.*, 2004). It has been also reported that Si increases the plant growth and yield of cucumber (Miyake and Takahashi, 1983; Voogt and Kreuzer, 1989).

Si increases rigidity of the mature leaves, which have a rougher texture and are held more horizontally, delays leaf senescence and increases chlorophyll content and ribulose, 1-5- biphosphate carboxylase activity (Adata and Besford, 1986). Moreover the induction of antioxidant enzymes and their protective role of membranes caused increasing the tolerance of plant to damages (Liang, 1999).

Salt stress decreased chlorophyll as compared to the non-saline conditions. These results are similar to those of Downton *et al.* (1985), Stepien and Klobus (2006) and Yildirim *et al.*, (2008) who indicated that chlorophyll content significantly decreased in the leaves of spinach and



cucumber plants with increasing NaCl concentration. In the present study, Si treated plants showed greater chlorophyll values than non-treated plants. These results are similar with Al-aghabary *et al.* (2004) that reported Si treatments caused to increase chlorophyll content of leaves of tomato under salt stress.

Salinity impairs membrane permeability and increases electrolyte leakage. However, application of Si partly maintained membrane permeability (Table 4B). Present study showed that Si reduced the amount of ion leakage in salt stressed calendula plants and Si protected the maintenance of membrane functions under stress conditions. Supporting evidence was shown when Si reduced electrolyte leakage salt stress condition in rice (Agarie *et al.*, 1998) leaves.

Salt stress decreased flower number per plant and flower diameter as compared to the non-saline conditions. In this experiment, different concentrations of Si increased flower number per plant and flower diameter. This is in agreement with the works by Savvas *et al.* (2002) and Kamenidou *et al.* (2010) that previously reported an increased gerbera flower quality when potassium silicate was added to the hydroponic nutrient solution. Hwang *et al.* (2005) also reported that applications of potassium silicate improved the growth and quality of cut flower miniature rose 'Pinocchio' in the rockwool culture system in agreement with the present results. It is possible that higher chlorophyll contents in Si treatments resulted in photosynthetic activity improvement and higher productivity (Table 4A).

## CONCLUSION

Based on the present results, Si alleviates the negative effect of salt stress on growth, chlorophyll reading values, electrolyte leakage and flower quality depending on the concentration of Si. Maximum alleviation of salt stress was found with 100mg/l Si application.

## Literature Cited

- Adatia, M.H. and Besford, R. T. 1986. The effects of silicon on cucumber plants grown in recirculating nutrient solution. *Annals of Botany*. 58: 343-351.
- Agarie, S., Hanaoka, N., Ueno, O., Miyazaki, A., Kubota, F., Agata, W. and Kaufman, P.B. 1998. Effects of silicon on tolerance to water deficit and heat stress in rice plants (*Oryza sativa* L.), monitored by electrolyte leakage. *Production Science*. 1: 96-103.
- Ahmad, R.R., Zaheer, H.S. and Ismail, S. 1992. Role of silicon in salt tolerance of wheat (*Triticum aestivum* L.). *Plant Science*. 85: 43-50.
- Al-Aghabary, K., Zhu, Z.J. and Shi, Q.H. 2004. Influence of silicon supply on chlorophyll content, chlorophyll fluorescence and antioxidative enzyme activities in tomato plants under salt stress. *Journal of Plant Nutrition*. 27: 2101-2115.
- Alarcon, J.J., Sanchez-Blanco, M.J., Bolarin, M.C. and Torrecillas, A. 1994. Growth and osmotic adjustment of two tomato cultivars during and after saline stress. *Plant Soil*. 166: 75-82.
- Alpaslan, M. and Gunes, A. 2001. Interactive effects of boron and salinity stress on the growth, membrane permeability and mineral composition of tomato and cucumber plants. *Plant Soil*. 236: 123-128.
- Belanger, R.R., Bowen, P.A., Ehret, D.L. and Menzies, J.G. 1995. Soluble silicon: Its role in crop and disease? *Management of greenhouse crops*. *Plant Disease*. 79: 329-336.
- Bradbury, M. and Ahmad, R. 1990. The effect of silicon on the growth of *Prosopis juliflora* growing in saline soil. *Plant Soil*. 125: 71-74.
- Carter, C.T., Grieve, C.M., Poss, J.A. and Suarez, D.L. 2005. Production and ion uptake of *Celosia argenta* irrigated with saline wastewaters. *Scientia Horticulturae*. 106: 387-394.
- Chartzoulakis, K., Loupassaki, M., Bertaki, M. and Androulakis, I. 2002. Effects of NaCl salinity on growth, ion content and CO<sub>2</sub> assimilation rate of six olive cultivars. *Scientia Horticulturae*. 96: 235-247.

- Dole, J.M. and Wilkins, H.F. 2004. Floriculture: Principles and species. Prentice Hall, Upper Saddle River, New Jersey.
- Downton, W.J.S., Grant, W.J.R. and Robinson, S.P. 1985. Photosynthetic and stomatal responses of spinach leaves to salt stress. *Plant Physiology*. 77: 85-88.
- Elawad, S.H. and Green, V.E. 1979. Silicon and the rice plant environment: A review of recent research. *Riso*. 28: 235-253.
- Epstein, E. 1994. The anomaly of silicon in plant biology. *Proceeding of the National Academy of Science*. 91: 11-17.
- Hwang, S.J., Hamayun, M., Kim, H.Y., Na, C.I., Kim, K.U., Shin, D.H., Kim, S.Y. and Lee, I.J. 2008. Effect of nitrogen and silicon nutrition on bioactive gibberellin and growth of rice under field conditions. *Journal of Crop Science and Biotechnology*. 10: 281-286.
- Hwang, S.J., Jeong, B.R. and Park, H.M. 2005. Effects of potassium silicate on the growth of miniature rose 'Pinocchio' grown on rock wool and its cut flower quality. *American Society for Horticultural Science*. 74: 242-247.
- Jaleel, C.A., Gopi, R., Manivannan, P. and Panneerselvam, R. 2007a. Antioxidative potentials as a protective mechanism in *Catharanthus roseus* (L.) G. Don. plants under salinity stress. *Turkish Journal of Botany*. 31: 245-251.
- Jaleel, C.A., Gopi, R., Manivannan, P. and Panneerselvam, R. 2007b. Responses of antioxidant defense system of *Catharanthus roseus* (L.) G. Don. To paclobutrazol treatment under salinity. *Acta Physiologica Plant*. 29: 205-209.
- Jones, L.H.P. and Handreck, K.A. 1967. Silica in soils, plants and animals. *Adv. Agron*. 19: 107-149.
- Kamenidou, S., Cavins, T.J. and Marek, S. 2009. Evaluation of silicon as a nutritional supplement for greenhouse zinnia production. *Scientia Horticulturae*. 119: 297-301.
- Kamenidou, S., Cavins, T.J. and Marek, S. 2010. Silicon supplements affect floricultural quality traits and elemental nutrient concentrations of greenhouse produced gerbera. *Scientia Horticulturae*. 119: 297-301.
- Karathanasis, A.D. 2002. Mineral equilibrium in environmental soil systems. pp. 109-151. In: J.B Dixon and S.B. Weed (eds.), *Soil mineralogy with environmental applications*. Soil Science Society of America, Madison.
- Kaya, C., Higgs, D., Ince, F., Amador, B.M., Cakir, A. and Sakar, E. 2003. Ameliorative effects of potassium phosphate on salt-stressed pepper and cucumber. *Journal of Plant Nutrition*. 26: 807-820.
- Liang, Y.C., Chen, Q., Liu, Q., Zhang, W.H. and Ding, R.X. 2003. Exogenous silicon increases antioxidant enzyme activity and reduces lipid peroxidation in roots of salt-stressed barley (*Hordeum vulgare* L.). *Journal of Plant Physiology*. 160: 1157-64.
- Liang, Y.C., Ma, T.S., Li, F.J. and Feng, Y.J. 1994. Silicon availability and response of rice and wheat to silicon in calcareous soils. *Communication in soil Science and Plant Analysis*. 25: 2285-2297.
- Liang, Y.C., Shen, Q.R., Shen, Z.G. and Ma, T.S. 1996. Effects of silicon on salinity tolerance of two barely cultivars. *Journal of Plant Nutrition*. 19: 173-183.
- Liang, Y.C. 1998. Effects of silicon on leaf ultrastructure, chlorophyll content and photosynthetic activity in barley under salt stress. *Pedosphere*. 8: 289-296.
- Liang, Y.C. 1999. Effects of silicon on enzyme activity and sodium, potassium and calcium concentration in barely under salt stress. *Plant Soil*. 29: 217-224.
- Lutts, S., Kinet, J.M. and Bouharmont, J. 1996. NaCl-induced senescence in leaves of rice (*Oryza sativa* L.) cultivars differing in salinity resistance. *Annals of Botany*. 8: 389-398.
- Ma, J., Nishimara, K. and Takahashi, E. 1989. Effect of silicon on the growth of rice plants at different growth days. *Soil Science and Plant Nutrition*. 35: 347-356.
- Matoh, T., Kairusmee, P. and Takahashi, E. 1986. Salt-induced damage to rice plants and alleviation

- effect of silicate. *Soil Science and Plant Nutrition*. 32: 295-304.
- Miyake, Y. and Takahashi, E. 1983. Effect of silicon on the growth of cucumber plant in soil culture. *Soil Science and Plant Nutrition*. 29: 463-471.
- Munns, R. 2002. Comparative physiology of salt and water stress. *Plant, Cell & Environment*. 25: 239-250.
- Okuda, A. and Takahashi, E. 1965. The role of silicon in the mineral nutrition of the rice plant. *Proc. Symposium of the International Rice Research Institute*. Johns Hopkins Press. Baltimore, M.D. p. 123-146.
- Parida, A.K. and Das, A.B. 2005. Salt tolerance and salinity effects on plants: A review. *Ecotoxicology and Environmental Safety*. 60: 324-349. Safety.
- Sanchez-Blanco, M.J., Rodriguez, P., Olmos, E., Morales, M.A. and Torrecillas, A. 2003. Contrasting physiological responses of dwarf sea-lavender and marguerite to simulated sea aerosol deposition. *Journal of Environment Quality*. 32: 3338-3344.
- Savant, N.K., Korndorfer, G.H., Datnoff, L.E. and Snyder, G. 1999. Silicon nutrition and sugarcane production: A review. *Plant Nutrition*. 22: 1853-1903.
- Savvas, D., Gizas, G., Karras, G., Lydakakis-Simantiris, N., Salahas, G., Papadimitriou, M. and Tsouka, N. 2007. Interactions between silicon and NaCl-salinity in a soilless culture of roses in greenhouse. *European Journal of Horticultural Science*. 72: 73-79.
- Savvas, D., Manos, G., Kotsiras, A. and Souvaliotis, S. 2002. Effects of silicon and nutrient induced salinity on yield, flower quality, and nutrient uptake of gerbera grown in a closed hydroponic system. *Journal of Applied Botany*. 76: 153-158.
- Senaratna, T., Touchell, D., Bunn, E. and Dixon, K. 2000. Acetyl salicylic acid (aspirin) and salicylic acid induce multiple stress tolerance in bean and tomato plants. *Plant Growth Regulation*. 30: 157-161.
- Sivritepe, N., Sivritepe, H.O. and Eris, A. 2003. The effects of NaCl priming on salt tolerance in melon seedlings grown under saline conditions. *Scientia Horticulturae*. 97: 229-237.
- Stepien, P. and Klobus, G. 2006. Water relations and photosynthesis in *Cucumis sativus* L. Leaves under salt stress. *Biologia Plantarum*. 50: 610-616.
- Takahashi, E., Ma, L.F. and Miyake, Y. 1990. The Possibility of silicon as an essential element for higher plants. *Comments Agriculture Food Chemistry*. 2: 99-122.
- Voogt, W. and Kreuzer, A. 1989. Cucumbers silicon raises yield but is difficult to supply. *Groenten-en-Fruit*. 45: 48-49.
- Wang, Y., Stass, A. and Horst, W. 2004. Apoplastic binding of aluminum is involved in silicon-induced amelioration of aluminum toxicity in maize. *Physiology*. 136: 3762-70.
- Yeo, A.R., Flowers, S.A., Rao, G., Welfare, K., Senanayake, N. and Flowers, T.J. 1999. Silicon reduces sodium uptake in rice (*Oryza sativa* L.) in saline conditions and this is accounted for by a reduction in the transpiration bypass flow. *Plant Cell Environment*. 22: 559-65.
- Yildirim, E., Turan, M. and Guvenc, I. 2008. Effect of foliar salicylic acid applications on growth, chlorophyll and mineral content of cucumber (*Cucumis sativus* L.) grown under salt stress. *Journal of Plant Nutrition*. 31: 593-612.
- Zhu, Z., Wei, G., Li, J., Qian, Q. and Yu, J. 2004. Silicon alleviates salt stress and increases antioxidant enzymes activity in leaves of salt-stressed cucumber (*Cucumis sativus* L.). *Plant Science*. 167: 527-33.