

# An Efficient *In Vitro* Propagation, Antioxidant and Antimicrobial Activities of *Aphyllorchis Montana* (Reichenb.f.)

Ganesan Mahendran

Research scholar, Department of Botany, Bharathiar University, Coimbatore- 641046, India.

Received: 11 June 2014

Accepted: 20 November 2014

\*Corresponding author's email: [mahendran0007@gmail.com](mailto:mahendran0007@gmail.com)

An *in vitro* plant regeneration protocol was successfully established in *Aphyllorchis montana*, a saprophytic achlorophyllous orchid by culturing immature seeds. Among the six basal media evaluated for seed germination, BM-TM medium was found to be the best followed by KC medium. After 40 days, all the media turned brown and the growths of the protocorms were arrested. Activated charcoal, 1 g/l in half strength BM-TM was found to be suitable for further development of protocorms. Half strength BM-TM medium was supplemented with different growth regulators either individually or in combinations for multiplication of shoots. Of the five cytokinins tested, TDZ at 6.8  $\mu\text{M}$  was found to be most effective for multiple shoot induction yielding  $17.24 \pm 0.27$  shoots after 10 weeks of culture. Addition of low concentration of NAA (1.3  $\mu\text{M}$ ) in MS medium supplemented with the cytokinin TDZ (6.8  $\mu\text{M}$ ) favoured shoot multiplication. A mean number of  $27.56 \pm 0.54$  shoots with  $3.92 \pm 0.11$  number of roots were produced per explant. The response of the seed derived protocorm to the different types of organic additives viz., peptone and yeast extract and coconut water was also evaluated. The addition of these organic additives to the medium containing TDZ enhanced the number of shoot regeneration. The plantlets were acclimatized in plastic pots containing sterilized vermiculite. The survival rate was 100 % when maintained in the culture room condition ( $25 \pm 2$  °C). Screening of the antibacterial, antioxidant activity and estimation of total phenolics and flavonoid content of methanolic extracts of micropropagated plants were also carried out and compared with that of the wild-grown plants. In all the tests, methanolic extract from wild-grown plants showed higher antioxidant, antimicrobial activity, total phenolics and flavonoid content than *in vitro* propagated plants.

Abstract

**Keywords:** Antimicrobial activity, Antioxidant activity, *Aphyllorchis montana*, Asymbiotic seed germination, *In vitro* propagation, Plant growth regulators, 2,2-Diphenyl-1-picrylhydrazyl.

**Abbreviations:** IAA (Indol-3-acetic acid), BM-TM (BM-1 Terrestrial orchid medium), IBA (Indole-3-butyric acid), NAA ( $\alpha$ -Naphthaleneacetic acid), TDZ (Thidiazuron), BA (6-Benzyl adenine), KIN (6-Furfurylaminopurine), 2-iP (2-Isopentenyladenine), GA<sub>3</sub> (Gibberellic acid), Zt (Zeatin), DPPH (2,2-diphenyl-1-picrylhydrazyl), ABTS (2,2'-azino-bis (3-ethylbenzothiazoline-6-sulfonic acid) diammonium salt), TPTZ (2,4,6-tripyridyl-S-triazine), EDTA (Ethylenediamine tetraacetic acid).

## INTRODUCTION

Orchids belong to the most diverse plant family known to man. They have complex lifecycle, mycorrhizal association (seed germination) and specific pollination syndrome. It is a family of considerable economic importance particularly in horticulture and floristry. Apart from the horticultural value, orchids are used in traditional herbal medicine. Many orchid species are threatened globally by over collection from the natural habitat for horticultural purpose. Plant tissue culture and micropropagation techniques play an important role in conservation programs and management of botanical collection.

*Aphyllorchis montana* (family Orchidaceae) is native to the region and rare in natural habitat (Sinu *et al.*, 2012) has drawn much attention in recent years. The extract of it has been in use since olden times to cure cough, cold, anemia and for its vitality strengthening properties (Prajapati *et al.*, 2003). *A. montana* is used in traditional Indian systems for diabetic activity (Bhavani *et al.*, 2012; Sreenu *et al.*, 2013). The species is also one of the important components of 'Astvarga' and used in the preparation of 'Chyavanprash', a highly popular Ayurvedic tonic. The medicinal properties are possibly due to production of secondary metabolites, including phenolic compounds. The role of antioxidants due to the phenolic compounds in this species cannot be ignored. To fulfill the high demand of pharmaceutical industries, at present raw material is largely being drawn from the wild; this has severely affected its availability in natural habitat and the species has been considered rare in the Western Ghats (Sinu *et al.*, 2012). In spite of this, there has been hardly any effort to commercially cultivate this species.

The vegetative growth of the terrestrial mycoheterotrophic orchid is absolutely an underground mechanism; it challenges to locate them in vegetative condition, species-specific fungi for seed germination, poor seed viability and low rate of germination in natural condition and poor rooting ability of vegetative cuttings of rhizomes. In spite of this, there has been hardly any effort to commercially cultivate this species. Inevitably, therefore, rapid multiplication of this important drug yielding genotype is imperative. Alternatively, *in vitro* micropropagation would be beneficial in accelerating large scale multiplication and conservation of this important plant species. *In vitro* seed germination has been suggested as a suitable propagation method for conservation of orchids (Kauth *et al.*, 2006; Stewart and Kane, 2006). Therefore, development of rapid protocol for high frequency *in vitro* plant regeneration in this important medicinal herb became necessary in order to reduce the existing pressure on natural populations and continuous supply of plant materials for the pharmaceutical industry. As tissue culture technique has now become a well-established method for large scale plants developed for commercial utilization of several endangered medicinal plants. The major goal of the present investigation was to standardize the best media for seed germination, growth regulators combinations for high frequency plantlet production from protocorm explants of *A. montana* species. In order to enhance the shoot bud multiplication rate, various growth regulators were examined to identify the best growth hormone combinations for maximum number of shoot buds production and to evaluate the antioxidant, antimicrobial activity of both wild-grown and *in vitro* regenerated plants.

## MATERIALS AND METHODS

### Plant material, explant preparation and surface sterilization

Immature seeds of *Aphyllorchis Montana* (Reichenb.f.) were collected from Vellingiri Hills (longitude 60-40' and 70-10' E and latitude 10o-55 and 11o-10' N 1200) at an altitude of 1000-1250 msl. Tamilnadu, India. Freshly collected green capsules were washed thoroughly under running tap water. The pods were then immersed in Teepol3 - 5%(v/v) for 2 - 5 minutes by continuous shaking and then rinsed with sterile distilled water thrice and the pod were pretreated with 0.1% (W/V) of Bavistin, a fungicide for 5 minutes and rinsed in double distilled water. Then the explants were surface sterilized in 0.01% mercuric chloride solution for 5 minutes and rinsed thoroughly with sterile distilled water (5-7 times). The capsules were dipped in 70% ethanol for 30 seconds

and flamed. The surface sterilized pods were cut opened with sterile blade and seeds were extracted using sterile forceps and spread as thin film in test tubes containing 20 ml of culture media. Cultures were maintained at  $25 \pm 2^\circ\text{C}$  under cool white fluorescent tubes at a light intensity of  $50 \mu\text{mol m}^{-2} \text{s}^{-1}$  with 16/8-h L/D photoperiod.

### Seed viability test

Seed viability was tested according to Vellupillai *et al.* (1997). For enumerating seed viability percentage, seeds from the fresh capsules were treated with 1% (w/v) 2, 3, 5-triphenyl tetrazolium chloride (pH 7.0) in the dark overnight. Treated seeds were observed with a light microscope and scored as either viable (red embryo) or nonviable (white embryo).

### Optimization of culture medium for asymbiotic seed culture

Immature seeds of *A. Montana* were inoculated on Knudson C modified Morel (Morel 1965; KCM), Lindemann (Lindemann *et al.*, 1970; LM), Mitra medium (Mitra *et al.*, 1976; M), Knudson C medium (Knudson, 1946; KC), Murashige and Skoog medium (Murashige and Skoog, 1962, MS) and BM-1-Terrestrial Orchid media (Van Waes and Debergh, 1986 procured from Hi-Media Laboratories Mumbai, India) initially to find out the suitable medium for maximum seed germination. The best medium for seed germination was selected for further studies. All media contained 2% sucrose and were solidified with 0.8% agar (Hi Media Laboratories, India). Activated charcoal (Hi-Media Laboratories Mumbai, India) was added at 0.5, 1.0 or 2.0 g/l to half strength basal BM-TM medium to arrest phenolic exudation and further development of the protocorms. The pH of the media was adjusted to 5.6–5.8 with 1 N NaOH or HCl before autoclaving at  $121^\circ\text{C}$ , 105 kPa for 20 min.

### Multiplication of protocorms

For the multiplication of protocorms, BM-TM medium was supplemented with cytokinins such as BA (1.10 to 8.80  $\mu\text{M}$ ), Kn (1.15 to 9.20  $\mu\text{M}$ ), 2-iP (1.01 to 8.12), Zt (1.1 to 9.1  $\mu\text{M}$ ) and TDZ (1.1 to 9.0  $\mu\text{M}$ ) either individually or in combination with NAA (1.3  $\mu\text{M}$ ). All media contained 20  $\text{g l}^{-1}$  sucrose and were solidified with 0.8% agar (Hi Media Laboratories, India). The cultures were maintained at  $25 \pm 2^\circ\text{C}$  temperature with 75–80% relative humidity and a 16/8 h (light/dark) photoperiod provided with diffuse light ( $50 \mu\text{mol m}^{-2} \text{s}^{-1}$ ). Final observation on the number of multiple shoot and the shoot length, root number and root length were recorded after 70 days of culture.

### Ex vitro plant establishment

For *ex vitro* establishment, well-developed plantlets were rinsed thoroughly with tap water to remove residual nutrients and agar from the plant body and transplanted to plastic pot containing vermiculite. The plastic pots were covered initially with polyethylene bags and maintained for two months inside the culture room for acclimatization under cool white tubular fluorescent lights (40 W, 220 V, Philips Electronics India Ltd.) at  $50 \mu\text{mol}^{-1} \text{m}^{-2} \text{s}^{-1}$  with a 16 h photoperiod at  $25 \pm 2^\circ\text{C}$ . The transplants were transferred to National Orchidarium, Yearcaud, Tamil Nadu.

### Preparation of methanol extracts

The biomass from *in vitro* propagated plants (8 week old culture, BM-TM medium supplemented with TDZ (6.8  $\mu\text{M}$ ) and NAA (1.3  $\mu\text{M}$ ) shoots were collected) and wild grown plants (whole plants) were washed under tap water and dried in oven at  $60^\circ\text{C}$  for two days. The material was powdered by using electric blender and stored in clean labelled airtight bottles. The powder (100g) was extracted by maceration in 300 ml of methanol (100%) for 3 days with frequent agitation. The mixture was filtered through Whatman No. 1 filter paper and the filtrate was concentrated and dried in petridishes at  $60^\circ\text{C}$  in the oven.

### **Estimation of total phenolics (TPC) and flavonoids content (TFC)**

The total phenolics content of the extracts was determined and calculated as gallic acid equivalent (GAE) in mg/g DW from the calibration curve according to method described by Sidhuraju and Becker (2003). The total flavonoids content of sample extract was determined following a colorimetric method and values were expressed as mg/g rutin equivalent (RE) of extract according to the method described by Zhishen *et al.* (1999).

### ***In vitro* antioxidant activity**

The free radical scavenging activity of the *A. Montana* methanol extracts of wild-grown plants and *in vitro* propagated plants were evaluated by using DPPH• (Blois, 1958), ABTS<sup>+</sup> cation radical (Re *et al.*, 1999) and ferric reducing antioxidant power (FRAP) activity (Pulido *et al.*, 2000) were measured using standard methods.

### **Antimicrobial activity**

#### **Test bacteria**

The antibacterial activity of isolated compounds 1-6 were evaluated against 6 pathogenic bacteria such as *Escherichia coli* (ATCC 25922), *Pseudomonas aeruginosa* (ATCC 27853), *Staphylococcus aureus* (ATCC 29213), *Streptococcus pneumoniae* (ATCC 33400) *Klebsiella pneumoniae* (ATCC 10031) and *Bacillus subtilis* (ATCC 6633) procured from the Institute of Microbial Technology (IMTECH) Chandigarh, India. All the strains were stored in the appropriate medium before use.

#### **Disc diffusion method**

Disc diffusion method (Joshi *et al.*, 2010) was used for the evaluation of antibacterial activity of *A. Montana* methanol extracts of wild-grown plants and *in vitro* propagated plants using 100 µl of suspension containing 10<sup>8</sup> CFU/ml of bacteria spread on the inoculated agar. A sterile cotton swab was dipped into the inoculum suspension to remove the excess of fluid. Whatmann filter paper discs (6 mm diameter) were prepared at the concentration of 25 µg/disc for wild-grown plants and *in vitro* propagated plants extracts and 10 µg/disc reference antibiotic (Ciprofloxacin). A disc prepared with only the corresponding volume of DMSO was used as negative control. The petriplates were then incubated and antimicrobial activity was evaluated by measuring the diameter of the zones of inhibition around the disc. The experiments were repeated in triplicate and the result was expressed as average value.

#### **Minimum inhibitory concentration**

The minimum inhibitory concentration (MIC) of wild-grown plants and *in vitro* propagated plants of *A. montanawas* determined using the micro-dilution assay in 96-well microplates (Siddiqi *et al.*, 2011). Briefly, 500 µl of each re-suspended sample (1.0 mg/ml) in DMSO (2%). Serial two-fold dilutions were prepared from the stock solution to give concentrations ranging from 500 µg to 3.90 µg/ml of the wild-grown plants and *in vitro* propagated plants extracts. The highest concentration of DMSO remaining after dilution (5%, v/v) caused no inhibition to bacterial growth. DMSO served as negative control. Streptomycin and Ciprofloxacin were served as positive controls. An aliquot of 100 µl standardized suspension of the test bacteria (10<sup>8</sup> CFU/ml) was transferred to a well of 96 plates. Then, 100 µl diluted samples were also added to each well and incubated at 37°C for 24h. The MIC was defined as the lowest concentration of samples which inhibited the visible growth of tested microorganisms. For further re-confirmation, 20 µl of MTT reagent (1mg/ml) was added as an indicator for microbial growth to each well of the microtitre plates, followed by 20 min incubation at 37°C. The reagent-suspension colour will remain clear or yellowish indicating complete inhibition activity as opposed to dark blue for growth (Eloff, 1998). The MIC was recorded as the most repeatable minimum concentration of triplicate.

## Experimental design and statistical analysis

The percent of seed germination was recorded after 10, 20 and 30 days of culture. Percent germination was calculated by dividing the number of germinated seeds by the total number of seeds inoculated. Number of multiple shoots, height, root number and root length was recorded after 70 days of culture. The experiments were repeated thrice and each set had five replicates. The significance of differences among means was assessed out using Duncan's multiple range test at  $P <$




Fig.2. Asymbiotic seed germination and multiple shoot development of *A. montana* (Bar-1cm)

- Seed derived protocorms developed on BM-TM medium after 30 days of culture
- Phenolic exudation on BM-TM medium
- Development of protocorms on half strength BM-TM medium with activated charcoal 1 g/l
- Development of multiple protocorm on half strength BM-TM medium supplemented with TDZ at  $9.8 \mu\text{M}$
- Multiple protocorm development on half strength BM-TM medium fortified with TDZ ( $6.8 \mu\text{M}$ ) and 20% CW
- Regenerated plantlet
- Hardened plantlet
- Hardened plantlet after 2 months under *ex vitro* condition




Fig.1. Effect of various culture media on seed germination of *A. montana*  
Error bars indicates  $\pm$  S.E

0.05 (ANOVA). Data of the antioxidant, total phenolic and flavonoid assays were expressed as the mean  $\pm$  standard deviation (SD) of three independent measurements. Correlation analysis was performed between phenolics and flavonoids with antioxidant activity using Pearson correlation two-tailed. The results were analyzed statistically using SPSS Version 17 (SPSS Inc., Chicago, USA).

## RESULTS

### *In vitro* seed germination

Tetrazolium (TZ) viability test indicated a mean embryo viability of 90%. Seeds germinated on all the media tested, however, the percentage of seed germination and protocorm development was not consistent and varied in different media. The embryos enlarged and occupied the entire seed coat within 10 days after sowing. Germination as evidenced by enlargement of the embryo was first observed in BM-TM medium, followed by both KC and KCM medium. The seed germination was low in MS, Mitra and Lindmann orchid medium (Fig. 1).

After 30 days, the embryos by repeated cell divisions emerged and rupturing the testa in BM-TM medium (Fig. 2a). The percentage seed germination was 79% in BM-TM basal medium (Fig. 2a), 52% in KC medium and 42.50% in KCM medium. After 40 days, all the media turned brown possibly indicating phenolic exudation and further development of protocorms were arrested (Fig. 2b). The above medium was reduced half strength and supplemented with activated charcoal at various concentrations (0.5 to 3 g/l) to pre-

Table 1. Effect of cytokinins on multiple shoot induction from seed derived protocorm of *A. montana*.

BA ( $\mu\text{M/l}$ )	Kn ( $\mu\text{M/l}$ )	Z ( $\mu\text{M/l}$ )	TDZ ( $\mu\text{M/l}$ )	2ip ( $\mu\text{M/l}$ )	No. of hoots/explant	Shoot length (cm)	No. of Root/explant	Length of root (cm)
1.10	-	-	-	-	1.20±0.71 <sup>g</sup>	2.60±0.66 <sup>bc</sup>	3.76±0.23 <sup>b</sup>	1.53±0.22 <sup>d</sup>
2.20	-	-	-	-	2.56±0.54 <sup>f</sup>	2.00±0.12 <sup>d</sup>	3.25±0.45 <sup>c</sup>	1.09±0.46 <sup>de</sup>
3.30	-	-	-	-	5.12±0.33 <sup>d</sup>	2.41±0.34 <sup>c</sup>	2.30±0.76 <sup>de</sup>	1.22±0.34 <sup>de</sup>
4.40	-	-	-	-	9.51±0.49 <sup>c</sup>	2.01±0.52 <sup>d</sup>	2.38±0.25 <sup>de</sup>	2.19±0.63 <sup>c</sup>
6.60	-	-	-	-	8.51±0.12 <sup>c</sup>	1.16±0.22 <sup>f</sup>	2.13±0.67	2.72±0.62 <sup>b</sup>
8.80	-	-	-	-	5.76±0.21 <sup>d</sup>	1.00±0.40 <sup>f</sup>	1.84±0.12 <sup>f</sup>	3.30±0.20 <sup>a</sup>
-	1.15	-	-	-	1.10±0.73 <sup>g</sup>	1.00±0.56 <sup>f</sup>	1.90±0.71 <sup>f</sup>	3.09±0.90 <sup>a</sup>
-	2.32	-	-	-	1.32±0.26 <sup>g</sup>	2.11±0.77 <sup>d</sup>	2.18±0.29 <sup>e</sup>	3.12±0.17 <sup>a</sup>
-	3.45	-	-	-	2.65±0.12 <sup>f</sup>	2.19±0.14 <sup>d</sup>	2.64±0.21 <sup>d</sup>	2.16±0.24 <sup>c</sup>
-	4.64	-	-	-	3.97±0.75 <sup>de</sup>	2.52±0.40 <sup>bc</sup>	1.89±0.78 <sup>f</sup>	2.06±0.87 <sup>c</sup>
-	6.90	-	-	-	5.14±0.65 <sup>d</sup>	2.82±0.49 <sup>b</sup>	3.65±0.62 <sup>bc</sup>	1.72±0.89 <sup>d</sup>
-	9.20	-	-	-	7.50±0.30 <sup>cd</sup>	2.42±0.43 <sup>c</sup>	3.12±0.62 <sup>cd</sup>	1.16±0.51 <sup>de</sup>
-	-	1.1	-	-	4.60±0.16 <sup>de</sup>	2.63±0.87 <sup>bc</sup>	1.34±0.52 <sup>g</sup>	1.71±0.41 <sup>d</sup>
-	-	2.2	-	-	1.25±0.50 <sup>g</sup>	2.61±0.54 <sup>bc</sup>	2.00±0.70 <sup>e</sup>	1.54±0.26 <sup>d</sup>
-	-	3.3	-	-	2.53±0.11 <sup>f</sup>	3.00±0.20 <sup>a</sup>	3.12±0.41 <sup>cd</sup>	2.73±0.31 <sup>b</sup>
-	-	4.5	-	-	5.54±0.64 <sup>d</sup>	2.81±0.28 <sup>b</sup>	3.89±0.63 <sup>b</sup>	2.39±0.32 <sup>bc</sup>
-	-	6.8	-	-	7.20±0.81 <sup>cd</sup>	3.00±0.79 <sup>a</sup>	3.60±0.29 <sup>bc</sup>	3.66±0.12 <sup>a</sup>
-	-	9.1	-	-	12.65±0.16 <sup>b</sup>	2.84±0.65 <sup>b</sup>	2.72±0.56 <sup>d</sup>	2.53±0.34 <sup>bc</sup>
-	-	-	1.1	-	2.30±0.83 <sup>f</sup>	3.16±0.66 <sup>a</sup>	2.29±0.12 <sup>de</sup>	1.75±0.34 <sup>d</sup>
-	-	-	2.2	-	3.65±0.76 <sup>e</sup>	3.00±0.79 <sup>a</sup>	2.69±0.34 <sup>d</sup>	2.00±0.00 <sup>c</sup>
-	-	-	3.3	-	5.65±0.76 <sup>d</sup>	2.71±0.54 <sup>b</sup>	2.93±0.18 <sup>de</sup>	2.19±0.11 <sup>c</sup>
-	-	-	4.5	-	8.97±0.75 <sup>c</sup>	2.01±0.02 <sup>d</sup>	2.31±0.51 <sup>de</sup>	2.30±0.13 <sup>bc</sup>
-	-	-	6.8	-	17.24±0.27 <sup>a</sup>	1.46±0.02 <sup>e</sup>	2.12±0.11 <sup>e</sup>	2.70±0.71 <sup>b</sup>
-	-	-	9.0	-	12.30±0.65 <sup>b</sup>	1.00±0.40 <sup>f</sup>	4.94±0.22 <sup>a</sup>	2.00±0.92 <sup>c</sup>
-	-	-	-	1.01	2.15±0.58 <sup>f</sup>	2.11±0.57 <sup>d</sup>	2.12±0.34 <sup>e</sup>	1.09±0.92 <sup>de</sup>
-	-	-	-	2.03	2.87±0.98 <sup>f</sup>	2.41±0.54 <sup>e</sup>	3.81±0.70 <sup>b</sup>	1.48±0.44 <sup>d</sup>
-	-	-	-	3.04	3.87±0.70 <sup>e</sup>	3.00±0.69 <sup>a</sup>	3.61±0.07 <sup>bc</sup>	2.00±0.23 <sup>c</sup>
-	-	-	-	4.06	5.82±0.23 <sup>d</sup>	3.00±0.10 <sup>a</sup>	3.53±0.75 <sup>bc</sup>	2.50±0.32 <sup>bc</sup>
-	-	-	-	6.09	9.70±0.65 <sup>c</sup>	2.42±0.40 <sup>c</sup>	3.50±0.65 <sup>bc</sup>	2.43±0.43 <sup>c</sup>
-	-	-	-	8.12	12.65±0.45 <sup>b</sup>	2.84±0.92 <sup>b</sup>	3.12±0.29 <sup>c</sup>	2.87±0.69 <sup>b</sup>

Values represent mean  $\pm$  S.E. of twice repeated experiments each with 5 replications. Means in a column with the different letter (superscript) are significantly different according to DMRT ( $P < 0.05$ ).

vent the phenolic exudation and induce further development of protocorms (Fig. 2c). Among the different concentrations of activated charcoal tested, 1g/l completely arrested phenolic exudation and it was found to be the optimum concentration for the development of protocorms.

### Effect of cytokinins on development of multiple shoots

The seed derived protocorm explants cultured on half strength BM-TM medium without any growth regulators formed a single shoot. In the presence of cytokinin, the explants responded positively in producing multiple shoots (multiple seedlings). Of the five cytokinins tested, TDZ was found to be more efficient than other cytokinins with respect to initiation and subsequent proliferation of shoots (Table 1). Among the different levels TDZ tested, the maximum number of shoots was observed on the half strength BM-TM medium containing 6.8  $\mu\text{M}$  of TDZ (17.24  $\pm$  0.27) (Table 1, Fig. 2d). The shoot buds first appeared as small white protuberances at basal surface at the protocorm, which eventually developed into multiple shoots within 30-35 days. The number of shoot buds increased with increasing concentration of TDZ up to an optimal level of 6.8  $\mu\text{M}$ . Among the various concentrations of Zt and 2-ip tested, maximum number of the multiple shoots was recorded in half strength BM-TM medium supplemented with 9.1  $\mu\text{M}$  Z and 8.12  $\mu\text{M}$  2-ip (12.65 shoots/explant). NAA (1.3  $\mu\text{M}$ ) in combination with cytokinins also favored shoot formation. However, the response for multiple shoot induction varied (Table 2). Among the different combinations, maximum number of shoots were produced in half strength BM-TM medium supplemented with TDZ (6.8  $\mu\text{M}$ ) and NAA (1.3  $\mu\text{M}$ ).

### Effect of TDZ and growth adjuvants on multiple shoot induction

Multiple shoots were also formed when seed derived protocorms were cultured on BM-TM medium supplemented with TDZ at 6.8  $\mu\text{M}$  in combination with growth adjuvants like pep-

Table 2. Effect of cytokinins and auxin on multiple shoot developed from seed derived protocorm of *A. montana*.

BA (μM)	Kn (μM)	Z (μM)	TDZ (μM)	2-ip (μM)	NAA (μM)	No. of shoot /explant	Shoot length (cm)	No. of Root/ex-plant	Root length (cm)
1.10	-	-	-	-	1.3	1.48±0.61 <sup>h</sup>	2.06±0.12 <sup>bc</sup>	4.56±0.11 <sup>bc</sup>	1.00±0.11 <sup>d</sup>
2.20	-	-	-	-	1.3	3.76±0.54 <sup>fg</sup>	2.34±0.87 <sup>b</sup>	5.12±0.34 <sup>bc</sup>	1.22±0.54 <sup>d</sup>
3.30	-	-	-	-	1.3	7.62±0.53 <sup>e</sup>	2.65±0.66 <sup>b</sup>	6.74±0.54 <sup>b</sup>	1.76±0.76 <sup>d</sup>
4.40	-	-	-	-	1.3	12.47±0.34 <sup>d</sup>	2.70±0.33 <sup>b</sup>	10.12±0.61 <sup>a</sup>	1.00±0.12 <sup>d</sup>
6.60	-	-	-	-	1.3	10.51±0.54 <sup>d</sup>	3.11±0.44 <sup>b</sup>	7.09±0.12 <sup>b</sup>	1.22±0.62 <sup>d</sup>
8.80	-	-	-	-	1.3	6.76±0.63 <sup>e</sup>	1.00±0.31 <sup>c</sup>	4.17±0.44 <sup>bc</sup>	2.00±0.43 <sup>c</sup>
-	1.15	-	-	-	1.3	1.76±0.21 <sup>h</sup>	1.22±0.31 <sup>c</sup>	1.30±0.41 <sup>e</sup>	1.56±0.23 <sup>d</sup>
-	2.32	-	-	-	1.3	2.11±0.46 <sup>g</sup>	2.23±0.54 <sup>bc</sup>	2.32±0.18 <sup>d</sup>	1.12±0.21 <sup>d</sup>
-	3.45	-	-	-	1.3	3.54±0.14 <sup>fg</sup>	2.43±0.54 <sup>b</sup>	3.11±0.65 <sup>c</sup>	2.16±0.32 <sup>c</sup>
-	4.64	-	-	-	1.3	4.32±0.80 <sup>f</sup>	2.65±0.10 <sup>b</sup>	4.50±0.23 <sup>bc</sup>	2.06±0.43 <sup>c</sup>
-	6.90	-	-	-	1.3	5.64±0.55 <sup>f</sup>	3.13±0.23 <sup>a</sup>	4.21±0.11 <sup>bc</sup>	1.10±0.54 <sup>d</sup>
-	9.20	-	-	-	1.3	7.50±0.37 <sup>e</sup>	3.61±0.55 <sup>a</sup>	4.82±0.62 <sup>bc</sup>	1.21±0.11 <sup>d</sup>
-	-	1.1	-	-	1.3	1.60±0.56 <sup>h</sup>	2.97±0.16 <sup>ab</sup>	1.02±0.11 <sup>e</sup>	1.65±0.20 <sup>d</sup>
-	-	2.2	-	-	1.3	2.62±0.84 <sup>g</sup>	3.20±0.19 <sup>a</sup>	2.34±0.45 <sup>d</sup>	1.23±0.43 <sup>d</sup>
-	-	3.3	-	-	1.3	2.70±0.65 <sup>g</sup>	3.61±0.89 <sup>a</sup>	2.23±0.41 <sup>d</sup>	2.73±0.31 <sup>b</sup>
-	-	4.5	-	-	1.3	4.47±0.07 <sup>f</sup>	3.44±0.29 <sup>a</sup>	3.89±0.63 <sup>c</sup>	2.39±0.32 <sup>c</sup>
-	-	6.8	-	-	1.3	5.32±0.66 <sup>f</sup>	3.87±0.64 <sup>a</sup>	3.60±0.29 <sup>c</sup>	3.66±0.12 <sup>a</sup>
-	-	9.1	-	-	1.3	7.33±0.87 <sup>e</sup>	3.98±0.16 <sup>a</sup>	2.72±0.56 <sup>d</sup>	2.53±0.34 <sup>b</sup>
-	-	-	1.1	-	1.3	4.81±0.33 <sup>f</sup>	3.16±0.62 <sup>a</sup>	2.29±0.12 <sup>d</sup>	1.75±0.34 <sup>d</sup>
-	-	-	2.2	-	1.3	7.12±0.87 <sup>e</sup>	3.45±0.69 <sup>a</sup>	2.69±0.34 <sup>d</sup>	2.00±0.00 <sup>c</sup>
-	-	-	3.3	-	1.3	12.57±0.61 <sup>d</sup>	2.67±0.21 <sup>b</sup>	2.93±0.18 <sup>cd</sup>	2.19±0.11 <sup>c</sup>
-	-	-	4.5	-	1.3	19.78±0.81 <sup>bc</sup>	2.35±0.98 <sup>b</sup>	3.31±0.51 <sup>c</sup>	2.30±0.13 <sup>c</sup>
-	-	-	6.8	-	1.3	27.56±0.54 <sup>a</sup>	1.56±0.33 <sup>c</sup>	3.92±0.11 <sup>bc</sup>	2.70±0.71 <sup>b</sup>
-	-	-	9.0	-	1.3	22.65±0.11 <sup>b</sup>	1.34±0.54 <sup>c</sup>	4.23±0.32 <sup>bc</sup>	3.41±0.12 <sup>a</sup>
-	-	-	-	1.01	1.3	2.98±0.12 <sup>g</sup>	2.43±0.67 <sup>b</sup>	2.12±0.34 <sup>d</sup>	1.32±0.14 <sup>d</sup>
-	-	-	-	2.03	1.3	1.41±0.17 <sup>h</sup>	2.78±0.88 <sup>b</sup>	3.78±0.32 <sup>c</sup>	1.98±0.35 <sup>cd</sup>
-	-	-	-	3.04	1.3	3.25±0.24 <sup>fg</sup>	3.22±0.23 <sup>a</sup>	3.24±0.34 <sup>c</sup>	2.23±0.34 <sup>c</sup>
-	-	-	-	4.06	1.3	5.98±0.35 <sup>f</sup>	3.54±0.82 <sup>a</sup>	3.76±0.87 <sup>c</sup>	2.17±0.41 <sup>c</sup>
-	-	-	-	6.09	1.3	10.70±0.89 <sup>d</sup>	2.17±0.10 <sup>bc</sup>	3.87±0.76 <sup>c</sup>	2.21±0.91 <sup>c</sup>
-	-	-	-	8.12	1.3	16.43±0.54 <sup>c</sup>	1.65±0.92 <sup>c</sup>	3.15±0.33 <sup>c</sup>	2.11±0.43 <sup>c</sup>

Values represent mean ± S.E. of twice repeated experiments each with 5 replications. Means in a column with the different letter (superscript) are significantly different according to DMRT (P < 0.05).

Table 3. Effect of growth adjuvants on multiple shoot induction from seed derived protocorm of *A. montana*.

TDZ (μM)	Peptone (g/l)	Coconut water (%)	Yeast extract (g/l)	No. of shoots	Shoot length (cm)	Root number	Root length (cm)
6.8	Control			17.24±0.27 <sup>e</sup>	1.46±0.02 <sup>c</sup>	2.12±0.11 <sup>d</sup>	2.70±0.71 <sup>a</sup>
6.8	-	-	-	18.66±0.11 <sup>e</sup>	1.11±0.31 <sup>e</sup>	2.19±0.32 <sup>d</sup>	1.92±0.22 <sup>b</sup>
6.8	0.5	-	-	20.21±0.26 <sup>de</sup>	1.16±0.82 <sup>e</sup>	1.98±0.18 <sup>e</sup>	1.70±0.43 <sup>b</sup>
6.8	1.0	-	-	23.82±0.32 <sup>c</sup>	0.97±0.12 <sup>f</sup>	1.50±0.12 <sup>f</sup>	1.40±0.65 <sup>c</sup>
6.8	2.0	-	-	20.89±0.44 <sup>de</sup>	0.46±0.14 <sup>g</sup>	1.00±0.43 <sup>g</sup>	0.98±0.36 <sup>e</sup>
6.8	3.0	-	-	21.71±0.54 <sup>d</sup>	1.76±0.23 <sup>b</sup>	2.12±0.22 <sup>d</sup>	1.10±0.12 <sup>d</sup>
6.8	-	5	-	23.61±0.11 <sup>c</sup>	1.49±0.62 <sup>c</sup>	3.92±0.31 <sup>c</sup>	1.87±0.11 <sup>b</sup>
6.8	-	10	-	27.24±0.76 <sup>b</sup>	1.17±0.39 <sup>e</sup>	5.54±0.52 <sup>a</sup>	1.72±0.18 <sup>b</sup>
6.8	-	15	-	28.33±0.27 <sup>a</sup>	1.98±0.42 <sup>a</sup>	7.21±0.29 <sup>a</sup>	1.45±0.13 <sup>c</sup>
6.8	-	20	-	17.89±0.61 <sup>e</sup>	1.21±0.26 <sup>d</sup>	1.00±0.32 <sup>g</sup>	0.50±0.17 <sup>f</sup>
6.8	-	-	0.5	14.32±0.79 <sup>f</sup>	1.11±0.13 <sup>e</sup>	1.00±0.11 <sup>g</sup>	0.40±0.11 <sup>f</sup>
6.8	-	-	1.0	12.18±0.43 <sup>g</sup>	1.00±0.43 <sup>f</sup>	0.91±0.11 <sup>g</sup>	0.10±0.03 <sup>g</sup>
6.8	-	-	2.0	9.29±0.99 <sup>h</sup>	1.00±0.54 <sup>f</sup>	0.00±0.00 <sup>h</sup>	0.00±0.00 <sup>h</sup>
6.8	-	-	3.0				

Values represent mean ± S.E. of twice repeated experiments each with 5 replications. Means in a column with the different letter (superscript) are significantly different according to DMRT (P < 0.05).

tone, yeast extract and coconut water (0.5 g- 3 g/l or 5-20%). The mean number of multiple shoots varied among the treatments (Table 3). A significantly higher number of multiple shoots (28.33 ± 0.27) were obtained on medium supplemented with TDZ (6.8 μM) and 20% CW (Fig. 2e). Other organic additives were also effective in inducing multiple shoots, however their number remained low when compared to CW (20%). The mean length of the shoot per explant was in the range of 0.46 ± 0.14 - 1.76 ± 0.23 cm which is lower when compared to the medium supplemented with TDZ (6.8 μM) individually (Table 1).

Table 4. Comparison of total phenol, flavonoids content and antioxidant activity of *in vitro* propagated plant and wild grown plant of *A. montana*

Sample	Total phenolics content (TPC) (mg/ga)	Total flavonoids content (TFC) (mg/gb)	DPPH IC <sub>50</sub> (µg/ml)	FRAP (mmol Fe(II)/g extractc)	ABTS (µmol trolox/g extractd)
Wild grown plant extract	121.34 ± 1.23	93.62 ± 1.17	26.87 <sup>a</sup>	2987.10 ± 19.33 <sup>a</sup>	38912.19 ± 456.12 <sup>a</sup>
<i>In vitro</i> propagated plants	61.89 ± 3.19	34.59 ± 1.14	41.45 <sup>c</sup>	1034.23 ± 67.44 <sup>c</sup>	20246.53 ± 291.24 <sup>c</sup>
BHA			34.59 <sup>b</sup>	1213.89 ± 80.86 <sup>b</sup>	21682.08 ± 359.47 <sup>b</sup>

Values are mean of three replicate determinations standard deviation. Mean values followed by different superscript in a column are significantly different ( $P < 0.05$ ).

<sup>a</sup> gallic acid equivalents (GAE)

<sup>b</sup> rutin equivalent (RE)

<sup>c</sup> mmol of ferrous equivalents / g extract;

<sup>d</sup> µmol of trolox equivalents/ g extract;

BHA- Butylated hydroxyl anisole

Table 5. Correlation between phenolics, flavonoids and different antioxidant parameters of *in vitro* regenerated and wild grown plant methanol extract of *A. Montana*.

Parameters	Phenolics		Flavonoids	
	Wild grown plant extract	<i>In vitro</i> propagated plants	Wild grown plant extract	<i>In vitro</i> propagated plants
DPPH	-0.987**	0.478	-0.971**	0.669
ABTS	0.980**	0.898	0.973**	0.732
FRAP	0.992**	-0.672	0.966*	0.980**

### *Ex vitro* establishment of plantlets

The plantlets (Fig. 2f) were transferred to the potting medium containing vermiculite and covered with polythene bag. After 2 months, the cover was gradually loosened, thus dropping the humidity (65–70%). This procedure subsequently resulted in *in vitro* hardening of the plants. The survival rate was 100% when maintained in culture room condition (25 ± 2 °C). New shoot and root were formed after two months of *in vitro* hardening (Fig. 2g & h). The hardened plantlets are maintained in the National Orchidarium, Tamil Nadu, India with 80% field establishment rate.

### Total phenolics and flavonoid content of the samples

The results of the phenolics and flavonoid content of both methanol extract of the wild plant and *in vitro* regenerated plants are shown in Table 4. The results revealed that the phenolics content of the wild grown plant of *A. montana* (121.34 ± 1.23 mg/1g) was significantly ( $P < 0.05$ ) higher than that of the *in vitro* regenerated plant of *A. montana* (61.89 ± 3.19 mg/1g). Also, the flavonoid content of the methanol extracts as revealed in Table 4 indicated that wild grown plant *A. montana* (93.62 ± 1.17 mg/1g) had significantly ( $p < 0.05$ ) higher flavonoid content than *in vitro* regenerated plants (34.59 ± 1.14 mg/1g).

### Antioxidant activity

In the DPPH assay, the wild-grown plants of *A. montana* showed higher scavenging activity (IC<sub>50</sub> = 26.87 mg/ml) compared to the positive standard (BHT) (IC<sub>50</sub> = 34.59 mg/ml) and *in vitro* propagated plant (IC<sub>50</sub> = 41.45 mg/ml) (Table 4). To determine the relationship between the levels of the total phenolics and the antioxidant capacity of the extracts, correlation and regression analysis was performed. Total phenolic content of wild grown plant methanolic extract correlated with radical scavenging activity against DPPH ( $r^2 = -0.987$ ,  $P < 0.01$ ) and flavonoid ( $r^2 = -0.971$ ,  $P < 0.01$ ) are presented in Table 5 whereas, in *in vitro* regenerated plant methanolic extract, the significant correlation was not observed.



Table 6. Evaluation of antimicrobial activity of methanolic extract of *A. montana* by agar disc diffusion method and microdilution method.

Micro-organ	Inhibition zone diameters of the test compounds (mm)			Minimum inhibitory concentration (MIC, µg/ml)			
	Wild grown plant extract 25µg/disc	<i>In vitro</i> propagated plants 25µg/disc	Ciprofloxacin 10 µg/disc	Wild grown plant extract	<i>In vitro</i> propagated plants	Ciprofl oxacin	Streptomycin
<i>E. coli</i> (ATCC 25922)	8.11±0.91	4.61±0.23	21.77±0.11	31.25	125	3.90	1.90
<i>P. aeruginosa</i> (ATCC 27853)	7.11±0.34	5.54±0.21	20.47±0.17	62.5	125	1.90	3.90
<i>S. aureus</i> (ATCC 29213)	11.87±0.13	6.42±0.98	20.77±0.11	15.62	31.25	7.81	1.90
<i>S. pnumoniae</i> (ATCC 33400)	6.55±0.62	4.54±0.31	21.07±0.05	62.5	125	3.90	3.90
<i>K. pnumoniae</i> (ATCC 10031)	10.21±0.91	8.12±0.67	19.17±0.11	7.81	15.62	1.90	3.90
<i>B. subtilis</i> (ATCC 6633)	10.17±0.23	9.97±0.11	22.67±0.11	15.62	31.25	3.90	7.81

The Trolox equivalent antioxidant capacity (TEAC) was measured using the improved ABTS radical cation decolorization assay. The decolorization of ABTS<sup>•+</sup> cation radical is an unambiguous way to measure the total equivalent antioxidant capacity of test compounds or plant samples. Since, TEAC is a measurement of the effective antioxidant activity of the extract. A higher TEAC value would imply greater antioxidant activity of the sample. Similarly to DPPH assay, the wild grown plant showed the highest amount of ABTS<sup>•+</sup> radical quenching ability and then BHA and *in vitro* plant material (Table 4). From the correlation analysis, it is conceived that phenolics ( $r^2 = 0.980$ ,  $P < 0.01$ ) and flavonoids ( $0.973$ ,  $P < 0.01$ ) of *A. montana* wild grown plant methanol extract was the main contributors for their reducing activity (Table 5).

Antioxidative activity has been proposed to be related to reducing power. Therefore, the antioxidant potential of *A. montana* wild grown plant and *in vitro* propagated methanolic extract was estimated for their ability to reduce TPTZ–Fe (III) complex to TPTZ–Fe (II) (Table 4). The FRAP activities of wild-grown plant of *A. montana* ( $2987.10 \pm 19.33$  mmol Fe (II)/g extract) and *in vitro* propagated plant ( $1034.23 \pm 67.44$  mmol Fe (II)/g extract) are presented in Table 4. Similar to DPPH and ABTS radical scavenging activity, the reducing power of wild-grown plant was significantly higher than *in vitro* samples. From the correlation analysis, it is conceived that phenolics ( $r^2 = 0.992$ ,  $P < 0.01$ ) and flavonoids ( $r^2 = 0.966$ ,  $P < 0.05$ ) of wild grown plant methanolic extract of *A. montana* were the main contributors for their reducing activity (Table 5).

### Antimicrobial activity

Antibacterial activity was compared between *in vitro* propagated and wild-grown plants. The results of antibacterial assay are presented in Table 6. The result from the disc diffusion method measured in inhibition zone (IZ in mm) of *in vitro* propagated and wild-grown plants methanol extracts against bacterial strains ranged from  $4.54 \pm 0.31$  to  $22.67 \pm 0.11$  mm (Table 6). Methanol extract of wild-grown plants of *S. corymbosa* showed high antimicrobial activity against *S. aureus*, *B. subtilis* and *K. pnumoniae* at 25 µg/ml. The antimicrobial activity of *in vitro* propagated and wild-grown plants methanol extracts of *A. montana* was quantitatively assessed by MIC against the six bacterial strains at various concentration range from 500 to 3.90 µg/ml. The wild-grown plants methanol extract exhibited potent growth inhibitory activity against *K. pnumoniae*, *B. subtilis* and *S. aureus* with MIC value of 7.81 and 15.62 µg/ml respectively (Table 6).

## DISCUSSION

Tetrazolium staining indicated that *A. montana* seed viability was higher than that of observed during germination experiments. However, many studies on the Orchidaceae indicate that the viability testing often is not a good indicator of germinability (Johnson *et al.*, 2007; Mahendran and Narmatha Bai, 2009; Mahendran *et al.*, 2013). Because of this, the viability estimates should be confirmed by germination tests. Mass propagation of orchids through asymbiotic seed germination is a tool for conservation of the declining orchid propagation in nature (Kauth *et al.*, 2006; Stewart and Kane, 2006; Mahendran and Narmatha Bai, 2009; Roy *et al.*, 2011; Mahendran *et al.*, 2013). The success of asymbiotic seed germination is dependent upon identification of suitable medium and abiotic conditions. The nutritional requirements of most orchids vary due to their enormous diversity and complex mycorrhizal interactions (Arditti *et al.*, 1990). Burgeff (1959) has reported that under culture condition the seedling of saprophytic species do not develop beyond the first leaf primordia and the first root. However in the present study fully developed seedlings were regenerated on BM-TM medium supplemented with growth regulators. In *A. montana*, among the six basal media evaluated for asymbiotic seed germination, BM-TM medium was found to be the best for seed germination followed by KC medium. Though all the media favoured seed germination, the growth of the protocorm was arrested beyond protocorm stage due to browning. Browning, a problem in *in vitro* culture of orchids, results from the accumulation of phenolic compounds and causes the loss of growth capacity and tissue death during culture (Rittirat *et al.*, 2012), but can be induced by different factors. In *Grammatophyllum speciosum*, as much as 70% PLB browning occurred after 8% glucose was added as the carbon source to MS medium (Pimsen and Kanchanapoom, 2011) while in *Phalaenopsis cornu-cervi*, New Dogashima medium induced up to 30% PLB browning (Rittirat *et al.*, 2012). MS medium containing 0.5 mg/l NAA was the optimal medium for *Cymbidium faberi*, 74% of PLBs turning green and 26% of PLBs undergoing browning (Tao *et al.*, 2011). The application of TDZ (3 mg/l) in half-strength MS medium with 60-day subculture interval stimulated browning in as many as 45% of *Phalaenopsis* somatic embryos (Gow *et al.*, 2009a). In a separate study, Gow *et al.*, (2009b) found that incubation of *Phalaenopsis* leaf explants in the dark for 15 days followed by 45 days under light induced browning in as many as 90% of embryos after 60 days. In *Eria bambusifolia*, MS medium reduced PLB browning more effectively than Knudson C medium (52% and 64%, respectively) with subsequently higher percentage germination (48% and 36%, respectively) (Basker and Narmatha Bai, 2010). However, in this study, protocorms browning was presumably caused by the high salt concentration of full-strength medium. Since terrestrial orchids usually require the medium with lower salt concentrations for seed germination (Rasmussen, 1995), the diluted macro-elements of MS medium (1/2, 1/4, or 1/8 MS) was preferable in several orchid species (Lee, 1998). In the present study, half strength BM-TM medium supplemented with activated charcoal was necessary for further growth of the protocorm in *A. montana*. This promotive effect has been attributed to the ability of activated charcoal to absorb phenolic compounds released by the plantlet into the media (Pan and van Staden, 1998).

The varied responses in different media might be due to the composition of the media. All the media used in the present study have different mineral composition. BM-TM medium is enriched with organic nitrogen sources such as casin hydrolysate, amino acids and vitamins, thus conditions are suspected to be responsible for enhancing seed germination which is in agreement with many workers (Kauth *et al.*, 2006; Stewart and Kane, 2006; Roy *et al.*, 2011; Mahendran *et al.*, 2013). These findings indicate the existence of species–medium specificity. Such species – specific medium for seed germination have also been reported by many workers (Arditti and Ernst, 1993; Roy *et al.*, 2011; Mahendran *et al.*, 2013).

Orchids needauxin or cytokinin for the formation of new protocorms and plantlets development. The type and concentrations of growth regulators play an important role during *in vitro* multiplication of many orchid species (Arditti and Ernst, 1993). In *A. montana*, the protocorm developed into multiple shoots directly on half strength BM-TM medium fortified with cytokinins or cytokinin in combination with NAA. Among the cytokinins, TDZ was more effective for inducing maximum

number of multiple shoots ( $17.24 \pm 0.27$ ). TDZ also found to be suitable for protocorm proliferation for *Dendrobium* hybrids, *Dendrobium candidum*, *Satyrium nepalense* and *Phalaenopsis gigantean* (Martin and Madassery, 2006; Zhao *et al.*, 2007; Mahendran and Narmatha Bai, 2009; Niknejad *et al.*, 2011). In *A. montana* TDZ in combination with NAA enhanced the number of secondary protocorms/multiple shoots. Combined effects of cytokinin (BA or TDZ) and auxin (NAA) proved to be useful in induction of protocorm and seedling in many orchid species like *Phalaenopsis* and *Doritaenopsis* (Park *et al.*, 2002; 2003) and *Dendrobium candidum* (Zhao *et al.*, 2008). The ratio of auxin and cytokinin for the initiation of shoot buds or PLB formation varies from species to species (Teng *et al.*, 1997). Cytokinin and auxin in the ratio of 2:1 produced shoot buds in *Rhynchostylis gigantea* (Van Le *et al.*, 1999), *Vanda spathulata* (Decruse *et al.*, 2003), *Phalaenopsis* and *Renanthera imschootiana* and *Vanda coerulea* (Seeni and Latha, 2000). An auxin in combination with cytokinin has been well documented for multiplication in various species (Javed *et al.*, 2013) as observed in our study. However, the concentration of auxin required was found to be very low which indicate a high endogenous level of auxin in the explants. This may be due to reinforced by high requirement of exogenous cytokinin as the auxin has been shown to inhibit the transcription of cytokinin biosynthesis genes suggested by Tanaka *et al.* (2006) and Javed *et al.* (2013).

The organic supplements have been shown to stimulate seed germination and seedling growth of many orchids (Arditti and Ernst, 1993; Zeng *et al.*, 2012; Zhang *et al.*, 2013). However, their effects are complex and may vary depending on species, the types of explants or the developmental stages. In the present study the combined effect of organic additives with TDZ ( $6.8 \mu\text{M}$ ), various organic additives (peptone, yeast extract and coconut water) at different concentrations (0.5, 1.0, 2.0, 3.0 g/l or 5, 10, 15 and 20%) were used together with the optimal TDZ concentration (Table 3). This study was needed so as to familiarize their ability to affect the shoot induction and multiplication rate and to optimize the medium composition for maximum plantlet regeneration. Addition of organic additives (peptone, yeast extract and coconut water) at various concentrations (0.5, 1.0, 2.0, 3.0 g/l or 5, 10, 15 and 20%) in combination with the optimum concentrations of cytokinins was found to be superior for shoot initiation and multiplication. Among the different levels of organic additives tested, TDZ ( $6.8 \mu\text{M}$ ) in combination with coconut water (20%) was found to be the best for multiplication of shoots ( $28.33 \pm 0.27$ ) with a mean number of  $1.98 \pm 0.42$  shoot length per explant (Table 3). This increase in the number of multiple shoots could be attributed to the biochemical compounds of the tender coconut milk such as amino acids, organic acids, inorganic ions, vitamins, sugars, lipids, nitrogenous compounds and hormones. Similarly, the effectiveness of coconut water for promoting shoots differentiation has also been reported in *Phalaenopsis gigantean* (Murdad *et al.*, 2006), *Gastrochilus calceolaris* (Pathak *et al.*, 2011), *Vanda* Kasem's Delight (Gnasekara *et al.*, 2012) and *Dendrobium* AlyaPink (Nambiar *et al.*, 2012).

In the present study, wild-grown plant gave higher phenolic compounds and total flavonoid content probably due to a higher stress associated to their growth conditions. Otherwise, *in vitro* cultured seem to be under less stress producing lower amounts of phenolics and total flavonoids compounds (secondary metabolites). In fact, *in vitro* cultured plant grown with controlled light, temperature and nutrients. Similarly early researchers found that lower amounts of phenolic compounds *in vitro* cultured samples in *Poliomintha glabrescens* (Garcia-Perez *et al.*, 2012), *Hypericum* spp (Danova *et al.*, 2012), *Cichorium pumilum* (Khateeb *et al.*, 2012), *Hypericum undulatum* (Rainha *et al.*, 2013), *Melissa officinalis* (Barros *et al.*, 2013) and *Swertia corymbosa* (Mahendran and Narmatha Bai, 2014).

Free-radical scavenging activity (FRSA) was determined to evaluate the antioxidant potential of *in vitro* propagated plants and these were compared with levels in wild-grown plants. Considering the fact that the mechanisms of antioxidant processes are complex, therefore, an approach with multiple assays in screening work is highly advisable (Matkowski, 2008; Amoo *et al.*, 2012; Mahendran and Narmatha Bai, 2014). In the present study, various methods of *in vitro* assays were performed to determine the antioxidant activity of *in vitro* propagated plants and wild grown plants of *A. montana*.

The DPPH free radical scavenging assay has been widely used to evaluate antioxidant capac-

ities. Antioxidants react with DPPH, reducing a number of DPPH molecules equal to the number of available hydroxyl groups (Matthaus, 2002). The degree of discoloration indicates that the samples to scavenge DPPH radical due to its ability to donate hydrogen proton. Similar to DPPH, the decolorization of ABTS<sup>•+</sup> radical reflects the capacity of an antioxidant species to donate electrons or hydrogen atoms to inactivate this radical species. The ABTS<sup>•+</sup> radical cation is generated from the reaction of ABTS with potassium persulfate overnight in water (Re *et al.*, 1999). In the present study, the methanolic extract of plants growing wild seemed to exhibit a higher antioxidant potential compared to plants growing *in vitro* under controlled conditions. The differences between the responses of methanolic extracts prepared from wild and *in vitro* growing explants could be attributed to the differences in the chemical constituents of the plant cells that are likely to develop in response to the surrounding environmental conditions and different components are accumulated during different growth phases or due to mutations (Parsaeimehr *et al.*, 2010; Mohanty *et al.*, 2011). The radical scavenging activity of extracts can be credited to the presence of its major phenolic compounds (Guimaraes *et al.*, 2010). The antioxidant activity of phenolic compounds is related to the hydroxyl groups linked to the aromatic ring, which are capable of donating hydrogen atoms with electrons and stabilizing free radicals (Dorman *et al.*, 2003; Yanishlieva *et al.*, 2006). In *P. glabrescens*, *C. pumilum*, *M. officinalis* and *S. corymbosa* higher antioxidant activity correlated with higher total phenolic contents (Garcia-Perez *et al.*, 2012; Khateeb *et al.*, 2012; Barros *et al.*, 2013; Mahendran and Narmatha Bai, 2014).

## CONCLUSION

In conclusion, the present study is the first to report seed derived protocorm explants and the successful plant regeneration in *A. montana*. The protocol described here could be applied in a propagation program for genetic resource conservation and commercial purposes. The total phenolics, flavonoid content and antioxidant activity of *in vitro* propagated plants methanol extract were lower than that of wild-grown plants' methanol extract. In addition, we provide evidence that anti-bacterial activity of *A. montana* extracts revealed that wild-grown plants' methanol extract at 25 µg/ml exhibited strong antibacterial activity against *S. aureus*, *B. subtilis* and *K. pneumoniae* with MIC values of 7.81 and 15.62 µg/ml respectably. Further work on the type of phytoconstituents and purification of individual groups of bioactive components can reveal the exact potential of the plant to inhibit pathogenic microbes and induce the antioxidation. Tissue culturing and the *in vitro* propagation of *Aphyllorchis* explants could be a possible method for the large scale commercial production of biologically active components.

## ACKNOWLEDGEMENTS

The financial support by University Grants Commission (UGC-F.No-37-97/2009(SR)), New Delhi is gratefully acknowledged.

## Literature Cited

- Amoo, S.O., Aremu, A.O. and Van Staden, J. 2012. *In vitro* plant regeneration, secondary metabolite production and antioxidant activity of micropropagated *Aloe arborescens* Mill. Plant Cell, Tissue and Organ Culture, 111: 345–358.
- Arditti, J. and Ernst, R. 1993. Micropropagation of orchids. John Wiley and Sons. New York.
- Arditti, J., Ernst, R., Yam, T.W. and Glabe, C. 1990. The contribution to orchid mycorrhizal fungi to seed germination: A speculative review. Lindleyana, 5: 249-255.
- Barros, L., Duenas, M., Dias, M.I., Sousa, M.J., Santos-Buelga, C. and Ferreira, I.C.F.R. 2013. Phenolic profiles of cultivated, *in vitro* cultured and commercial samples of *Melissa officinalis* L. infusions. Food Chemistry, 136: 1–8.
- Basker, S., and Narmatha Bai V. 2010. *In vitro* propagation of an epiphytic and rare orchid *Eria bambusifolia* Lindl. Research in Biotechnology, 1: 15–20.

- Bhavani, P., Sreenu, T., Tharangini, K., Geethanjali, J., Govinda Reddy, T. and Venkata Lakshmi, D. 2012. Hypoglycemic activity of ethanolic extract of *Aphyllorchis montana* on alloxan induce diabetes in rats. *International Journal of Chemistry and Pharmaceutical Sciences*, 3(3): 27-29.
- Blois, M.S. 1958. Antioxidant determinations by the use of a stable free radical. *Nature* 26: 1199-1200.
- Burgeff, H. 1959. Mycorrhiza of orchids, pp. 361-395. In: Withner C.L. (ed), *The Orchids: A Scientific Survey*, The Ronald Press Company, New York.
- Danova, K., Nikolova-Damianova, B., Denev, R., and Dimitrov, D. 2012. Influence of vitamins on polyphenolic content, morphological development, and stress response in shoot cultures of *Hypericum* spp. *Plant Cell, Tissue and Organ Culture*, 110: 383–393.
- Decruse, S.W., Gangaprasad, A., Seeni S. and Menon V.S. 2003. Micropropagation and ecorestoration of *Vanda spathulata*, an exquisite orchid. *Plant Cell, Tissue and Organ Culture*, 72: 199-202.
- Dorman, H.J.D., Peltoketo, A., Hiltunen, R. and Tikkanen, M.J. 2003. Characterisation of the antioxidant properties of deodorized aqueous extracts from selected lamiaceae herbs. *Food Chemistry*, 83: 255–262.
- Eloff, J.N. 1998. A sensitive and quick microplate method to determine the minimal inhibitory concentration of plant extracts for bacteria. *Planta Medica*, 64: 711–713.
- Garcia-Perez, E., Gutierrez-Urbe, J.A. and Garcia-Lara, S. 2012. Luteolin content and antioxidant activity in micropropagated plants of *Poliomintha glabrescens* (Gray). *Plant Cell, Tissue and Organ Culture*, 108: 521–527.
- Gnasekaran, R., Poobathy, R., Mahmood, M., Samian, M.R. and Subramaniam S. 2012. Effects of complex organic additives on improving the growth of PLBs of *Vanda Kasem's Delight*. *Australian Journal of Crop Science*, 6: 1245-1248.
- Gow, W.P., Chen, J.T. and Chang, W.C. 2009a. Enhancement of direct somatic embryo-genesis and plantlet growth from leaf explants of *Phalaenopsis* by adjusting culture period and explant length. *Acta Physiologiae Plantarum*, 32: 621–627.
- Gow, W.P., Chen J.T. and Chang, W.C. 2009b. Effects of genotype, light regime, explants position and orientation on direct somatic embryogenesis from leaf explants of *Phalaenopsis orchids*. *Acta Physiologiae Plantarum*, 31: 363–369.
- Guimaraes, R., Sousa, M.J. and Ferreira, I.C.F.R. 2010. Contribution of essential oils and phenolics to the antioxidant properties of aromatic plants. *Industrial Crops and Products*, 32: 152–156.
- Javed, S.B., Anis, M., Khan P.R. and Aref, I.M. 2013. *In vitro* regeneration and multiplication for mass propagation of *Acacia ehrenbergiana* Hayne: a potential reclament of denude arid lands. *Agroforestry Systems*, 87: 621–629.
- Johnson, T.R., Stewart, S.L., Daniela, D., Kane, M.E. and Richardson, L. 2007. A symbiotic and symbiotic seed germination of *Eulophia alta* (orchidaceae) - Preliminary evidence for the symbiotic culture advantage. *Plant Cell, Tissue and Organ Culture*, 90: 313-323.
- Joshi, S.C., Verma, A.R. and Mathela, C.S. 2010. Antioxidant and antibacterial activities of the leaf essential oil of himalayan lauraceae species. *Food and Chemical Toxicology*, 48: 37–40.
- Kauth, P.J., Vendrame, W.A. and Kane, M.E. 2006. *In vitro* seed culture and seedling development of *Calopogon tuberosus*. *Plant Cell, Tissue and Organ Culture*, 85: 91–102.
- Khateeb, W.A., Hussein, E., Qouta L., Aludatt, M., Al-Shara, B. and Abu-zaiton, A. 2012. *In vitro* propagation and characterization of phenolic content along with antioxidant and antimicrobial activities of *Cichori umpumilum* Jacq. *Plant Cell, Tissue and Organ Culture*, 110: 103–110.
- Knudson, L. 1946. A new nutrient solution for the germination of orchid seed. *American Orchid Society Bulletin*, 15: 214–217.
- Lee, Y.I. 1998. The study of embryo development and seed germination *in vitro* in slipper orchids. M.Sc thesis. National Taiwan University, Taiwan.
- Lindemann, E.G.P., Gunckel, J.E. and Davidson, O.W. 1970. Meristem culture of *Cattleya*. *American Orchid Society Bulletin*, 39: 1002-1004.

- Mahendran, G., Muniappan, V., Ashwini, M., Muthukumar, T. and Narmatha Bai, V. 2013. Asymbiotic seed germination of *Cymbidium bicolor* Lindl. (Orchidaceae) and the influence of mycorrhizal fungus on seedling development. *Acta Physiologiae Plantarum*, 35: 829-840.
- Mahendran, G. and Narmatha Bai, V. 2009. Mass propagation of *Satyrium nepalense* D. Don- A medicinal orchid via seed germination. *Scientia Horticulturae*, 119: 203-207.
- Mahendran, G. and Narmatha Bai, V. 2014. Micropropagation, antioxidant properties and phytochemical assessment of *Swertia corymbosa* (Griseb.) Wight ex C. B. Clarke: a medicinal plant. *Acta Physiologiae Plantarum*, 36: 589-603.
- Martin, K.P. and Madassery, J. 2006. Rapid *in vitro* propagation of *Dendrobium* hybrids through direct shoot formation from foliar explants, and protocorm-like bodies. *Scientia Horticulturae*, 108: 95-99.
- Matkowski, A. 2008. Plant *in vitro* culture for the production of antioxidants – A review. *Biotechnology Advances*, 26: 548-560.
- Matthaus, B. 2002. Antioxidant activity of extracts obtained from residues of different oil seeds. *Journal of Agricultural and Food Chemistry*, 50: 3444-3452.
- Mitra, G.C., Prasad, R.N. and Chowdhury, A.R. 1976. Inorganic salts and differentiation of protocorms in seed callus of an orchid and correlated changes in its free amino acid content. *Indian Journal of Experimental Biology*, 14: 350-351.
- Mohanty, S., Parida, R., Singh, S., Joshi, R., Subudhi, E. and Nayak, S. 2011. Biochemical and molecular profiling of micropropagated and conventionally grown *Kaempferia galanga*. *Plant Cell, Tissue and Organ Culture*, 106: 39-46.
- Morel, G.M. 1965. Clonal propagation of orchids by meristem culture. *Cymbidium Society News*, 20: 3-11.
- Murashige, T. and Skoog, F. 1962. A revised medium for rapid growth and bio-assays with tobacco tissue cultures. *Physiologia Plantarum*, 15: 473-497.
- Murdad, R., Hwa, K.S., Seng, C.K., Latip, M.A., Aziz, Z.A. and Ripin, R. 2006. High frequency multiplication of *Phalaenopsis gigantea* using trimmed bases protocorms technique. *Scientia Horticulturae*, 111: 73-79.
- Nambiar, N., Tee, C.S. and Maziah, M. 2012. Effects of organic additives and different carbohydrate sources on proliferation of protocorm like bodies in *Dendrobium* Alya Pink. *Plant Omics Journal*, 5: 10-18.
- Niknejad, A., Kadir, M.A. and Kadzimin, S.B. 2011. *In vitro* plant regeneration from protocorms-like bodies (PLBs) and callus of *Phalaenopsis gigantea* (Epidendroideae: Orchidaceae). *African Journal of Biotechnology*, 10: 11808-11816.
- Pan, M.J. and van Staden, J. 1998. The use of charcoal in *in vitro* culture – A review. *Plant Growth Regulation*, 26: 155-163.
- Park, S.Y., Murthy, H.N. and Paek, K.Y. 2002. Rapid propagation of *Phalaenopsis* from floral stalk-derived leaves. *In vitro Cellular & Developmental Biology – Plant*, 38: 168-172.
- Park, S.Y., Murthy, H.N. and Paek, K.Y. 2003. Protocorm-like body induction and subsequent plant regeneration from root tip cultures of *Doritaenopsis*. *Plant Science*, 164: 919-923.
- Parsaeimehr, A., Sargsyan, E. and Javidnia, K. 2010. A comparative study of the antibacterial, antifungal and antioxidant activity and total content of phenolic compounds of cell cultures and wild plants of three endemic species of *Ephedra*. *Molecules* 15: 1668-1678.
- Pathak, P., Piri, H., Vij, S.P., Mahant, K.C. and Chauhan, S. 2011. *In vitro* propagation and mass scale multiplication of a critically endangered epiphytic orchid, *Gastrochilus calceolaris* (Bhch.-Ham ex J.E.Sm) D. Don using immature seeds. *Indian Journal of Experimental Biology*, 49: 711-716.
- Pimsen, M., and Kanchanapoom, K. 2011. Effect of basal media and sugar types on *in vitro* regeneration of *Grammatophyllum speciosum* Blume. *Notulae Scientia Biologicae*, 3(3): 101-104.

- Prajapati, N.S., Purohit, S.S., Sharma, A.K. and Kumar, T. 2003. A handbook of medicinal plants. A complete source book. Agrobios.
- Pulido, R., Bravo, L. and Sauro-Calixto, F. 2000. Antioxidant activity of dietary polyphenols as determined by a modified ferric reducing/antioxidant power assay. *Journal of Agricultural and Food Chemistry*, 48: 3396-3402.
- Rainha, N., Koci, K., Coelho, A.V., Lima, E., Baptista, J. and Ferreira, F. 2013. HPLC–UV–ESI-MS analysis of phenolic compounds and antioxidant properties of *Hypericum undulatum* shoot cultures and wild-growing plants. *Phytochemistry*, 86: 83–91.
- Rasmussen, H.N. 1995. Terrestrial orchids: from seed to mycotrophic plant. Cambridge University Press., Cambridge.
- Re, R., Pellegrini, N., Proteggente, A., Pannala, A., Yang, M. and Rice- Evans, C. 1999. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Biology and Medicine*, 26: 1231-1237.
- Rittirat, S., Thammasiri, K. and Te-chato, S. 2012. Effect of media and sucrose concentrations with or without activated charcoal on the plantlet growth of *Phalaenopsis cornu-cervi* (Breda) Blume & Rchb. f. *International Journal of Agricultural Technology*, 8(6): 2075–2085.
- Roy, A.R., Patel, R.S., Patel, V.V., Sajeev, S. and Deka, B.C. 2011. Asymbiotic seed germination, mass propagation and seedling development of *Vanda coerulea* Griff ex. Lindl. (BlueVanda): An *in vitro* protocol for an endangered orchid. *Scientia Horticulturae*, 128: 325-331.
- Seeni, S. and Latha, P.G. 2000. *In vitro* multiplication and eco rehabilitation of the endangered Blue *Vanda*. *Plant Cell, Tissue and Organ Culture*, 61: 1–8.
- Siddhuraju, P. and Becker, K. 2003. Studies on antioxidant activities of *Mucuna* seed (*Mucunapruriens* var. *utilis*) extracts and certain non-protein amino/imino acids through *in vitro* models. *Journal of the Science of Food and Agriculture*, 83: 1517-1524.
- Siddiqi, R., Naz, S., Ahmad, S. and Sayeed, S.A. 2011. Antimicrobial activity of the polyphenolic fractions derived from *Grewia asiatica*, *Eugenia jambolana* and *Carissa carandas*. *Journal of Food Science and Technology*, 46: 250–256.
- Sinu, P.A., Sinu, N. and Chandrashekar, K. 2012. Ecology and population structure of a terrestrial mycoheterotrophic orchid, *Aphyllorchis montana* Rchb.f. (Orchidaceae) in soppinabetta forests of the western ghats. *Journal of Threatened Taxa*, 4(9): 2915–2919.
- Sreenu, T., VenkataRamana, K., Jyothisbasu, T. and Delhiraj, N. 2013. Hypoglycemic activity of ethanolic extract of *Aphyllorchis montana* induced by streptozocin in rats. *International Journal of Pharmacy Technology Research*, 5: 320-323.
- Stewart, S.L. and Kane, M.E. 2006. Symbiotic germination of *Habenariama croceratitis* (Orchidaceae), a rare Florida terrestrial orchid. *Plant Cell, Tissue and Organ Culture*, 86: 159–167.
- Tanaka, M., Takei, K., Kojima, M., Sakakibara, H. and Mori, H. 2006. Auxin controls local cytokinin biosynthesis in the nodal stem in apical dominance. *The Plant Journal*, 45(6): 1028–1036.
- Tao, J., Yu, L., Kong, F. and Zhao, D. 2011. Effects of plant growth regulators on *in vitro* propagation of *Cymbidium faberi* Rolfe. *African Journal of Biotechnology*, 10(69): 15639–15646.
- Teng, W.L., Nicholson, L. and Teng, M.C. 1997. Micropropagation of *Spathoglottis plicata*. *Plant Cell Reports*, 16:831-835.
- Van Le, B., Hang Phuong, N.T., Anh Hong, L.T. and Tran Thanh Van, K. 1999. High frequency shoot regeneration from *Rhynchostylis gigantea* (Orchidaceae) using thin cell layers. *Plant Growth Regulation*, 28: 179-185.
- Van Waes, J.M. and Debergh, P.C. 1986. *In vitro* germination of some western european orchids. *Physiologia Plantarum*, 67: 253-261.
- Vellupillai, M., Swaru, P.S. and Goh, C.J. 1997. Histological and protein changes during early stage of seed germination in the orchid *Dendrobium curmenatum*. *Journal of Horticultural Sciences*, 76: 941-948.
- Yanishlieva, N.V., Marinova, E. and Pokorny, J. 2006. Natural antioxidants from herbs and spices.

European Journal of Lipid Science and Technology,108: 776–793.

- Zeng, S., Wu, K., Teixeira da Silva, J.A., Zhang, J., Chen, Z., Xia, N. and Duan, J. 2012. Asymbiotic seed germination, seedling development and reintroduction of *Paphiopedilum wardii* Sumerh., an endangered terrestrial orchid. *Scientia Horticulturae*,138: 198–209.
- Zhang, Y., Lee, Y.L., Deng, L. and Zhao, S. 2013. Asymbiotic germination of immature seeds and the seedling development of *Cypripedium macranthos* Sw., an endangered lady's slipper orchid. *Scientia Horticulturae*, 164: 130–136.
- Zhao, P., Wang, W., Feng, F.S., Wu, F., Yang, Z.Q. and Wang, W.J. 2007. High frequency shoot regeneration through transverse thin cell layer culture in *Dendrobium candidum* Wall. Ex. Lindl. *Plant Cell, Tissue and Organ Culture*, 90: 131-139.
- Zhao, P., Wu, F., Feng, F.S. and Wang, W.J. 2008. Protocorm-like body (PLB) formation and plant regeneration from the callus culture of *Dendrobium candidum* Wall ex Lindl. *In Vitro Cellular & Developmental Biology – Plant*, 44: 178-185.
- Zhishen, J., Mengcheng, T. and Jianming, W. 1999. The determination of flavonoid contents in mulberry and their scavenging effects on superoxide radicals. *Food Chemistry*, 64: 555-559.

Archive of SID



# یک تکثیر درون شیشه‌ای کارآمد همراه با فعالیتهای آنتی‌اکسیدانی و ضد میکروبی گیاه *Aphyllorchis Montana* (Reichenb.f.)

گانشان ماهندران

محقق گروه گیاهشناسی دانشگاه بهاراتیار، هندوستان

تاریخ تایید: ۲۹ آبان ۱۳۹۳

تاریخ دریافت: ۲۱ خرداد ۱۳۹۳

\* ایمیل نویسنده مسئول: mahendran0007@gmail.com

یکی دستورالعمل باززایی درون شیشه‌ای برای گیاه *Aphyllorchis Montana* (Reichenb.f.) که یک ارکید بدون کلروفیل اسپورفیت است، از طریق کشت بذرهای نابالغ به صورت موفقیت آمیز انجام شد. در بین ۶ محیط کشت پایه که برای جوانه زنی بذرهای ارزیابی شد، محیط کشت BM-TM و سپس KC بهترین بودند. ۴۰ روز بعد، همه‌ی محیط‌های کشت قهوه‌ای شده و رشد پیش‌پدازه‌ها متوقف شد. زغال فعال به مقدار یک گرم در لیتر در محیط کشت ۵۰٪ BM-TM برای نمو بهتر پیش‌پدازه‌ها بهتر بود. این محیط کشت با تنظیم کننده‌های رشد مختلف به تنهایی یا در ترکیب با هم برای تکثیر شاخساره‌ها تکمیل شد. از بین ۵ سیتوکنین آزمایش شده، TDZ در غلظت ۶/۸ میکرومول برای القای تکثیر شاخساره موثرترین بود و پس از ۱۰ روز  $17/24 \pm 0/27$  شاخساره تولید شد. افزودن مقدار کمی NAA (۱/۳ میکرومول) به محیط کشت MS تکمیل شده با سیتوکنین TDZ (۶/۸ میکرومول) برای شاخه‌زایی مطلوب بود. میانگین  $27/56 \pm 0/54$  شاخساره با  $3/92 \pm 0/11$  ریشه در هر نمونه گیاهی بدست آمد. واکنش بذرهای حاصل از پیش‌پدازه‌ها به انواع مختلف ترکیبات آلی اضافه شده مثل پیتون، عصاره مخمر و آب نارگیل نیز ارزیابی شد. افزودن این ترکیبات به محیط کشت حاوی TDZ تعداد شاخساره را افزایش داد. گیاهچه‌ها در گلدان‌های پلاستیکی حاوی ورمی‌کولیت استریل شده سازگار شدند. میزان بقاء گیاهچه‌ها در اتاق کشت ۲۵ درجه سانتی‌گراد، صد در صد بود. ارزیابی فعالیت ضد باکتریایی و آنتی‌اکسیدانی و تخمین مواد فنولیک و مقدار فلاونوئید عصاره متانولی گیاهان ریز ازدیاد شده نیز انجام و با گیاهان خودرو مقایسه شد. در همه آزمایش‌ها، عصاره متانولی گیاهان خودرو مقادیر بیشتری از ترکیبات فوق نسبت به گیاهان درون شیشه‌ای داشتند.

چیتیه

**کلید واژگان:** فعالیت ضد باکتریایی، فعالیت آنتی‌اکسیدانی، *Aphyllorchis Montana* (Reichenb.f.)، جوانه‌زنی غیرهمزیستی بذر، تکثیر درون شیشه‌ای، تنظیم کننده‌های رشد، ۲،۲-دی‌فنیل-۱-بیکریل‌هیدرازیل.

www.SID.ir