

Effect of Foliar Application of Silicon on Physiological Responses of Chrysanthemum (*Dendranthema ×grandiflorum*) at Two Different Growth Stages

Hadi Hajipour and Zohreh Jabbarzadeh*

Department of Horticultural Science, College of Agriculture, Urmia University, P.O. Box 165, Iran

Received: 26 October 2015

Accepted: 31 December 2015

*Corresponding author's email: z.jabbarzadeh@urmia.ac.ir

Abstract

To evaluate the physiological responses of chrysanthemum to silicon spray in non-stress conditions, this study was performed in two separate experiments (at 4 to 5 and 8 to 12-leaf stages) as a factorial randomized completely design with two factors: silicon source (in two levels: sodium silicate and calcium silicate) and the concentration of silicate at five levels (0, 50, 100, 150 and 200 mg L⁻¹) in cocopeat- perlite (1:1 v/v) medium in four replications and two observations. Physiological traits such as catalase (CAT), ascorbate peroxidase (APX), guaiacol peroxidase (GPX) activity, proline, malondialdehyde (MDA) and hydrogen peroxide (H₂O₂) content were measured. The results showed that the foliar spray of sodium and calcium silicate increased the activity of CAT, APX and GPX enzymes and MDA, but proline and H₂O₂ were decreased. Generally, in the stage of 8-12 leaves, sodium silicate at high concentrations (150 and 200 mg L⁻¹) was the most effective in increasing antioxidant enzymes activity and reducing proline, MDA and H₂O₂. At 4-5 leaf stage sodium silicate at 100 mg L⁻¹ increased antioxidant enzymes activity and calcium silicate at 150 mg L⁻¹ reduced proline, MDA and H₂O₂.

Keywords: Ascorbate peroxidase, Calcium silicate, Catalase, Guaiacol peroxidase, Hydrogen peroxide, Sodium silicate.

INTRODUCTION

Chrysanthemum (*Dendranthema × grandiflorum*) is one of the most important ornamental plants and its commercial production is often affected by biotic and abiotic stresses (Zhang *et al.*, 2005).

After oxygen, silicon is the second most abundant element in the crust of the earth. Plants can use only the Si(OH)₄ (Epstein, 1999). Silicon causes resistance to pathogens, biotic and abiotic stresses. Biological stresses cause the accumulation of ROS (Reactive Oxygen Species) including superoxide radicals (O₂⁻), hydroxyl radicals (•HO) and hydrogen peroxide (H₂O₂) in plants (Molassiotis *et al.*, 2006). Antioxidant enzymes such as guaiacol peroxidase (GPX), ascorbate peroxidase (APX) and catalase (CAT) seems to play an important role in reducing the harmful effects of ROS (Li-Ping *et al.*, 2006). Silicon can cause modulation of the oxidative stress, enhance membrane stability and reduce the electrolyte leakage (Karmollachaab *et al.*, 2013). A positive relationship between silicon concentration and activity of CAT and APX has been reported in plants (Mohaghegh *et al.*, 2011). Catalase is a major enzyme that can scavenge ROS and inhibit lipid peroxidation, while in plant cells can control the levels of hydrogen peroxide and the damage to cell membranes and chlorophyll. Catalase can control hydrogen peroxide levels in plant cells and seems to play an important role in the photosynthesis process (Khelifa *et al.*, 2011). APX is belonging to a group of peroxidases that have been found in chloroplasts, microbodies and cytosols and its main role is to remove the hydrogen peroxide. APX is protecting the plant cells against oxidative stress by using ascorbate as an electron donor (Ghamsari *et al.*, 2007; Rosa *et al.*, 2002). GPX is considered as a ROS scavenger because of its tendency to destroy hydrogen peroxide better than catalase (Brigelius-Flohe and Flohe, 2003). MDA is a breakdown product of unsaturated fatty acids and is widely used as a parameter for assay of lipid peroxidation (Mittler, 2002). It is estimated that more than 75% of the MDA is derived from α-Linoleic acid (Weber *et al.*, 2004). Accumulation of proline is stimulated in response to environmental stress, osmotic conditions, aging and other factors. Proline plays an important role in osmoregulation (Ahmad and Hellebust, 1988). The aim of this study was to investigate the role of silicon, by foliar application, on the physiological and antioxidant response of chrysanthemum.

MATERIALS AND METHODS

Plant material and growth conditions

This study was performed in two separate experiments. Every experiment had four replications and two observations. In the first experiment (at 4-5-leaf stage of plant growth cycle), there were two factors: the first factor was the two silicone source (sodium silicate and calcium silicate) and the second factor was the concentration of silicate at five levels (0, 50, 100, 150 and 200 mg). The second experiment was done at 8-12-leaf stage with the same treatments as in the first experiment. The medium that was used for the research was a mixture of cocopeat and perlite (1:1 v/v). Plants were grown in a greenhouse in 2 L pots in cocopeat-perlite (1:1 v/v) medium under 17/14°C (day/night) and light intensity of 55-75 μmol m⁻²s⁻¹. Sodium and calcium silicate foliar application was done weekly for 10 weeks. During this period the standard Hoagland nutrient solution (1/4 concentration) was added weekly to the media.

Enzyme extraction and assay

Preparation of plant extracts for determination of CAT, APX and GPX activity was carried out based on Kang and Saltveit protocol (2002) with slight modifications. 0.5 g of leaves was placed into a cold mortar/pestle. Enzyme extraction was done with 3 ml of Tris buffer (pH =7.5) that contained 3 mM and 1 mM EDTA (and extraction buffer for measuring the activity of APX was consisted of 0.2 mM ascorbate). The homogenates were centrifuged for 20 min at 4°C at 5000 rpm. The supernatant was used as a crude extract for measuring the activity of antioxidant enzymes.

The activity of CAT was measured according to Aebi (1984), APX according to Nakano and Asada (1981) and the GPX according to the protocol of Upadhyaya *et al.* (1985).

To determine the proline content, 0.5 g of fresh leaves were homogenized with 5 ml of 95% ethanol. The upper phase of filtrate was separated and its sediments were washed with 5 ml of 70% ethanol twice and its upper phase added to the previous one. The mixture was centrifuged at 3500 g for 10 min at 4°C and the supernatant was recovered and the alcoholic extract kept at 4°C (Paquin and Lechasseur, 1979). One ml of alcoholic extract was diluted with 10 ml of distilled water and 5 ml of ninhydrin (0.125 g ninhydrin, 2 ml of 6 mM NH₃PO₄, 3 ml of glacial acetic acid) and 5 ml of glacial acetic acid was added. The mixture was placed in boiling water bath for 45 min at 100°C. The reaction was stopped by placing the test tubes in cold water. The samples were vigorously mixed with 10 ml benzene. The light absorption of benzene phase was measured at 515 nm.

Lipid peroxidation in the leaf tissues was determined in terms of malondialdehyde (MDA) content by the thiobarbituric acid (TBA) method, as described by Popham and Novacky (1990). Briefly, 0.2 g of leaf tissue were homogenized in 5 ml of 1% (w: v) trichloroacetic acid (TCA) and then centrifuged at 8000 g for 10 min. One ml of the supernatant was added in to 4 ml of 20 % TCA that contained 0.5 % TBA and the solution was heated for 30 min at 95 °C in water bath. The samples were cooled on ice for 5 min and re-centrifuged for 5 min at 8000 g. Absorbance was measured at 532 nm. Malondialdehyde was calculated according to the following formula and extinction coefficient, mM⁻¹Cm⁻¹ 155.

$$\text{MDA } (\mu\text{mol/g FW}) = [\text{A}_{532} - \text{A}_{600}/155] \times 1000$$

Hydrogen peroxide levels were determined according to the Velikova *et al.*, (2000) protocol. 0.1 g fresh weight of leaf tissue was homogenized in 5 ml of 0.1% TCA solution. After centrifugation at 12000 g for 15 min, 0.5 ml of the supernatant was added to the reaction mixture containing 0.5 ml of 10mM potassium phosphate buffer (pH=7.0) and 1 ml of 1M potassium iodide (KI). The reaction mixture was placed at room temperature in the dark for one hour and then the absorbance was determined at 390 nm.

Statistical analysis of experimental data was performed using SAS 9.1 software and the means were compared with Duncan's multiple range test (DMRT) at 5% probability level.

RESULTS AND DISCUSSION

Catalase activity

At both growth stages, the highest activity of CAT was related to sodium silicate. The results showed that 200 and 100 mg L⁻¹ silicon were most effective in increasing catalase activity, at 8-12 and 4-5-leaf stage, respectively (Fig. 1 A, B). In 8-12-leaf stage at all levels catalase activity in sodium silicate was higher than calcium silicate. In 8-12-leaf stage with increasing sodium silicate CAT activity was also higher so that the highest catalase activity was observed at 200 mg L⁻¹ with 8.574 mM/min. Generally, for increasing catalase activity greater concentration of sodium silicate is needed at 8-12-leaf stage than at 4-5-leaf stage.




Fig. 1. Effect of sodium and calcium silicate at 8- 12 (A) and 4- 5-leaf stage (B) on catalase activity. Means with the same letter are not significantly different at 5% level of Duncan's test.




Fig. 2. Effect of sodium and calcium silicate at 8- 12 (A) and 4- 5-leaf stage (B) on ascorbate peroxidase activity. Means with the same letter are not significantly different at 5% level of Duncan's test.

Silicon may increase CAT by affecting catalase production path in peroxisomes. Increasing of CAT production by application of silicon provides conditions for the activation of another antioxidant enzyme such as APX and GPX. Silicon also can increase catalase activity in non-stress conditions. Catalase is able to convert hydrogen peroxide into water and oxygen. Silicon increased catalase activity in wheat (Tale Ahmad and Haddad, 2011), cotton (Moldes *et al.*, 2013) and tomato (Al-Aghabary *et al.*, 2004). Catalase activity in plants treated with silicon was 29.2% higher than in silicon-deprived plants (Gang *et al.*, 2008). Liang *et al.* (1996) showed that in barley (salt sensitive and resistant cultivars), silicon increased activity of CAT, SOD and peroxidase in roots.

Ascorbate peroxidase activity

At 8-12-leaf stage calcium silicate was more effective than sodium silicate in increasing APX activity (Fig. 2A) but at 4-5-leaf stage sodium silicate was more effective than calcium silicate (Fig. 2B). By application of sodium silicate at both stages, highest APX rate was obtained at 100 mg L⁻¹. At 4-5-leaf stage the maximum APX amount was obtained at 100 mg L⁻¹ sodium silicate with 5.192 mM/min. The most effective concentrations at the 8-12 and 4-5-leaf stages were 200 and 100 mg L⁻¹, respectively.

Silicon likely acted, in the presence of reactive oxygen species, as signal to increase the production of ascorbate peroxidase. Thus, the APX activity was increased by ROS signals; it also may play a role in the suppression of cellular ROS. CAT tendency to hydrogen peroxide (mM) is lower than ascorbate peroxidase (μM). So, APX activity in non-stress conditions; which rate of hydrogen peroxide is little; is more than CAT. APX may act as a signal in the regulation of ROS. APX is the first enzyme in the pathway to remove hydrogen peroxide and accelerated hydrogen peroxide convert into the water. Activity of APX enhanced production of other antioxidant enzymes such as CAT, SOD and glutathione reductase (GR). Application of silicon can increase antioxidant enzymes activity in *Helianthus annuus* (Moghadam *et al.*, 2013), soybean (Miao *et al.*, 2010) and wheat (Gharineh and Karmollachaab, 2013). Results of Moldes *et al.* (2013) showed that potassium silicate treatments increased GPX, CAT and APX activity in cotton, in hydroponic systems.

Guaiacol peroxidase activity

The highest level of GPX activity at 8-12 and 4-5-leaf stage was observed in sodium and calcium silicate, respectively (Fig. 3A, B). At 8-12-leaf stage, GPX activity, with sodium silicate at all levels was higher than with calcium silicate. At 4-5-leaf stage, there was no significant difference in GPX activity among sodium silicate concentrations, but in the case of calcium silicate 50 and 100 mg L⁻¹ were more effective than the others. GPX is one of the important enzymes that can protect the plant against oxidative stress. Silicon treatment increased SOD, GPX, CAT and POD activity compared to control in sunflower (Moghadam *et al.*, 2013). Silicon increases the antioxidant enzymes such as catalase and peroxidase and decreases hydrogen peroxide and malondialdehyde concentrations (Al-Aghabary *et al.*, 2004). Another study showed that silicon increases




Fig. 3. Effect of sodium and calcium silicate in 8- 12 (A) and 4- 5-leaf stage (B) on guaiacol peroxidase activity. Means with the same letter are not significantly different at 5% level of Duncan's test.

the amount of SOD, CAT, GPX and APX and reduces H_2O_2 and MDA in linen grown in hydroponic conditions (Bharwana *et al.*, 2013).

Proline

At 8-12 and 4-5-leaf stages lowest proline levels (0.420 and 0.696 $mg\ g^{-1}$), were achieved by sodium and calcium silicate, respectively (Fig. 4A, B). At 8-12-leaf stage at all levels of silicon, proline content in sodium silicate was less than in calcium silicate while at the 4-5-leaf stage the opposite was observed. Proline amount was decreased at both stages with 150 $mg\ L^{-1}$ sodium and calcium silicate and again increased with 200 $mg\ L^{-1}$ of both.

Silicon reduces stress effects and activates antioxidant systems in plant. It can reduce the amount of hydrogen peroxide and malondialdehyde that provide the context for proline decrease. Foliar spray of Actisil Hydro Plus decreased free proline in Solomon's seal (*Polygonatum multiflorum*) in non-stress conditions (Rubinowska *et al.*, 2014). Similar results of free proline reduction were obtained silicon treatments Kaya *et al.* (2006) and Tuna *et al.* (2008). Proline content was decreased significantly by application of sodium silicate in bean (Muhammad and Shaheed, 2012). Silicon at 215 mM in tomato and 5 mM in spinach caused a significant decrease in MDA, membrane permeability, proline and hydrogen peroxide (Hsieh *et al.*, 2002).

Malondialdehyde

At 8-12-leaf stage, silicon reduced MDA at low concentrations but at 4-5-leaf stage at high concentrations. In both 8-12 and 4-5-leaf stages highest MDA level was found in control (0.546 and 0.669 $\mu mol\ g^{-1}$ FW, respectively). At 4-5-leaf stage and 8-12-leaf stage the greatest reduction in MDA was caused by 150 $mg\ L^{-1}$ sodium silicate and 50 $mg\ L^{-1}$ calcium silicate (Fig. 5A, B). Cell membrane stability is decreased by lipid peroxidation. Silicon by decreasing malondialdehyde causes increasing stability of cell membranes. Silicone increases cell membrane stability through




Fig. 4. Effect of sodium and calcium silicate at 8- 12 (A) and 4- 5-leaf stage (B) on proline content. Means with the same letter are not significantly different at 5% level of Duncan's test.




Fig. 5. Effect of sodium and calcium silicate at 8- 12 (A) and 4- 5-leaf stage (B) on malondialdehyde content. Means with the same letter are not significantly different at 5% level of Duncan's test.

changing the ratios of unsaturated and saturated fatty acids which maintained function and integrity of cytoplasmic membrane. Also, silicon increased SOD and APX activity that resulted in reduction in membrane degradation, increased membrane stability and reduced EL (Electrolyte Leakage) and thus increased plant tolerance to stress. Silicon, probably by increasing the potassium ions, protected the cell membrane against damage caused by oxidative stress. The effect of free radical attack on lipids and aldehydes was due to aldehydes including MDA. The increase in this biomarker could be due to the decreased activity of catalase and superoxide dismutase. Silicon decreased malondialdehyde levels compared to control in gerbera cut flowers (Kazemi *et al.*, 2012). Silicon and acetyl salicylic applied in carnation caused a decrease in malondialdehyde levels in non-stress conditions (Kazemi *et al.*, 2012). Analysis of the cytoplasmic membrane showed that silicon decreased EC (Electrical conductivity) and malondialdehyde in *Polygonatum multiflorum* (Rubinowska *et al.*, 2014). Liang *et al.* (2003) in *Hordeum vulgare* and Kim *et al.* (2014) in *Oryza sativa* showed that application of silicon decreased MDA amount. Recent results indicate that silicon increased activity of SOD, POD and CAT but decreased MDA in barley, tomato and maize (Al-Aghabary *et al.*, 2004; Mussa 2006; Liang *et al.*, 2007). In silicon-treated plants in non-stressed conditions, the activity of SOD, CAT was increased and APX activity slightly decreased (Tale Ahmad and Haddad, 2011).

H₂O₂

At 8-12-leaf stage sodium silicate reduced hydrogen peroxide effectively but at the 4-5-leaf stage calcium silicate was better (Fig. 6A, B). In both stages, 150 mg L⁻¹ silicate caused a reduction




Fig. 6. Effect of sodium and calcium silicate at 8- 12 (A) and 4- 5-leaf stage (B) on H₂O₂. Means with the same letter is not significantly different at 5% level of Duncan's test.

in hydrogen peroxide to its lowest rate. Ascorbate peroxidase, by using two ascorbic acid molecules, converts H₂O₂ to 2 MDHA molecules and water. Silicon by stimulating antioxidant enzymes effects on hydrogen peroxide causes the conversion hydrogen peroxide into water molecules and therefore the decrease in hydrogen peroxide concentration. According to Gunes *et al.* (2008) silicon treatment of sunflower reduced H₂O₂ amount. Sodium silicate causes a decrease in H₂O₂ and MDA in soybean (Miao *et al.*, 2010). Zhu *et al.* (2005) reported that Si decreased malondialdehyde and H₂O₂ in wheat. Mussa (2006) showed that silicon increased CAT activity and decreased H₂O₂ in corn.

Literature Cited

- Aebi, H. 1984. Catalase *in vitro*. Methods in Enzymology, 105: 121-126.
- Ahmad, I. and Hellebust, A. 1988. The relationship between inorganic nitrogen metabolism and proline accumulation in osmoregulatory responses of two euryhaline microalgae. Plant Physiology, 88: 348-354.
- Al-Aghbary, K., Zhu, Z. and Shi, Q. 2004. Influence of silicon supply on chlorophyll content, chlorophyll fluorescence and antioxidative enzyme activities in tomato plants under salt stress. Journal of Plant Nutrition, 27(12): 2101-2115.
- Bharwana, S.A., Ali, S., Farooq, M.A., Iqbal, N., Abbas, F. and Ahmad, M.S.A. 2013. Alleviation of lead toxicity by silicon is related to elevated photosynthesis, antioxidant enzymes suppressed lead uptake and oxidative stress in cotton. Journal of Bioremediation and Biodegradation, 4: 187-198.
- Brigelius-Flohe, R. and Flohe, L. 2003. Is there a role of glutathione peroxidases in signaling and differentiation? Biofactors, 17: 93-102.
- Epstein, E. 1999. Silicon. Annual review of plant physiology and plant molecular biology. 50: 641-664.
- Gang, L.U., Jian, W., Zhang, J., Zhou, Y. and Cao, J. 2008. Suppressive effects of silicon nutrient on phomopsis stem blight development in asparagus. HortScience, 43(3): 811-817.
- Ghamsari, L., Keyhani, E. and Golkhoo, S. 2007. Kinetics properties of guaiacol peroxidase activity in *Crocus sativus* L. corm during rooting. Iranian Biomedical Journal, 1: 137-146.
- Gharineh, M.H. and Karmollachaab, A. 2013. Effect of silicon on physiological characteristics wheat growth under water-deficit stress induced by PEG. International Journal of Agronomy and Plant Production, 4 (7): 1543-1548.
- Gunes, A., Pilbeam, D.J., Inala, A. and Cobana, S. 2008. Influence of silicon on sunflower cultivars under drought stress, I: growth, antioxidant mechanisms, and lipid peroxidation. Communications in Soil Science and Plant Analysis, 39: 1885-1903.
- Hsieh, T.H., Lee, J.T., Charng, Y.Y. and Chan, M.T. 2002. Tomato plants ectopically expressing *Arabidopsis* cbf1 show enhanced resistance to water deficit stress. Plant Physiology, 130: 618-626.
- Kang, H.M. and Saltveit, M.E. 2002. Chilling tolerance of maize, cucumber and rice seedling (leaves and roots) and differentially affected by salicylic acid. Physiologia Plantarum, 115: 577-579.
- Karmollachaab, A., Bakhshandeh, A., Gharineh, M.H., Moradi Telavat, M.R. and Fatahi, G. 2013. Effect of silicon application on physiological characteristics and grain yield of wheat under drought stress condition. International Journal of Agronomy and Plant Production, 4 (1): 30-37.
- Kaya, C., Tuna, L. and Higgs, D. 2006. Effect of silicon on plant growth and mineral nutrition of maize grown under water-stress conditions. Journal of Plant Nutrition, 29: 1469-1480.
- Kazemi, M., Gholami, M., Asadi, M. and Aghdasi, S. 2012. Efficiency of silicon, nickel and acetylsalicylic acid reduced senescence and extended vase life of cut rose flowers. Trends in Applied Sciences Research, 7: 590-595.
- Khelifa, S., M'hamdi, M., Rejeb, H., Belbahri, L. and Souayah, N. 2011. Relation between catalase activity, salt stress and urban environments in *Citrus aurantium* L. Journal of Horticulture and Forestry, 3(6): 186-189.
- Kim, Y., Khan, A.L., Kim, D., Lee, S., Kim, K., Waqas, M., Jung, H., Shin, J., Kim, J. and Lee, I. 2014. Silicon mitigates heavy metal stress by regulating P-type heavy metal ATPases, *Oryza sativa* low silicon genes and endogenous phytohormones. Plant Biology, 14: 1-13.
- Liang, Y., Chen, Q., Liu, Q., Zhang, W. and Ding, R. 2003. Exogenous silicon (Si) increases antioxidant enzyme activity and reduces lipid peroxidation in roots of salt-stressed barley (*Hordeum vulgare* L.). Journal of Plant Physiology, 160: 1157-1164.
- Liang, Y.C., Sun, W.C., Zhu, Y.G. and Christile, P. 2007. Mechanisms of silicon mediated alleviation of abiotic stresses in higher plants: a review. Environmental Pollution, 147: 422-428.

- Liang, Y.C., Zhenguo, S.Q. and Tongsheng, M. 1996. Effect of silicon on salinity tolerance of two barley genotypes. *Journal of Plant Nutrition*, 19: 173-183.
- Li-Ping, B., Fang-Gong, S., Ti-Da, G. and Zhao-Hui, S. 2006. Effect of soil drought stress on leaf water status, membrane permeability and enzymatic antioxidant system of maize. *Soil Science*, 16: 326-332.
- Miao, B., Han, X. and Zhang, W. 2010. The ameliorative effect of silicon on soybean seedlings grown in potassium-deficient medium. *Annals of Botany*, 105: 967-973.
- Mittler, R. 2002. Oxidative stress, antioxidants and stress tolerance. *Trends in Plant Science*, 7: 405-410.
- Moghadam, A., Mohamadi, A., Ramezani, A., Mansurifar, S., Sadat Asilan, K., Moradi-Gahderijani, M. and Sadat Jamian, S. 2013. Application of silicon ameliorates salinity stress in sunflower (*Helianthus annuus* L.) plants. *International Journal of Agriculture and Crop Sciences*, 6: 1367-1372.
- Mohaghegh, P., Khoshgoftarmanesh, A.H., Shirvani, M., Sharifnabi, B. and Nili, N. 2011. Effect of silicon nutrition on oxidative stress induced by *Phytophthora melonis* infection in cucumber. *The American Phytopathological Society*, 95(4): 455-460.
- Molassiotis, A., Sotiropoulos, T., Tanou, G., Diamantidis, G. and Therios, I. 2006. Boron-induced oxidative damage and antioxidant and nucleolytic responses in shoot tip cultures of apple rootstock (*Malus domestica* Borkh). *Environmental and Experimental Botany*, 56: 54-62.
- Moldes, C.A., Fontaode Lima, O., Camina, J.M., Kiriachek, G.S., Molas, M.L. and Tsai, S.M. 2013. Assessment of the effect of silicon on antioxidant enzymes in cotton plants by multivariate analysis. *Journal of Agriculture and Food Chemistry*, 61(47): 11243-11249.
- Muhammad, A.J. and Shaheed, A.I. 2012. Effects of salicylic acid and silicon on oxidative damage and antioxidant activity in mung bean cuttings under boron toxicity. *Journal of Babylon University/Pure and Applied Sciences*, 22: 335-348.
- Mussa, H.R. 2006. Influence of exogenous application of silicon on physiological response of salt-stressed maize (*Zea mays* L.). *Agriculture and Biology Journal*, 2: 293-297.
- Nakano, Y. and Asada, K. 1981. Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in spinach chloroplasts. *Plant Cell Physiology*, 22(5): 867-880.
- Paquin, R. and Lechasseur, P. 1979. Observation sur une méthode de dosage de la proline libérée dans les extraits de plantes. *Canadian Journal of Botany*, 57: 1851-1854.
- Popham, P.L. and Novacky, A. 1990. Use of dimethylsulfoxide to detect hydroxyl radical during bacteria-induced hypersensitive reaction. *Plant Physiology*, 96: 1157-1160.
- Rosa, S.B., Caverzan, A., Teixeira, F.K., Lazzarotto, F., Silveira, J.A.G., Ferreira-Silva, S.L., Shigeoka, S., Ishikawa, T., Tamoi, M., Miyagawa, Y., Takeda, T., Yabuta, Y. and Yoshimura, K. 2002. Regulation and function of ascorbate peroxidase isoenzymes. *Journal of Experimental Botany*, 35: 1305-1319.
- Rubinowska, K., Pogroszewska, E., Laskowska, H., Szot, P., Zdybel, A., Stasiak, D. and Kozak, D. 2014. The subsequent effect of silicon on physiological and biochemical parameters of *Polygonatum multiflorum* (L.) All. 'Variegatum' cut shoots. *Acta Scientiarum Polonorum Hortorum Cultus*, 13(1): 167-178.
- Tale Ahmad, S. and Haddad, R. 2011. Study of silicon effects on antioxidant enzyme activities and osmotic adjustment of wheat under drought stress. *Czech Journal of Genetic and Plant Breeding*, 47(1): 17-27.
- Tuna, A.L., Kaya, C., Higgs, D., Murillo-Amador, B., Aydemir, S. and Girgin, A.R. 2008. Silicon improves salinity tolerance in wheat plants. *Environmental and Experimental Botany*, 62: 10-16.
- Upadhyaya, A., Sankhla, D., Davis, T.D., Sankhla, N. and Smith, B.N. 1985. Effect of paclobutrazol on the activities of some enzymes of activated oxygen metabolism and lipid peroxidation in senescing soybean leaves. *Journal of Plant Physiology*, 121: 453-461.
- Velikova, V., Yordancv, I. and Edreva, A. 2000. Oxidative stress and some antioxidant systems in acid rain-treated bean plants. Protective role of exogenous polyamines. *Plant Science*, 151: 59-66.

- Weber, H., Chetelat, A., Reymond, P. and Farmer, E.E. 2004. Selective and powerful stress gene expression in *Arabidopsis* in response to malondialdehyde. *Plant Journal*, 37: 877–889.
- Zhang, C.Q., Hong, B., Li, J.K. and Gao, J.P. 2005. A simple method to evaluate the drought tolerance of ground-cover chrysanthemum rooted cuttings. *Scientia Agriculture Sinica*, 38: 789-796.
- Zhu, X., Gong, H., Chen, G., Wang, S. and Zhang, C. 2005. Different solute levels in two spring wheat cultivars induced by progressive field water stress at different developmental stages. *Journal of Arid Environment*, 62: 1–14.

Archive of SID