

## Assessment of Biochemical and Histopathological Changes in the Liver of Chick Embryo Treated with a Commercial Deltamethrin Formulation

Research Article

N. Bhaskar<sup>1</sup> and L. Shahani<sup>1\*</sup><sup>1</sup> Department of Zoology, IIS University, Jaipur, Rajasthan, India

Received on: 11 Aug 2014

Revised on: 26 Nov 2014

Accepted on: 15 Jan 2015

Online Published on: Sep 2015

\*Correspondence E-mail: [lata.shahani@iisuniv.ac.in](mailto:lata.shahani@iisuniv.ac.in)

© 2010 Copyright by Islamic Azad University, Rasht Branch, Rasht, Iran

Online version is available on: [www.ijas.ir](http://www.ijas.ir)

### ABSTRACT

We studied the hepatotoxic effects of a deltamethrin containing formulation (Decis®) on developing embryo of *Gallus domesticus*. Fertilized eggs were immersed in three different concentrations of deltamethrin (12.5, 25 and 50 mg/L) for 60 min at 37 °C prior to incubation (ED 0) and on the 4<sup>th</sup> day of incubation (ED 4) and incubated till embryonic day (ED) 16. Exposure to 50 mg/L of deltamethrin resulted in a significant decrease in total protein and total glycogen content in the liver when exposed on ED 4 and ED 0 respectively, while total glutathione content was reduced at concentrations applied on both the exposure days. Among enzymes, liver alkaline phosphatase activity was significantly increased at 25 and 50 mg/L deltamethrin when embryos were exposed on ED 0 and 4, whereas the glutamate pyruvate transaminase activity showed a marked increase in animals exposed to 50 mg/L deltamethrin only at ED 4. Histopathological evaluation revealed mild to marked cellular lesions in liver sections of insecticide treated embryo. The magnitude of most of these changes appeared to be dose dependent being more severe on exposure to highest dose of deltamethrin. The observed pathological changes include degeneration and necrosis of hepatocytes, cytoplasmic vacuolization, enlarged and congested sinusoidal spaces, leucocyte infiltrations and congestion and dilation of the central vein.

**KEY WORDS** biochemical, chick embryo, deltamethrin, hepatotoxic, pathological, synthetic pyre-throid.

### INTRODUCTION

Synthetic pyrethroids are rapidly replacing conventional pesticides like organochlorines, organophosphates and carbamates because of the worldwide concern about their safety (Datta and Kaviraj, 2003). These pyrethroids are synthetic analogues of pyrethrins extracted from *Chrysanthemum cinerariaefolium* flowers, having insecticidal properties; they came on the market in the early 1970 s for agricultural purposes. Deltamethrin, [(S)-a-cyano-3-phenoxybenzyl-(1R)-cis-3-(2, 2-dibromovinyl)-2, 2-dimethylcyclopropane carbo-xylate] type II synthetic pyrethroid, was synthesized in 1974 and since then, it has been

applied for a range of commercial crops and by extension controls a variety of pests. It is used as an active ingredient in the number of commercial available insecticide formulations, which are mostly used in cotton belt and also in veterinary medicinal formulations to treat cattle. Deltamethrin is considered the most powerful and therefore the most toxic pyrethroid. Even though this chemical is broken down via UV and sun light, it is quite resistant to storage and can preserve its activity for 6 months at 40 °C. For this reason it may represent a risk to both mammals and the ecosystem as whole (Özkan and Üstüner, 2012). Deltamethrin has varying degree of toxicological impacts in different animal species, such as fish (Datta and Kaviraj, 2003; Ural and Sa-

glam, 2005; Köprücü *et al.* 2006; Velišek *et al.* 2006; Sharma and Ansari, 2011; Amin and Hashem, 2012), Japanese quail (Martin, 1990), south American toad (Salibián, 1992) and rats (Manna *et al.* 2005; El-Maghraby *et al.* 2010).

The avian egg and developing embryo has been widely used in toxicological research due to the properties such as small size, well known embryonic development, ease of accessibility of animals during experimentation, short incubation period (21 days), minimal expenditure of time and money and possibility of experimenting on large scale for statistically valid results, which proved this animal as a promising experimental model for preliminary teratological screening of various toxicants (Jelinek, 1982; Kotwani, 1998). Therefore, the present study was undertaken to investigate the possible adverse effects of a commercial formulation of deltamethrin on the liver of the developing embryo of *Gallus domesticus*.

## MATERIALS AND METHODS

### Toxicant

The pesticide used in the present study was a concentrate formulation containing an emulsion of deltamethrin-(25 g deltamethrin/L; Decis®-Bayer Crop Science Limited, Gujarat, India).

### Test animal

Fertilized eggs of *Gallus domesticus* (BV 300 breed) were obtained from a commercial hatchery (Kewalramani Hatcheries, Ajmer, India), cleaned and placed in an incubator at 38±1 °C with relative humidity of 70-80%. The eggs were turned (90 degrees) periodically during storage. Use of the chick embryos was in conformity with the policies of Institutional Animal Care and Use Committee (IACUC, 2008). All the experimental procedures conducted on the animals, including the housing, care and maintenance of experimental animals / eggs, were approved by the Animal Ethical Committee of Institute.

### Experimental design

The eggs were immersed in different aqueous emulsions of insecticide prepared in distilled water (vehicle) for 60 min at 37 °C temperature, at two moments: prior to incubation (ED 0) and on embryonic day 4 (stage 24, Hamburger and Hamilton, 1951) (ED 4). The insecticides were used at the concentrations 12.5, 25 and 50 mg/L of deltamethrin, based on the recommended dose (25 mg/L) for crop protection. Two other groups represented the control group I (Untreated eggs) and II (eggs immersed in vehicle, i.e. distilled water). Twenty eggs were assigned for each treatment groups.

All the eggs were kept for incubation and candled daily for breakdown examination of clear and dead embryos. At the end of experiment, the liver was excised from all of the sacrificed chick embryos on embryonic day 16 (stage 42, Hamburger and Hamilton, 1951) and submitted to biochemical and histopathological analyses.

### Biochemical study

Dissected liver from each of sacrificed animal was washed with saline solution, weighed and cut into two parts. One liver fragment was homogenized (10%) with a known volume of ice-cold phosphate buffer saline (PBS, pH 7.4) for estimating the total protein content, total GSH content and the enzymatic activity. The other fragment of the liver was used for analysis of total cholesterol and total glycogen content. Total protein content was estimated according to the method of Lowry *et al.* (1951). The Liebermann and Burchardt reaction (Henry and Henry, 1974) was used for estimating total cholesterol content. Glycogen content was extracted and estimated according to Montgomery procedure (1957). The reduced form of glutathione (GSH) was determined according to Moron *et al.* (1979). The activities of phosphatases (Alkaline Phosphatase- ALP; orthophosphoric monoester phosphohydrolase, EC 3:1:3:1 and Acid Phosphatase -ACP; monoester phosphohydrolase, EC 3:1:3:2) were estimated according to the method of Kind and King (1954), while amino transaminases activities (Glutamate pyruvate transaminase-GPT; L-alanine 2-oxoglutarate aminotransferase; EC 2.6.1.2 and Glutamate oxaloacetate transaminase-GOT; L-aspartate, 2-oxoglutarate aminotransferase; EC 2.6.1.1) were estimated according to King's method (1965).

### Histopathological study

Samples of Bouin's fixed liver tissue were cut into smaller pieces after 24 hours of fixation. Tissues were processed in a series of graded ethanol and xylene and embedded in a mixture of paraffin and bee wax (3:1); liver sections were cut at 5 µm and stained with haematoxylin and eosin. The stained sections were examined and scoring of histopathological changes was done as follow: (-) absent; (+) mild; (++) moderate; (+++) severe and (++++) extremely severe (Bancroft *et al.* 1996).

### Statistical analysis

All the obtained values of biochemical parameters were presented as mean ±Standard error (SE) and statistical significance was performed by IBM SPSS Statistics 17 analytic software using student "t" test. The value of *p* less than (≤) 0.05, 0.01 and 0.001 were considered to be significant, highly significant and very highly significant against control values.

## RESULTS AND DISCUSSION

### Biochemical changes

The biochemical profiles of the total hepatic protein content was significantly decreased ( $P \leq 0.05$ ) in embryos treated with 50 mg/L of deltamethrin on the 4<sup>th</sup> day of incubation, whereas the total glycogen content showed a marked decrease ( $P \leq 0.05$ ) in the animals treated with 50 mg/L of deltamethrin prior to incubation. The total cholesterol content of liver resulted unaffected following deltamethrin treatments on either embryonic days. Liver GSH content was significantly ( $P \leq 0.05$ ), highly significantly ( $P \leq 0.01$ ) and very highly significantly ( $P \leq 0.001$ ) decreased after exposure to 12.5, 25 and 50 mg/L deltamethrin treatment respectively on embryonic day 0, while when the exposure occurred on the 4<sup>th</sup> day of incubation, the liver GSH content was significantly depleted at 12.5 mg/L deltamethrin concentration and highly significantly depleted at 25 and 50 mg/L of deltamethrin, compared with their respective control values (Figure 1). With regards to enzymes, a statistically significant ( $P \leq 0.05$ ) and a highly significant ( $P \leq 0.01$ ) increase in liver ALP activity was observed in animals recovered from eggs treated prior to incubation with 25 and 50 mg/L of deltamethrin respectively, while it was significantly increased ( $P \leq 0.05$ ) following treatment with 50 mg/L of deltamethrin on embryonic day 4. ACP activity remained unaltered. Among liver transaminases, GPT activity showed a highly significant increase ( $P \leq 0.01$ ) at 50 mg/L deltamethrin concentration only when eggs were exposed at the 4<sup>th</sup> day of incubation, while the activity of GOT remained unchanged (Figure 2).


### Histopathological changes

The appearance of liver sections of untreated and vehicle treated control chick embryo was normal with clearly defined liver parenchyma formed of hepatic cords and hepatic sinusoids, arranged radially around the central vein (Table 1 and Figure 3). The hepatocytes were polygonal, with centrally located round vesicular nuclei. Blood sinusoids were narrow, followed their branching and anastomosed with each other. However, some mild alterations under the standard control range were observed in few liver sections of both the controls and control II treated on ED 0 and ED 4, respectively. The liver sections of chick embryos treated with 12.5, 25 and 50 mg/L of deltamethrin on embryonic day 0 revealed congestion and dilation of the central vein, degeneration of hepatocytes that showed pyknotic nuclei, enlarged blood sinusoids and dense inflammatory leukocyte infiltration. Activations of Kupffer cells were also observed in a few animals treated with 25 mg/L of deltamethrin. Hyalinization and necrosis of hepatocytes were more severe in the group exposed to 50 mg/L of deltamethrin.

The characteristic radial arrangement of hepatocytes was lost in this group. Exposure to deltamethrin at 12.5 mg/L on embryonic day 4 revealed a moderate increase in sinusoidal spaces with some degenerative changes in hepatocytes and leucocyte infiltration. Clear round areas in hepatic parenchyma indicated fatty changes. Increased blood sinusoidal spaces with large vacuoles in the hepatic parenchyma, disturbed hepatic architecture due to necrosis of hepatocytes and dilation of central vein were more prominent in liver sections from the 25 and 50 mg/L of deltamethrin treated groups. In addition, clumping of hepatocytes in rounded masses in mid-zonal area was also observed in liver sections of 2 chick embryos treated with 25 mg/L of deltamethrin. Pesticides are generally known to interfere with important biochemical and enzymatic processes such as carbohydrates, lipids and protein metabolism that regulate the normal physiology of animals. Any type of alteration in these processes can alter the homeostasis of the organism and also affect normal functioning of their organs (Ksheerasagar *et al.* 2011). The toxic responses occur frequently in the liver as it is a predominant organ for xenobiotic metabolism and detoxification, as well as the first organ to encounter ingested toxicants that enter portal circulation (Levi, 1987; Treinen-Moslen, 2001). In our study, insecticide mediated biochemical alterations indicate hepatic damage, which was well supported by observing pathological lesions in the liver section of developing chick embryo.

### Biochemical changes

The results of the present study exhibited a decrease in total liver protein content in chick embryo exposed to 50 mg/L of deltamethrin on the 4<sup>th</sup> day of incubation. This depletion in protein content may be attributed to degradation and possible use of these proteins in metabolic processes during the stress condition induced by the insecticide. Indeed it is possible that animals may need more energy to detoxify the toxicant and to overcome the stress. Omotuyi *et al.* (2006) and Okechukwo and Auta (2007) reported that the reduction in total protein level is likely due to proteolysis of structural proteins by using their amino acid for translation of detoxifying enzyme synthesis and / or conjugation of insecticide metabolite. Impaired incorporation of the amino acids into polypeptide chains may also affect the quantity of protein (Okechukwo and Auta, 2007). Moid *et al.* (2012) observed a decline in the protein content in different organs such as the liver, spleen and gastrocnemius muscles of mice after deltamethrin treatment, suggesting that low protein level may indicate liver dysfunction resulting from a change in enzymatic activity which may either cause reduced protein synthesis or their increased breakdown.



**Figure 1** Deltamethrin effect on: (a) the total protein content; (b) the total cholesterol content; (c) the total glycogen content and (d) the total GSH content in the liver of 16 day old chick embryo recovered from eggs treated with deltamethrin (Decis) on embryonic day 0 and embryonic day 4

Each value represents Mean $\pm$ SE

N= 5-6 animals per group


Statistical difference from the controls: \* significant at ( $P\leq 0.05$ ); \*\* highly significant at ( $P\leq 0.01$ ) and \*\*\* very highly significant using student "t" test

ED: embryonic day and GSH: reduced glutathione

The present results are in agreement with those reported by other investigators who studied the effect of deltamethrin in various models, like *Cyprinus carpio* (Velišek *et al.* 2006), rats (Yousef *et al.* 2006; El-Maghraby *et al.* 2010), *Danio rerio* (Sharma and Ansari, 2011) and *Clarias gariepinus* (Amin and Hashem, 2012) and found that this insecticide causes a significant reduction in total protein content. Presently, a marked decrease in the total liver glycogen content was observed in embryos exposed to the highest dose concentration (50 mg/L) of deltamethrin prior to incubation. Our results are in accordance with the finding of Anwar *et al.* (2004) who also observed a decrease in liver glycogen content of 16-day old chick embryo treated with permethrin (50, 100 and 200 ppm) on day "0" of incubation. According to them, the significant reduction in total glycogen content might have occurred as a result of its use to detoxify the insecticide or its metabolites through the glucuronidation process, a process by which toxic metabo-

lites combine with glucuronic acid component of UDP - glucuronic acid and are excreted from liver through bile. Furthermore, they also attributed this decrease to an increase in activity of the phosphorylase enzyme (phosphorylase-a and phosphorylase-b) involved in glycogen breakdown (glycogenolysis) for the glucuronidation process or to provide glucose into the circulatory system to cope with the energy requirement of cells which are under stress. Present findings are comparable to findings of Muthukumaravel *et al.* (2013) and Bhusan *et al.* (2013) who studied the toxic effect of  $\lambda$ -cyfluthrin in *Oreochromis mossambicus* and that of cypermethrin and beta cyfluthrin in the Wistar rats respectively, reporting a significant depletion in total hepatic glycogen content.

Reduced glutathione (GSH) is a major endogenous antioxidant and also a redox regulator playing an important protective role in detoxifying both endogenous molecules (free radicals) and xenobiotics.



**Figure 2** Effect of deltamethrin treatments on: (a) ALP activity; (b) ACP activity; (c) GPT activity and (d) GOT activity in the liver of 16 day old chick embryo recovered from eggs treated with deltamethrin (Decis) on embryonic day 0 and embryonic day 4

Each value represents Mean $\pm$ SE

N= 5-6 animals per group

Statistical difference from the controls: \* significant at ( $P \leq 0.05$ ) and \*\* highly significant at ( $P \leq 0.01$ ) using student "t" test

ED: embryonic day; ALP: alkaline phosphatase; ACP: acid phosphatase; GPT: glutamate pyruvate transaminase; GOT: glutamate oxaloacetate transaminase; KAU: king armstrong unit: the amount of enzymes that transforms one mg of phenol in 15 minutes and unit of GPT and GOT activity:  $\mu$ mol of pyruvate formed/min/g tissue

High levels of toxicants that are detoxified by GSH can cause its severe depletion, which is an important factor in determining cellular redox status in cells (Han *et al.* 2006). The GSH depletion in liver (major reservoir of GSH) can potentiate covalent binding of xenobiotics and its metabolites to DNA, protein and other biomolecules and cause hepatotoxicity (Levi, 1987). The present results demonstrated that exposure to insecticide deltamethrin depleted hepatic GSH content in chick embryos in a dose dependent manner. This reduction in total reduced glutathione content was either due to its decreased synthesis (Parthasarathy and Joseph, 2011) or to its inclusion in the detoxification process of reactive metabolites caused by the insecticide (Aouacheri *et al.* 2005). In poultry birds, a decrease in liver GSH content has been observed following treatment with permethrin for a period of 30 days (Ezeji *et al.* 2012).

The present results are also in line with the work reported by Rehman *et al.* (2006) who noted a reduction of the GSH content in kidney and liver of mice treated with deltamethrin.

Alkaline phosphatases (ALP) are found mainly in parenchymal cells of the liver and are responsible for mediating the active transport of metabolites across the membrane. These enzymes are also associated with protein synthesis and glycogen metabolism (Pilo *et al.* 1972; Ksheerasagar *et al.* 2011).


In the current study, administration of deltamethrin on embryonic day 0 and 4 resulted in a significant elevation of liver alkaline phosphatase activity.

The increased level of ALP activity might be due to its increased synthesis in the presence of increased biliary pressure (Gabriel *et al.* 2009).

**Table 1** Semi-quantitative scoring of histopathological findings in the liver of 16 day old chick embryo recovered from eggs treated with deltamethrin on embryonic day 0 and 4

Histopathological findings	Embryonic day 0					Embryonic day 4				
	Control		Deltamethrin			Control		Deltamethrin		
	I	II	12.5 mg/L	25 mg/L	50 mg/L	I	II	12.5 mg/L	25 mg/L	50 mg/L
Degeneration and necrosis of hepatocytes	-	-	++	++	+++	-	+	++	+++	+++
Cytoplasmic vacuolation and enlarged blood sinusoids	-	-	+++	++	++	-	+	++	++	+++
Leucocyte infiltration	+	+	+++	+++	+++	-	-	+	++	+
Fatty infiltration	-	-	-	-	-	-	+	++	++	+++
Hepatocytes with pycnotic nuclei	-	+	+	++	++	-	-	-	-	-
Kuffer cell activation	-	-	-	+	-	-	-	-	-	-
Congestion and/or dilation of central vein	+	+	++	++	+++	-	-	+	++	++

(-): indicates normal; (+): indicates mild; (++) indicates moderate and (+++): indicates severe.



**Figure 3** Photomicrographs of liver sections from 16 day old chick embryo (Haematoxylin and Eosin, magnification of 400x). [a] liver section of untreated control chick embryo showing normal histological structure of radially arranged hepatocytes (H) around central vein (CV). [b] section of the liver from treated with 12.5 mg/L of deltamethrin on ED 0 showing hepatocytes with pycnotic nuclei (PN) and leucocyte infiltrations (LI). [c] section of the liver from treated with 25 mg/L of deltamethrin on ED 0 showing necrotic hepatocytes (N) with pycnotic nuclei (PN) leading to enlarged blood sinusoids (BS). [d] section of the liver from treated with 50 mg/L of deltamethrin on ED 0 showing severe hypertrophy of hepatocytes leading to necrosis (N) and increased sinusoidal space with congestion and vacuolization (V). [e] section of the liver from treated with 12.5 mg/L of deltamethrin on ED 4 showing degenerating hepatocytes (DH) around dilated central vein (CV). [f] section of the liver from treated with 25 mg/L of deltamethrin on ED 4 showing clumping of necrotic hepatocytes (N) and leucocyte infiltrations (LI) in central vein (CV). [g] section of the liver from treated with 50 mg/L of deltamethrin on ED 4 showing vacuolization (V) and necrotic hepatocytes with enlarged blood sinusoidal space (BS) and fatty infiltration (FI) leads to loss of radial arrangement of hepatic cells.

Similarly, [Shakoori et al. \(1996\)](#) and [Panati et al. \(2012\)](#) observed elevation in above parameters in the liver of freshwater fish *Ctenopharyngodon idella* and in CNS and muscle of crab *Ozotelphusa senex senex* exposed to synthetic pyrethroid fenvalerate. The data from the present study revealed that the liver GPT activity of chick embryo was elevated after exposure to deltamethrin on the 4<sup>th</sup> day of incubation. The elevation in GPT activity indicates tissue damage due to formation of reactive oxygen species (ROS) and reactive intermediate after insecticide treatment ([Ksheerasagar et al. 2011](#)). It also indicates enhanced mobilization of amino acids towards gluconeogenesis and effective transamination to provide keto acids to serve as precursors in both the glycolytic pathway and tricarboxylic acid (TCA) cycle to the synthesis of essential constituents and also to cope with energy demands during insecticide induced toxicity ([Tiwari and Singh, 2004](#); [Savithri et al. 2010](#)). An elevation of the above mentioned parameter is in concurrence with the reports of several investigators who found increased GPT activity in the liver of *Ctenopharyngodon idella* exposed with fenvalerate ([Shakoori et al. 1996](#)) and in the liver of rat exposed with flumethrin ([Mishra et al. 2012](#)) and cypermethrin and  $\beta$ -cyfluthrin ([Bhusan et al. 2013](#)) respectively.

### Histopathological changes

The histopathological alterations observed herein on liver of chick embryo revealed mild to marked cellular changes upon the administration of different doses of the insecticide deltamethrin. However, some of these changes were also observed in both the control groups, which were considered to be within the standard normal range and might have occurred due to experimental and / or laboratory conditions. The most prevalent and severe types of lesions encountered included the degeneration and necrosis of hepatocytes, cytoplasmic vacuolization, enlarged and congested sinusoidal spaces, leucocyte infiltrations and congestion and / or dilation of the central vein. [Sahu and Ghatak \(2002\)](#) also observed these hepatic cellular changes such as cytoplasmic vacuolization with disintegration of liver cells in the chick embryo treated with dimecron prior to incubation. They suggested that vacuolated cells might have resulted from strong toxic effect of insecticide on cell membrane, which gets ruptured, and on nuclear degeneration, causing cytoplasmic vacuolization. [Anwar et al. \(2004\)](#) and [Anwar and Shakoori \(2010\)](#) reported that increased sinusoidal spaces are required to increase the blood flow in the liver lobule so to increase oxygen and nutrient supply to fulfill the requirements of hepatocytes under stress. The degenerative and necrotic changes evidenced in the hepatocytes might have occurred as a result of insecticide-mediated liver injury, associated with the formation of reactive metabolites

through hepatic cytochrome P-450 catalyzed oxidation ([Jain et al. 2011](#)). These reactive metabolites bind to cellular macromolecules such as protein and unsaturated lipids present on plasma membrane and change their biological properties ultimately resulting in the membrane destruction ([Levi, 1987](#)). Furthermore, the leucocyte infiltrations observed were considered as prominent responses of body tissues suggestive of irritability, inflammation and hypersensitivity toward the insecticide mediated toxic insult ([Khogali et al. 2005](#); [Sakr, 2007](#)).

Moreover, the exposure of developing chick embryo to deltamethrin prior to incubation resulted in mild to moderate occurrence of darkly stained pycnotic nuclei in hepatocytes, which were also described by [Anwar et al. \(2004\)](#) in liver sections of chick embryo treated with permethrin, whom consider this type of nuclei indicative of irreversible nuclear condensation and DNA fragmentation, which are a hallmark of programmed cell death (apoptosis) resulting from insecticide toxicity. In the present study, fatty infiltrations were observed in the liver of chick embryo exposed to deltamethrin on 4<sup>th</sup> day of incubation. [Moon et al. \(2007\)](#) reported that fatty infiltrations in the liver could be due to accumulation of high levels of triglycerides in parenchymal cells, caused by the defect in any steps from synthesis to release of triglyceride conjugated apoproteins mediated by hepatotoxic substances. Furthermore, an excess of lipid content can also result from an oversupply of free fatty acid from adipose tissue due to its impaired oxidation ([Levi, 1987](#); [Treinen-Moslen, 2001](#)). The pathological changes observed in the present study are also similar to those reported by other researchers in different experimental animal models under the influence of deltamethrin ([Tos-Luty et al. 2001](#); [Manna et al. 2005](#); [Yildirim et al. 2006](#); [Staicu et al. 2007](#)).

### CONCLUSION

The results of the present investigation clearly allow to conclude that the observed biochemical and histopathological changes in the liver of developing chick embryo are due to the toxic effect of the deltamethrin containing formulation (Decis®) under the used experimental conditions. Although these results cannot be directly extrapolated to mammals, they may be useful for comprehensively predict mechanisms of potential teratogenicity caused by these insecticides at preliminary screening level. Further studies with technical grade deltamethrin on mammals would no doubt aid in better understanding of the potential health effects of this insecticide. But notwithstanding with this concept, it can be recommended that the deltamethrin containing formulations should be used with cautions in the environment where pregnant animal or woman live as even

low concentration of such insecticides may interfere with the normal foetal development.

## ACKNOWLEDGEMENT

This work was supported by the University Grant Commission, Bhopal, India under Grant No. 140/304071/09-10/CRO.

## REFERENCES

- Amin K.A. and Hashem K.S. (2012). Deltamethrin-induced oxidative stress and biochemical changes in tissues and blood of catfish (*Clarias gariepinus*): antioxidant defense and role of alpha-tocopherol. *BMC Vet. Res.* **8**, 45.
- Anwar K., Ali S.S. and Shakoori A.R. (2004). Effect of a single sublethal dose of permethrin on the development of liver in chick embryo. *Pakistan J. Zool.* **36**, 59-68.
- Anwar K. and Shakoori A.R. (2010). Cypermethrin toxicity in the liver of developing chick embryo. *Pakistan J. Zool.* **42**, 725-733.
- Aouacheri W., Saka S. and Djafer R. (2005). Effect of alphamethrin on glutathione redox system in rats. *J. King. Saud. Univ.* **18**, 51-56.
- Bancroft G.D., Stevens A. and Turner D.R. (1996). Theory and Practice of Technique. Churchill: Livingston. New York.
- Datta M. and Kaviraj A. (2003). Acute toxicity to the synthetic pyrethroid deltamethrin to freshwater catfish *Clarias gariepinus*. *Bull. Environ. Contam. Toxicol.* **70**, 296-299.
- El-Maghraby S.I., Taha H.A. and Hassan N.S. (2010). Effect of *Anthum graveolens* extract on biochemical and histopathological alteration of deltamethrin in rats. *J. Bioanal. Biomed.* **2**, 8-12.
- Ezeji E., Anyalogbu E.A., Ezejiofor T.N. and Udensi J.U. (2012). Determination of reduced glutathione and glutathione S-transferase of poultry birds exposed to permethrin insecticide. *Am. J. Biochem.* **2**, 21-24.
- Gabriel U.U., Obomanu F.G. and Oveh O.D. (2009). Enzymes in selected tissues of cat fish hybrid exposed to aqueous extracts from *Lepidagathis alopecuroides* leaves. *Electron. J. Environ. Agric. Food Chem.* **8**, 856-864.
- Hamburger V. and Hamilton H.L. (1951). A series of normal stages in the development of the chick embryo. *J. Morphol.* **88**, 49-92.
- Han D., Hanawa N., Saberi B. and Kaplowitz N. (2006). Mechanisms of liver injury. III. Role of glutathione redox status in liver injury. *Am. J. Physiol. Gastrointest. Liver. Physiol.* **291**, 1-7.
- Henry R.J. and Henry M. (1974). Clinical Chemistry, Principles and Techniques. Harper and Row, New York.
- Jain T., Koley K.M., Vadlamudi V.P., Ghosh R.C., Roy S., Tiwar S. and Sahu U. (2011). Diclofenac- induced biochemical and histopathological changes in white leghorn birds (*Gallus domesticus*). *Indian J. Pharmacol.* **41**, 237-241.
- Jelinek R. (1982). Use of chick embryo in screening for embryotoxicity. *Teratog. Carcinog. Mutagen.* **2**, 255-261.
- Khogali F.A., Sheikh J.M., Rahman S.A., Rahim A.A. and Daghestani M.H. (2005). Histopathological and hematological effects of dimethoate 40EC on some organs of albino mice. *J. King. Saud. Univ.* **18**, 73-87.
- Kind P.R.N. and King E.J. (1954). Estimation of plasma phosphatase by determination of hydrolyzed phenol with amino antipyrine. *J. Clin. Pathol.* **7**, 322-326.
- King J. (1965). Practical Clinical Enzymology. Van Nostrand, London, D. Company.
- Köprücü S.S., Köprücü K. and Ural M.S. (2006). Acute toxicity of the synthetic pyrethroid deltamethrin to fingerling European Catfish, *Silurus glanis*. *Bull. Environ. Contam. Toxicol.* **76**, 59-65.
- Kotwani A. (1998). Use of chick embryo in screening for teratogenicity. *Indian J. Physiol. Pharmacol.* **42**, 189-204.
- Ksheerasagar R.L., Hiremath M.B. and Kaliwal B.B. (2011). Impairment of hepatic biochemical contents and enzymes activities during carbosulfan intoxication in albino mice. *Int. Multidisciplin. Res. J.* **1**, 6-15.
- Levi P.E. (1987). Toxic Action. Pp. 133-184 in A Textbook of Modern Toxicology. E. Hodgson and P.E. Levi, Eds. Elsevier, New York.
- Lowry O.H., Rosebrough N.J., Farr A. and Randall R.J. (1951). Protein measurement with folin phenol reagent. *J. Biol. Chem.* **193**, 265-273.
- Manna S., Bhattacharyya D., Mandal T.K. and Das S. (2005). Repeated dose toxicity of deltamethrin in rats. *Indian J Pharmacol.* **37**, 160-164.
- Martin P.A. (1990). Effects of carbofuran, chlorpyrifos and deltamethrin on hatchability, deformity, chick size and incubation time of Japanese quail (*Coturnix japonica*) eggs. *Environ. Toxicol. Chem.* **9**, 529-534.
- Mishra A., Dewangan G., Mahajan V. and Mandal T.K. (2012). Effect of flumethrin on tissue biochemistry following oral administration in wistar albino rats. *Int. J. Pharmacol. Biol. Sci.* **3**, 191-200.
- Moid N., Patel F., Shrimali A., Desai K.R. and Highland H.N. (2012). Toxicological implications of type II pyrethroid, deltamethrin on certain tissues of Swiss albino mice. *Int. J. Pharmaceut. Sci. Health Care.* **5**, 41-54.
- Montgomery R. (1957). Determination of glycogen. *Arch. Biochem. Biophys.* **67**, 378-387.
- Moon Y., Kim H.K., Suh H. and Chung D.H. (2007). Toxic alterations in chick embryonic liver and spleen by acute exposure to Fusarium- producing mycotoxin deoxynivalenol. *Biol. Pharm. Bull.* **30**, 1808-1812.
- Moron M.J., Dipierre J.W. and Mannerv K.B. (1979). Levels of glutathione, glutathione reductase and glutathione-S-transferase activities in rat lungs and liver. *Biochem. Biophys. Acta.* **582**, 67-71.
- Muthukumaravel K., Sathick O. and Raveendran S. (2013). Lambda cyhalothrin induced biochemical and histological changes in the liver of *Oreochromis mossambicus*. *Int. J. Pure. Appl. Zool.* **1**, 80-85.
- Okechukwo E.O. and Auta J. (2007). The effects of sub-lethal doses of lamda-cyhalothrin on some biochemical characteristics of African catfish *Clarias gariepinus*. *J. Boil. Sci.* **7**, 1473-



- 1477.
- Omotuyi I.O., Oluyemi K.A., Omofoma C.O., Josiah S.J., Adesanya O.A. and Saalu L.C. (2006). Cyfluthrin-induced hepatotoxicity in rats. *African J. Biotechnol.* **5**, 1909-1912.
- Özkan O. and Üstüner O. (2012). Investigations about genotoxicity of deltamethrin. *Kafkas. Univ. Vet. Fak. Derg.* **18**, 69-74.
- Panati K., Venkata R.N. and Venkatramana R.A.T. (2012). Changes in ATPase and phosphatases in different tissues of crab, *Oziotelphusa senex senex*, following exposure to fenvalerate. *Bioscan.* **7**, 487-489.
- Parthasarathy R. and Joseph J. (2011). Studies on the hepatic antioxidant defense system in  $\lambda$ -cyhalothrin-induced oxidative stress in fresh water tilapia (*Oreochromis mossambicus*). *African J. Environ. Sci. Technol.* **5**, 530-534.
- Pilo B., Asanani M.V. and Shah R.V. (1972). Studies on the wound healing and repair in pigeon liver III. Histological studies on acid and alkaline phosphatases during process. *J. Aim. Morphol. Physiol.* **19**, 205-212.
- Rehman H., Ali M., Atif F., Kaur M., Bhatia K. and Raisuddin S. (2006). The modulatory effect of deltamethrin on antioxidants in mice. *Clinica. Chimica. Acta.* **369**, 61-65.
- Sahu C.R. and Ghatak S. (2002). Effect of dimecron on developing chick embryo: malformations and other histopathological changes. *Anat. Histol. Embryol.* **31**, 15-20.
- Sakr S.A. (2007). Ameliorative effect of Ginger (*Zingiber officinale*) on mancozeb fungicide induced liver injury in albino rats. *Australian J. Basic. Appl. Sci.* **1**, 650-656.
- Salibián A. (1992). Effects of deltamethrin on the South American toad, *Bufo arenarum*, tadpoles. *Bull. Environ. Contam. Toxicol.* **48**, 616-621.
- Savithri Y., Sekhar P.R., Rani K.U. and Doss P.J. (2010). Toxicity of chlorpyrifos on total proteins, free amino acids and aminotransferase activity levels in liver and kidney tissues of albino rats. *Adv. Pharmacol. Toxicol.* **11**, 13-19.
- Shakoori A.R., Mughal A.L. and Iqbal M.J. (1996). Effects of sublethal doses of fenvalerate (a synthetic pyrethroid) administered continuously for four weeks on the blood, liver, and muscles of a freshwater fish, *Ctenopharyngodon idella*. *Bull. Environ. Contam. Toxicol.* **57**, 487-494.
- Sharma D.K. and Ansari B.A. (2011). Effect of deltamethrin and a neem based pesticide achool on some biochemical parameters in tissues liver, ovary and muscle of Zebrafish *Danio rerio* (Cyprinidae). *Res. J. Chem. Sci.* **1**, 125-134.
- Staicu A.C., Munteanu M.C., Costin D., Costache M. and Dinischiotu A. (2007). Histological changes in deltamethrin-induced intoxication in *Carassius auratus gibelio* (Pisces-cyprinidae). *Biotechnol. Anim. Husband.* **23**, 619-626.
- Tiwari S., Tiwari R. and Singh A. (2012). Impact of cypermethrin on fingerlings of common edible carp (*Labeo rohita*). *Scientific World. J.* **1**-7.
- Tos-Luty S., Haratym-Maj A., Latuszynska J., Obuchowska-Przebirowska D. and Tokarska-Rodak M. (2001). Oral toxicity of deltamethrin and fenvalerate in Swiss mice. *Ann. Agric. Environ. Med.* **8**, 245-254.
- Treinen-Moslen M. (2001). Toxic responses of the liver. Pp. 471-489 in Casarett and Doull's Toxicology: The Basic Science of Poisons. K.D. Curtis, Ed. New York.
- Ural M.S. and Saglam N. (2005). A study on the acute toxicity of pyrethroid deltamethrin on the fry rainbow trout (*Oncorhynchus mykiss* Walbaum, 1792). *Pest. Biochem. Physiol.* **83**, 124-131.
- Velíšek J., Dobšíková R., Svobodová Z., Modrá H. and Lusková V. (2006). Effect of deltamethrin on the biochemical profile of common carp (*Cyprinus carpio*). *Bull. Environ. Contam. Toxicol.* **76**, 992-998.
- Yildirim M.Z., Benli A.C., Selvi M., Ozkul A., Erkoç F. and Koçak O. (2006). Acute toxicity, behavioral changes, and histopathological effects of deltamethrin on tissues (gills, liver, brain, spleen, kidney, muscle, skin) of Nile tilapia (*Oreochromis niloticus*) fingerlings. *Environ. Toxicol.* **21**, 614-620.
- Yousef M.I., Awad T.I. and Mohamed E.H. (2006). Deltamethrin-induced oxidative damage and biochemical alterations in rat and its attenuation by vitamin E. *Toxicology.* **227**, 240-247.