

Investigating critical growth stage of cotton subject to water deficit stress

Fiaz Ahmad*, Shabab Ud Din, Asia Perveen and Mohammad Naveed Afzal

Central Cotton Research Institute (CCRI), Multan, Pakistan

Abstract

Critical growth stage of cotton crop was investigated by artificially imposing water stress under field conditions. The crop was given water deficit stress for a period of 30 days at squaring (SS) and first boll split (BS) phases by maintaining the leaf water potential (ψ_w) at -2.2 ± 0.2 MPa whereas in control plots ψ_w was maintained at -1.6 ± 0.2 MPa by irrigation scheduling. The average irrigation water applied during two years was 2432 m³, 2174 m³, and 2194 m³ in NS, SS, and BS treatments, respectively. The results revealed that the imposed water stress, at either stage of crop, had adverse effects on cotton crop performance. The main stem height decreased by 12% and 7.4%, inter-nodal length by 9.2% and 4.3% in SS and BS, respectively over NS treatment. The decrease in dry biomass production was 4% and 7% in leaf; 21% and 11% in stalk and 20% and 13% in fruit in SS and BS treatments, respectively. Gas exchange characteristics were also adversely affected by imposed water stress showing a decrease of 18% and 28% in stomatal conductance, 16% and 22% in transpiration rate and 24% and 30% in net photosynthetic rate in SS and BS treatments, respectively. Chlorophyll (SPAD values), electrolyte leakage and cell injury values increased over non stressed crop. Lint percentage and fibre strength increased whereas fibre length decreased significantly in BS treatment. It was concluded that the squaring phase is more critical to water deficit stress in cotton.

Keywords: cotton; stress; gas exchange; yield; lint quality

Ahmad, F., S. Din, A. Perveen and M. N. Afzal. 2013. 'Investigating critical growth stage of cotton subject to water deficit stress'. *Iranian Journal of Plant Physiology* 4 (1), 873-880.

Introduction

Undoubtedly, cotton is recognized as being biologically drought tolerant. But, despite its xerophytic distinctiveness, it is widely grown as an irrigated crop. Water deficit is one of the most yield limiting factors as it affects growth and development (Umebese et al. 2009) by decreasing vegetative development, leaf area, photosynthetic and transpiration rates due to stomatal closure (Cornic and Massacci, 1996; Mwanamwenge et al., 1999). Several lines of evidence indicate that а decrease in photosynthesis due to water deficits has been attributed to both stomatal and non-stomatal limitations (Shangguan et al., 1999). The effect of water deficit, however, varies with the variety, degree and duration of stress and the growth stage of the plant (Adejare and Umebese, 2007). In several plants, growth and yield are slightly affected at the vegetative stage but drastically reduced at the reproductive stage (Adejare and Umebese, 2007; Ma et al., 2006). Since cotton plant is indeterminate in nature, the

^{*}Corresponding author *E-mail address*: fiazdrccri@gmail.com Received: August, 2013 Accepted: September, 2013

compensation capacity of a variety would determine the critical growth stage to cope with the water deficit conditions. Plants adapt to water deficits by changes in morphology, altered patterns of development and cellular metabolism (Umebese et al., 2009).

With the changes in climate and increasing demand due to intensified agriculture, water has become a scarce commodity for crop production (Reddy et al., 1996; Umar, 2006). Hence there is a great need to utilize the water resources in an efficient and intelligent manner. Apart from selection of water, efficient varieties for cultivation and identification of critical growth stage of a cotton cultivar may help to sustain its production under water stress environment.

Materials and Methods

A field experiment on cotton (cv. CIM-499) was conducted during 2008 and 2009 with three water stress treatments i.e. no water stress (NS) at $-1.6\pm$ 0.2 MPa leaf water potential (ψ_w), stress at squaring (SS, 35 days after sowing) and stress at first boll split (BS, 85 days after sowing) phase. ψ_w was maintained at -2.2 ± 0.2 MPa in water stress treatments for a period of 30 days by irrigation scheduling. The crop was sown in third week of May during both the years at a plant configuration of 75 x 30 cm in RCBD layout. ψ_w was measured by water potential apparatus (Chas W. Cook Div., England) from fully expanded youngest leaf excised at 11.00 hours. Gas exchange characteristics were measured by Cl-

Table 1

340 hand held portable photosynthesis system (CID, USA). For dry biomass production plants from unit land area were harvested at maturity, partitioned into leaf, stalk, fruit portions and dried at 80 °C (Wells and Meredith Jr., 1984). Irrigation water applied was measured by "Cut Throat Flume". The irrigation water applied in 2008 and 2009 was 2632 m³ & 2573 m³, 2374 m³ & 2342 m³ and 2404 m³ & 2385m³ in NS, SS, and BS treatments. respectively. Annual precipitations received during the experiment years 2008 and 2009 were 64.5 mm and 79 mm, respectively.

Results

Impact on plant structure development

Imposition of water stress, at either stage of growth, caused a significant negative impact on plant structure development. Averaged across the years, main stem height ranged from 95 cm to 108 cm, nodes on main stem 29 to 30 and intermodal length from 3.16 cm to 3.48 cm in different treatments. With the imposition of water stress main stem height decreased, over NS, by 12% and 7.4% and inter-nodal length by 9.2% and 4.3% in SS and BS treatments, respectively. Although nodes on main stem decreased by 3.2% each in SS and BS treatment but the decrease was not statistically significant. Significant differences in plant structure development were observed between the years (Table 1).

Treatmonte	Plant structure development										
Treatments	Height (cm)			Node n	umbers	Inter-nodes (cm)					
NS SS	2008 101c [*] 87a	2009 115c 103a	Mean 108C [#] 95A	2008 30 ^{ns} 29 ^{ns}	2009 32 ^{ns} 31 ^{ns}	Mean 31 ^{ns} 30 ^{ns}	2008 3.37c 3.00a	2009 3.59c 3.32a	Mean 3.48C 3.16A		
BS	94b	106b	100B	29 ^{ns}	31 ^{ns}	30 ^{ns}	3.24b	3.42b	3.33B		
Mean	94A ^ş	108B		29.3A	31.3B		3.20A	3.44B			

Impact of water stress imposition on plant structure development at maturity

^{*}Values with different letters in each column are statistically significant at p<0.05

[#]Means with different letters in each column are statistically significant at p<0.05

^{\$}Means with different letters are statistically significant at p<0.05 among years

ns: non-significant; NS: no stress; SS: stress at squaring; BS: stress at boll split

Impact on dry biomass production

biomass production Dry decreased significantly (p<0.05) with the imposed water stress at squaring and first boll split phases of crop. Averaged across the years, the leaf biomass ranged from 112.5 to 121.5 gm⁻², stalk biomass from 150 to 189 gm⁻², fruit biomass from 462 to 575 gm^{-2} and the total biomass from 728.0 to 885.5 gm^{-2} in different treatments. Here again the minimum dry biomass was produced by cotton plant in SS treatment. A comparison of the decreased dry biomass production over NS treatment revealed that leaf biomass decreased by 4% and 7%, stalk biomass by 21% and 11%, fruit biomass by 20% and 13% and total biomass by 18% and 12% in SS and BS treatments, respectively (Table 2).

Impact on gas exchange characteristics

Gas exchange characteristics like stomatal conductance (g_s) , transpiration rate (E)

and net photosynthetic rate (P_N) were affected significantly with water stress. Averaged data of two year show that the g_s ranged from 129 to 184 m mol CO₂ m⁻²s⁻¹, *E* from 3.25 to 4.13 m mol H₂O m⁻²s⁻¹, P_N from 27.2 to 38.8 μ mol CO₂ m⁻²s⁻¹ and P_N/E from 8.4 to 9.4 m mol H₂O/ μ mol CO₂. With the imposition of water stress g_s decreased by 18% and 28%, *E* by 16% and 22%, P_N by 24% and 30% and water use efficiency (P_N/E) by 9.6% and 10.6% in SS and BS treatments, respectively (Table 3).

Impact on boll production and seed cotton yield

Production of bolls, boll weight and seed cotton yield varied considerably among the stress treatments. Averaged data of two years revealed that the number of bolls per plant ranged from 19.5 to 22.0, individual boll weight from 3.03 to 3.18 g and seed cotton yield from 1908 to 2277 kg ha⁻¹ in different treatments. The adverse effect of water stress was also reflected in decreased

Table 2

Impact of water stress imposition on dry biomass production at maturity

Tuestaseate		Dry biomass production (gm ⁻²)												
Treatments	Leaf			Stalk			Fruit			Total				
	2008	2009	Mean	2008	2009	Mean	2008	2009	Mean	2008	2009	Mean		
NS	118c [*]	125c	121.5C [#]	180c	198c	189C	545c	605c	575C	843c	928c	885.5C		
SS	113b	119b	116.0B	143a	157a	150A	400a	524a	462A	656a	800a	728.0A		
BS	110a	115a	112.5A	160b	176b	168B	448b	554b	501B	718b	845b	781.5B		
Mean	113.7A ^{\$}	119.7B		161A	177B		464A	561B		739A	857.7B			

 * Values with different letters in each column are statistically significant at p<0.05

[#]Means with different letters in each column are statistically significant at p<0.05

^sMeans with different letters are statistically significant at p<0.05 among years

Table 3

Impact of water stress imposition on stomatal conductance (gs), transpiration rate (E), net photosynthetic rate (P_N) and wateruse-efficiency (P_N/E)

_												
Treatments	gs (mmol CO ₂ m ⁻² s ⁻¹)			<i>E</i> (mmol H ₂ O m ⁻² s ⁻¹)			<i>P</i> _N (μmol CO ₂ m ⁻² s ⁻¹)			P _N /E (mmol H₂O/μmol CO₂)		
NS	2008 164c [*]	2009 203c	Mean 184C [#]	2008 4.06c	2009 4.20c	Mean 4.13C	2008 37.1b	2009 40.5b	Mean 38.8B	2008 9.1 ^{ns}	2009 9.7 ^{ns}	Mean 9.4 ^{ns}
SS	143b	156b	150B	3.37b	3.53b	3.45B	28.3a	30.4a	29.4A	8.4 ^{ns}	8.6 ^{ns}	8.5 ^{ns}
BS	125a	133a	129A	3.14a	3.34a	3.24A	25.9a	28.4a	27.2A	8.3 ^{ns}	8.5 ^{ns}	8.4 ^{ns}
Mean	144A ^{\$}	164B		3.52A	3.69B		30.4 ^{ns}	33.1 ^{ns}		8.6 ^{ns}	8.9 ^{ns}	

^{*}Values with different letters in each column are statistically significant at p<0.05

[#]Means with different letters in each column are statistically significant at p<0.05

 $^{
m S}$ Means with different letters are statistically significant at p<0.05 among years; ns means non-significant

production of bolls, their weight and total seed cotton yield. Over NS treatment, with imposed water stress bolls per plant decreased by 10% and 5%, boll weight by 4% and 5% and seed cotton yield by 16% and 10% in SS and BS treatments, respectively (Table 4).

Impact on chlorophyll, electrolyte leakage and cell injury

Accumulation of chlorophyll (SPAD values) was observed with water stress imposed at either stage of the crop. Averaged across the years, the chlorophyll content ranged from 45.0 to 47.9 in different treatment. The imposition of water stress at SS and BS phases caused an increase of 5% and 6%, respectively in chlorophyll content over NS. Chlorophyll content did not vary significantly between 2008 and 2009. Electrolyte leakage varied from 212 to 323 μ mhos cm⁻¹ and cell injury from 80 to 96% in different treatments. Average data of the two years revealed that the electrolyte leakage increased, over NS, by 47% in SS and 36% in BS. The cell injury varied from 80 to 96% in different treatment irrespective of the years. The imposition of water deficit stress increased cell injury by up to 14% & 10% in SS and BS, respectively. Here again the cell injury did not vary significantly among the years (Table 5).

Impact on lint quality

Lint percentage increased significantly in BS treatment in 2008 only and a non significant change was observed in 2009. The average data of two years also showed significant increase of lint (%) in BS over the other treatments. However, the fiber length decreased significantly (p<0.05) from 28.8 mm in NS to 28.4 mm in SS and 27.7 mm in BS treatments. Fiber strength varied from 92.9 to 94.5 thousand pound per square inch (tppsi) in different treatments, irrespective of the years. In BS treatment, fiber strength increased significantly when compared with the other treatments. Although micronaire increased with water stress imposed at both stages of the crop over non water stress, the differences were not significant. Among the years of study no significant variations in fiber quality characteristics were observed (Table 6).

Table 4

Inner	ot of	functor	ctrocc i	mnacitian	an hall	nraduction	and cood	cotton .	d ald
IIIIII	(1 0)	i waier	STRESST	mbosinon		Droduction	and seed	COLION	vieio
					0	p. 0 0 0 0 0 0 0 0		000000	,

T	Seed cotton yield and its components												
reatments	Bolls p	er plant		Boll weig	ht (g)		na⁻¹)						
	2008	2009	Mean	2008	2009	Mean	2008	2009	Mean				
NS	20b [*]	24b	22.0B [#]	3.23c	3.13c	3.18C	2151c	2403c	2277C				
SS	18a	21a	19.5A	3.15b	2.94b	3.05B	1730a	2086a	1908A				
BS	19ab	23b	21.0B	3.14a	2.92a	3.03A	1879b	2239b	2059B				
Mean	19A ^{\$}	22.7B		3.17B	3.00A		1920A	2243B					

Values with different letters in each column are statistically different at p<0.05

[#]Means with different letters in each column are statistically different at p<0.05

Table 5

Impact of water stress imposition on chlorophyll (SPAD values), electrolyte leakage and cell injury

Treatments	Chlorop	ohyll (SPA	D value)	Electro	lyte leakag	e (µS cm⁻¹)	Cell injury (%)			
	2008	2009	Mean	2008	2009	Mean	2008	2009	Mean	
NS	45.7a [*]	44.3a	45.0A [#]	216a	212a	214A	82a	80a	81.0A	
SS	47.8b	46.6b	47.2B	323c	303b	313C	96b	93b	94.5B	
BS	48.3b	47.4b	47.9B	294b	288b	291B	92b	90b	91.0B	
Mean	47.3 ^{ns}	46.1 ^{ns}		278 ^{ns}	268 ^{ns}		90.0 ^{ns}	87.7 ^{ns}		

^{*}Values with different letters in each column are statistically significant at p<0.05

[#]Means with different letters in each column are statistically significant at p<0.05

Discussion

Since water is the primary component of actively growing plants ranging from 70-90% of plant fresh mass (Gardner et al., 1984), limited water availability would have implications on the growth and physiological processes of all plants. Water deficit conditions have been reported to inhibit cell enlargement (Jaleel et al., 2009), thereby restricting the development of plant structure due to interruption of water flow from xylem to the surrounding elongating cells (Nonami, 1998). In the present studies the imposition of water deficit stress caused significant reduction in main stem height owing to the decreased intermodal length as nodes per plant did not change significantly among the treatments. The results are in conformity to the findings of Zhang et al. (2004) and Petropoulos et al. (2008) who also reported significant reduction in plant height in other crop plants due to water deficit stress. A comparison among the water stress treatments indicated that the maximum reduction in main stem height was observed when water stress was imposed at squaring phase. Variation in the plant response to water deficit among different varieties, degree and duration of stress and the growth stage of the plant has been observed (Adejare and Umebese, 2007; Jaleel et al., 2008). Since the squaring phase of cotton plant is the initial stage for plant architecture development, any stress at this stage of crop may limit the overall performance of short duration cotton varieties. The decreased dry biomass production was inevitable due to reduced plant growth and development as a result of water deficit stress. Reduced cell growth

by loss of turgor pressure (Anjum et al., 2011; Taiz and Zeiger, 2006) under water stress is considered to be the main causal effect for decreased biomass production (Farooq et al., 2009; Zhao et al., 2006). Cotton genotypes showing tolerance to water deficit conditions tend to divert from vegetative to reproductive organs. Therefore, the productivity of cotton under drought stress would be determined by the processes of dry matter partitioning and temporal biomass distribution (Kage et al., 2004).

Drought-induced reduction in leaf area is ascribed to suppression of leaf expansion through reduction in photosynthesis (Rucker et al., 1995). The imposed water stress adversely affected the physiological processes as g_s , E, and P_N decreased significantly. Similar results have been reported by Anjum et al. (2011a) who found that drought stress in maize led to considerable decline in net photosynthesis (33.22%), transpiration rate (37.84%), stomatal conductance (25.54%), water use efficiency (50.87%) and intercellular CO₂ (5.86%) as compared to well water control. Among the treatments, the stress imposed at first boll split phase affected the physiological phenomena (g_s, E) of cotton plant more than the stress at squaring phase, except the P_N which did not differ significantly among treatments. Gas exchange characteristics varied marginally among the years of study. Many studies have shown the decreased photosynthetic activity under drought due to stomatal or non-stomatal stress mechanisms (Del Blanco et al., 2000; Samarah et al., 2009). Stomatal closure, an abscisic acid mediated response (Borel and Simonneau, 2002) that results in decreased stomatal conductance and photosynthetic rate (Sunkar et al., 2003) and

Table 6

Impact of water stress imposition at various stages of growth on lint percentage, fiber length, fiber strength and micronaire values

Treatments	Lint (%)			L	Length (mm)			ength (tpp	si) [¥]	Micronaire (µg inch ⁻²)		
	2008	2009	Mean	2008	2009	Mean	2008	2009	Mean	2008	2009	Mean
NS	38.4a*	39.6 ^{ns}	39.0A [#]	29.2c	28.4b	28.8b	92.3a	93.4a	92.9A	4.48 ^{ns}	4.50 ^{ns}	4.49 ^{ns}
SS	38.5a	39.9 ^{ns}	39.2AB	28.7b	28.0b	28.4b	93.0a	94.2b	93.6A	4.50 ^{ns}	4.53 ^{ns}	4.52 ^{ns}
BS	39.0b	40.0 ^{ns}	39.5B	28.0a	27.4a	27.7a	94.4b	94.6b	94.5B	4.58 ^{ns}	4.59 ^{ns}	4.59 ^{ns}
Mean	38.6 ^{ns}	39.8 ^{ns}		28.6 ^{ns}	27.9 ^{ns}		93.2 ^{ns}	94.1 ^{ns}		4.52 ^{ns}	4.54 ^{ns}	

^{*}Values with different letters in each column are statistically significant at p<0.05

[#]Means with different letters in each column are statistically significant at p<0.05

ns: non-significant

less leaf transpiration due to water stress (Osmond et al., 1987) help the plants to survive under adverse environmental conditions.

The adverse effects of water deficit stress growth and development and on plant physiological phenomena were reflected in terms of reduced fruit production and seed cotton yield. Although seed cotton yield decreased under water deficit stress imposed at both stages of growth, the maximum decline was observed in SS treatment, owing to the less number of bolls per plant. The results are in conformity to the previous findings where decreased cotton (Gossypium hirsutum L.) lint yield was associated with more fruit abortion, due to increased levels of abscisic acid (Borel and Simonneau, 2002) and decreased boll production under moisture during the reproductive deficits growth (Pettigrew, 2004). Moreover, the hormonal imbalance in squares and bolls, under water stress, plays a significant role in fruit shedding (Guinn et al., 1990).

The imposed water stress caused increase in chlorophyll (SPAD value) over control plots, however the water stressed treatments (SS & BS) were not statistically different from each other. The accumulation of chlorophyll under water stress conditions has been found to be a common observation in plants (Hamada, 1996; Jabeen et al., 2008) in order to protect themselves from photo-damage, thereby reducing the extent of absorbed light by changes in chlorophyll content (Murchie and Horton, 1997). The increase in electrolyte leakage and cell injury values, reflect water stress associated increased temperature effect on cell membrane damage. Iswari and Palta (1989) also reported increased cell membrane damage, due to loss of water potential under water deficit conditions.

The percentage of lint and its quality characteristics, although, did not vary greatly with the imposed water stress at either stage of the crop; however, fiber length decreased while fiber strength increased in BS treatment. The lint percentage has been reported to be negatively affected with the increased soil moisture levels (Grimes, 1969), due to the development of healthy and mature seeds (Saranga et al., 1998) under adequate soil moisture conditions. The decrease in fiber length in water stressed plots could be attributed to the increased levels of abscisic acid that inversely affected the final fiber length (Dasani and Thaker et al., 2006). Pettigrew (2004) also reported up to 2% decrease in fiber length under moisture deficit stress. Although micronaire increased with water stress at both stages of the crop over non-water stress, the differences were not significant. Furthermore, among the years of study no significant variations in fiber quality characteristics were observed.

Conclusion

Water stress imposed for a period of 30 days at either stage of crop exerted negative impact on plant structure development, biomass accumulation, gas exchange characteristics and seed cotton production. Water stressed plants showed increased chlorophyll, cell injury and electrolyte leakage, lint percentage, and fiber strength whereas fiber length decreased with water stress. Squaring phase of cotton plant, i.e., 35 days after sowing (DAS) was found to be more critical for water deficit stress as compared to first boll split phase, i.e., 85 DAS.

References

- Adejare, F. B. and C. E. Umebese. 2007. 'Stomatal Resistance to Low Leaf Water Potential at Different Growth Stages Affects Plant Biomass in *Glycine max* L'. *Am. J. Agric. Biol. Sci.*, 2: 136-141.
- Anjum, S. A., X. Y. Xie, L. C. Wang, M. F. Saleem, C. Man and W. Lei. 2011. 'Morphological, physiological and biochemical responses of plants to drought stress (Review article)'. *African J. Agric. Res.*, 6(9): 2026-2032.
- Anjum, S. A., L. C. Wang, M. Farooq, M. Hussain,
 L. L. Xue and C. M. Zou. 2011a. 'Brassinolide application improves the drought tolerance in maize through modulation of enzymatic antioxidants and leaf gas exchange'. J. Agron. Crop Sci., doi: 10.1111/j.1439-037X.2010. 00459.x.
- **Borel, C.** and **T. Simonneau**. 2002. 'Is the ABA concentration in the sap collected by pressurizing leaves relevant for analyzing drought effects on stomata? Evidence from ABA-fed leaves of transgenic plants with

modified capacities to synthesize ABA'. J. Exp. Bot., 53(367): 287–296.

- **Cornic, C.** and **A. Massacci**. 1996. 'Leaf photosynthesis under drought stress'. *In: Photosynthesis and Environment*. Ed. N. R. Baker, Kluwer Acad. Publs: 347–366.
- **Cushman, J. C.** and **H. J. Bohnert**. 2000. 'Genomic approaches to plant stress tolerance'. *Plant Biol.*, 3(2): 117–124.
- Dasani, S. H. and V. S. Thaker. 2006. 'Role of abscisic acid in cotton fiber development'. *Russ. J. Plant Physiol.*, 53(1): 62-67.
- Del Blanco, I. A., S. Rajaram, W. E. Kronstad and M. P. Reynolds. 2000. 'Physiological performance of synthetic hexaploid wheat– derived populations'. *Crop Sci.*, 40(5): 1257-1263.
- Farooq, M., A. Wahid, N. Kobayashi, D. Fujita and S. M. A. Basra. 2009. 'Plant drought stress: effects, mechanisms and management'. Agron. Sustain. Dev., 29(1): 185–212.
- Flexas, J., J. Bota, J. Galmes, H. Medrano and M. Ribas-Carbo. 2006. 'Keeping a positive carbon balance under adverse conditions: responses of photosynthesis and respiration to water stress'. *Physiol. Plant.*, 127(3): 343-352.
- Gardner, W. R. and H. R. Gardner. 1983. 'Principles of water management under drought conditions'. *Agric. Water Manage.*, 7(1-3): 143-155.
- Grimes, D. W., W. L. Dickens and W. D.
 Anderson. 1969. 'Functions for cotton (*Gossypium hirsutum* L.) production from irrigation and nitrogen fertilization variables:
 II. Yield components and characteristics'. *Agronomy Journal*, 61(5): 773–776.
- Guinn, G., J. R. Dunlap and D. L. Brummett. 1990. 'Influence of water deficit on the abscisic Acid and Indole-3-acetic acid contents of cotton flower buds and flowers'. *Plant Physiol.*, 93(3): 1117-1120.
- Hamada, A. M. 1996. 'Effect of NaCl, water Stress or both on gas exchange and growth of wheat'. *Biol. Plant.*, 38(3): 405-415.
- Hussain, M., M. A. Malik, M. Farooq, M. Y. Ashraf and M. A. Cheema. 2008. 'Improving drought tolerance by exogenous application of glycinebetaine and salicylic acid in

sunflower'. J. Agron. Crop Sci., 194(3): 193-199.

- **Iswari, S.** and **J. P. Palta**. 1989. 'Plasma membrane Atpase activity following reversible and irreversible freezing injury'. *Plant Physiology*, 90(3): 1088–1095.
- Jabeen, F., M. Shahbaz and M. Ashraf. 2008. 'Discriminating some prospective cultivars of maize (*Zea mays* L.) for drought tolerance using gas exchange characteristics and proline contents as physiological markers'. *Pak. J. Bot.*, 40(6): 2329-2343.
- Jaleel, C. A., R. Gopi and R. Panneerselvam. 2008. 'Growth and photosynthetic pigments responses of two varieties of *Catharanthus Roseus* to triadimefon treatment'. *Comp. Rend. Biol.*, 331(4): 272–277.
- Jaleel, C. A., P. Manivannan, A. Wahid, M. Farooq, R. Somasundaram and R. Panneerselvam. 2009. 'Drought stress in plants: A review on morphological characteristics and pigments composition'. Int. J. Agric. Biol., 11(1): 100–105.
- Kage, H., M. Kochler and H. Stützel. 2004.'Root growth and dry matter partitioning of cauliflower under drought stress conditions: measurement and simulation'. *Eur. J. Agron.*, 20(4): 379–394.
- Kaya, M. D., G. Okçub, M. Ataka, Y. Çıkılıc and Ö
 Kolsarıcıa. 2006. 'Seed treatments to overcome salt and drought stress during germination in sunflower (*Helianthus annuus* L.)'. *Eur. J. Agron.,* 24(4): 291–295.
- Ma, Q. Q., W. Wang, Y. H. Li, D. Q. Li and Q. Zou. 2006. 'Alleviation of photoinhibition in drought-stressed wheat (*Triticum aestivum*) by foliar-applied glycinebetaine'. *J. Plant Physiol.*, 163(2): 165-175.
- **Murchie, E. H.** and **P. Horton**. 1997. 'Acclimation of photosynthesis to irradiance and spectral quality in british plant species: chlorophyll content, photosynthetic capacity and habitat preference'. *Plant Cell Environ.*, 20(4): 438-448.
- Mwanamwenge, J., S. P. Loss, K. H. M. Siddique and P. S. Cocks. 1999. 'Effect of water stress during floral initiation, flowering and podding on the growth and yield of faba bean (*Vicia faba* L.)'. *Europ. J. Agron.*, 11(1): 1–11.

- **Nonami, H.** 1998. 'Plant water relations and control of cell elongation at low water potentials'. *J. Plant Res.*, 111(3): 373–382.
- Osmond, C. B., M. P. Austin, J. A. Berry, W. D. Billings and W. E. Winner. 1987. 'Stress physiology and distribution of plants'. *Bioscience*, 37(1): 38-48.
- Petropoulos, S. A., D. Dimitra, M. G. Polissiou and H. C. Passam. 2008. 'The Effect of water deficit stress on the growth, yield and composition of essential oils of parsley'. *Sci. Hort.*, 115(4): 393–397.
- **Pettigrew, W. T.** 2004. 'Moisture deficit effect on cotton lint yield, yield components, and boll distribution'. *Agron. J.*, 96(2): 377-383.
- Reddy K. R., H .F. Hodges and J. M. McKinion. 1996. 'Food and agriculture in the 21st Century'. *World Resource Review*, 8(1): 80-97.
- Rucker K. S., C. K. Kvien, C. C. Holbrook and J. E. Hook. 1995. 'Identification of peanut genotypes with improved drought avoidance traits'. *Peanut Sci.*, 22(1): 14-18.
- Samarah, N. H., A. M. Alqudah, J. A. Amayreh and G. M. McAndrews. 2009. 'The Effect of late-terminal drought stress on yield components of four barley cultivars'. J. Agron. Crop Sci., 195(6): 427-441.
- Saranga, Y., I. Flash and D. Yakir. 1998.'Variation in water-use efficiency and its relation to carbon isotope ratio in cotton'. *Crop Science*, 38(3): 782–787.
- Shangguan, Z., M. Shao and J. Dyckmans. 1999. Interaction of osmotic adjustment and photosynthesis in winter wheat under soil drought'. J. Plant Physiol., 154(5-6): 753–758.

- Sunkar R., D. Bartels and H. H. Kirch. 2003. 'Overexpression of a stress-inducible aldehyde dehydrogenase gene from *Arabidopsis thaliana* in transgenic plants improves stress tolerance'. *Plant J.*, 35(4): 452-64.
- Taiz, L. and E. Zeiger. 2006. '*Plant Physiology'*. 4th edition, Sinauer Associates, Sunderland.
- **Umar, S.** 2006. 'Alleviating adverse effects of water stress on yield of sorghum, mustard and groundnut by potassium application'. *Pakistan Journal of Botany*, 38(5): 1373-1380.
- Umebese, C. E., T. O. Olatimilehin and T. A. Ogunsusi. 2009. 'Salicylic acid protects nitrate reductase activity, growth and proline in amaranth and tomato plants during water deficit'. *Am. J. Agri. Bio. Sci.*, 4(3): 224-229.
- Wells, R. and W. R. Meredith Jr. 1984. 'Comparative growth of obsolete and modern cotton cultivars: II. reproductive dry matter partitioning'. *Crop Science*, 24(5): 863-868.
- Zhang, M., L. Duan, Z. Zhai, J. Li, X. Tian, B. Wang, Z. He and Z. Li. 2004. 'Effects of plant growth regulators on water deficit-induced yield loss in soybean'. *Proceedings of the 4th International Crop Science Congress Brisbane, Australia.*
- Zhao, T. J., S. Sun, Y. Liu, J. M. Liu, Q. Liu, Y. B. Yan and H. M. Zhou. 2006. 'Regulating the drought-responsive element (DRE)-mediated signaling pathway by synergic functions of trans-active and transinactive DRE binding factors in *Brassica napus'*. J. Biol. Chem., 281 (16):10752-10759.