



Effect of salicylic acid on the quality of edible oil and fatty acids composition in different regions of sunflower (*Helianthus annuus* L.) heads

Hajar Khani Basiri, Mohammad Sedghi*, Raouf Seyed Sharifi

Department of Agronomy and Plant Breeding, Faculty of agriculture and natural resources, University of Mohaghegh Ardabili, Ardabil, Iran

Abstract

To study the effect of salicylic acid (SA) on the amount and quality of oil in different regions of sunflower inflorescence, a factorial experiment was conducted based on randomized complete block design with 3 replications at the University of Mohaghegh Ardabili, Ardabil, Iran. Salicylic acid at concentrations of 0, 0.1, and 0.2 g L⁻¹ were sprayed on whole plant at 3 stages (twice before flowering and a third time after flowering) and kernels were collected from different regions of capitulum (inner, middle, and outer layers). Samples were taken fifteen days after flowering with 5-day intervals and the amount of oil and four fatty acids (palmitic, stearic, oleic, and linoleic acids) were measured. Results showed that SA increased content of oil and unsaturated fatty acids but decreased the amount of saturated fatty acids and improved the quality of the kernels obtained from the mother plant. Linoleic acid content increased more than oleic acid by spraying of SA and the concentration of 0.2 g L⁻¹ SA was more effective. Kernel position had significant effect on the oil content and fatty acids, so that from the periphery toward the center there was a reduction trend in the oil and saturated fatty acid content.

Keywords: fatty acid; oil; salicylic acid; sunflower

Khani Basiri, H., M. Sedghi and R. Seyed Sharifi. 2017. 'Effect of salicylic acid on the quality of edible oil and fatty acids composition in different regions of sunflower (*Helianthus annuus* L.) heads'. *Iranian Journal of Plant Physiology* 8 (1), 2285-2292.

Introduction

Sunflower is an important source of edible oil after soybean, peanut, and canola in the world (Anonymous 2008). The oil contains large amounts of unsaturated fatty acids, which reduce cholesterol and prevent heart disease (Rathore, 2005). The oil has a pleasant taste and contains vitamins A, D, E, and K and is used in making margarine (Hussain et al., 2000). Seed oil content

varies between 25-42 percent, but can reach to 65% at very good conditions. Its protein content has been reported in the range of 10 to 25%. The most amounts of lipids accumulate in the sunflower seed cotyledons which contain about 78% of their weight and 7.4% of embryo weight. Oil from the seeds of this plant is drying oil with iodine index about 145-200 (Seyed Sharifi, 2007) which contains four major fatty acids including palmitic acid (16:0), stearic acid (18:0), oleic acid (18:1), and linoleic acid (18:2). Sunflower oil is rich

*Corresponding author

E-mail address: m_sedghi@uma.ac.ir

Received: March, 2017

Accepted: October, 2017

in linoleic acid and is mainly used in the food industry (Baydar and Erbas, 2005). The optimum temperature for oil synthesis is 18-22 °C (Anderson et al., 1978; Harris et al., 1978).

Environmental conditions and seed position on the mother plant are directly effective on the quality of produced seeds in mustard, sunflower and soybean (Guleria et al., 2007). In sunflower, kernel weight is reduced from the outer layers toward to the centre of capitulum (Goffner et al., 1988; Karadogan et al., 1998). The number of filled kernels in the centre is often low, while significantly reducing the sunflower yield. Low number of filled kernels in the centre (low yield at this area) is associated with vascular absorption (Durrieu, et al., 1985). Kernel filling in annual plants like sunflower is a fundamental process in reproduction and crop yield (Sedghi et al., 2008). Some plants such as sunflower in addition to protein and starch produce and accumulate large amounts of photo-assimilates during seed development in the form of triacylglycerol and store them as lipids which will convert to sucrose during germination through glyoxylate cycle (Taiz and Zeiger, 2006).

Accumulation of lipids during grain filling depends upon the transport of assimilates from the source to sink. In the process of grain filling and dry matter, oils and fatty acids accumulation from the periphery toward the centre of the capitulum have complex patterns (Munshi et al., 2003). Likely, capability of sunflower seeds produced from different positions of inflorescence in germination and emergence is due to the difference in the content of seed reserves (Munshi et al., 2007).

Salicylic acid (ortho-hydroxy benzoic acid, SA) is a natural phenolic compound and as a phytohormone plays an important role in the regulation of plant growth and development (Kang et al., 2003). SA stimulates the processes such as flowering, leaf stomatal closure, seed

germination, seedling growth, and activation of plant defense systems (Chandra et al., 2007). In addition to its effects on growth and resistance to stress and diseases, SA can increase fruit yield, glycolysis pathway, flowering in thermogenic plants, absorption and transport of ions, photosynthesis rate, stomatal conductance, and transpiration through the stomata (Hayat et al., 2010). The first report on the physiological effects of SA in plants was the *in vivo* induction of flowering and bud growth in cells of tobacco (Eberhard et al., 1989).

Salicylic acid significantly increased oil and protein content in sunflower and with increasing SA concentration from 75 to 100 mg L⁻¹ content of oil and protein significantly increased in comparison to control (Hayat et al, 2010). Also, saturated fatty acids decreased but unsaturated fatty acids increased (Dawood et al 2012). It was suggested that SA increases the concentration of auxin, gibberellin, zeatin, and zeatin ribosid while it reduces the abscisic acid contents in plants (Shehata et al., 2000, 2001; Zaghlool, 2002).

To the best of our knowledge there are few reports about the effect of SA on the lipid and fatty acid content in sunflower kernels from different positions, so this experiment designed for such purpose.

Materials and Methods

The experiment was conducted in the research field of the University of Mohaghegh Ardabili, Ardabil, Iran (15° 38' N and 15° 48' E). It is located in the semi-arid and cold climate with the mean precipitation of 400 mm annually. Soil texture was silty-loam with 1350 m altitude. Meteorological characteristics of the research field are shown in Table 1.

The experiment was conducted as factorial based on randomized complete block design with 3 replications. Treatments consisted

Table 1
Meteorological characteristics of the experimental field

| | Temperature (°C) | | | Humidity (%) | | | Precipitation (mm) |
|-----------|------------------|---------|---------|--------------|---------|---------|--------------------|
| | minimum | maximum | average | minimum | Maximum | average | |
| 22Apr-May | 6.6 | 21.4 | 14 | 37.9 | 92.2 | 65.0 | 18.9 |
| May-Jun | 10.3 | 25 | 17.7 | 39 | 90 | 65 | 25.6 |
| Jun-Jul | 12.7 | 23.9 | 18.3 | 57 | 95 | 76 | 8.7 |
| Jul-17Aug | 12.5 | 27.7 | 21 | 40 | 89 | 65 | 2.8 |

Table 2
Comparison of means for the oil content (%) of different treatments at sampling dates

| treatment | N ₁ * | N ₂ | N ₃ | N ₄ | N ₅ |
|-----------|---------------------|---------------------|---------------------|---------------------|---------------------|
| S1P1 | 0.0 ^f | 0.0 ^g | 0.0 ^h | 0.0 ^h | 0.600 ^g |
| S1P2 | 2.8667 ^c | 5.266 ^f | 9.700 ^f | 12.466 ^f | 16.233 ^f |
| S1P3 | 4.300 ^d | 11.100 ^c | 17.600 ^c | 22.233 ^c | 27.066 ^c |
| S2P1 | 0.0 ^f | 0.0 ^g | 0.766 ^g | 0.0 ^h | |
| S2P2 | 4.500 ^d | 7.333 ^e | 18.333 ^e | 13.166 ^e | |
| S2P3 | 7.600 ^b | 13.366 ^b | 19.766 ^b | 24.066 ^b | 29.633 ^b |
| S3P1 | 0.0 ^f | 0.0 ^g | 0.600 ^g | 0.800 ^g | 0.933 ^g |
| S3P2 | 5.800 ^c | 8.700 ^d | 12.333 ^d | 17.033 ^d | 21.266 |
| S3P3 | 11.367 ^a | 17.500 ^a | 22.666 ^a | 29.400 ^a | 33.433 ^a |

P1: inner, P2: middle, P3: outer position, S1: 0, S2: 0.1 and S3: 0.2 g L⁻¹ salicylic acid

* N is the samplings with 5 day intervals.

Table 3
Comparison of means for the oleic acid content (%) of different treatments at sampling dates

| treatment | N ₁ * | N ₂ | N ₃ | N ₄ | N ₅ |
|-----------|---------------------|---------------------|---------------------|---------------------|---------------------|
| S1P1 | 0.200 ^h | 0.533 ^h | 0.933 ^g | 1.166 ^g | 1.533 ^f |
| S1P2 | 4.500 ^f | 10.800 ^f | 16.033 ^f | 20.233 ^f | 24.533 ^e |
| S1P3 | 13.200 ^c | 21.100 ^c | 28.666 ^c | 34.833 ^c | 40.700 ^b |
| S2P1 | 0.400 ^{gh} | 0.800 ^{gh} | 1.333 ^g | 1.366 ^g | 1.633 ^f |
| S2P2 | 6.933 ^e | 13.666 ^e | 18.400 ^e | 21.333 ^e | 25.766 ^d |
| S2P3 | 17.133 ^b | 24.600 ^b | 31.300 ^b | 36.800 ^b | 41.166 ^b |
| S3P1 | 0.733 ^g | 0.966 ^g | 1.333 ^g | 1.433 ^g | 1.600 ^f |
| S3P2 | 9.900 ^d | 16.866 ^d | 20.733 ^d | 23.366 ^d | 27.433 ^c |
| S3P3 | 22.033 ^a | 27.333 ^a | 33.600 ^a | 38.133 ^a | 43.800 ^a |

P1: inner, P2: middle, P3: outer position, S1: 0, S2: 0.1 and S3: 0.2 g L⁻¹ salicylic acid

* N is the samplings with 5 day intervals.

of three levels of salicylic acid including 0, 0.1, and 0.2 g L⁻¹ and three separate positions for kernels in the head as inner, outer, and middle. Seeds of hybrid oilseed sunflower named Aline 122.1 × R-14 were planted in the field under favorable climatic conditions. Inter and intra row spacing was 65 and 30 cm, respectively. There were 5 rows in each plot. Average planting depth was 5 cm. In order to achieve the desired density four seeds were placed in each hole. Adjusted for density, thinning operation was carried out when the seedlings had two true leaves.

Salicylic acid was sprayed on leaves in all treatments except the control treatment during three stages (two times before flowering and another after flowering) at early morning or evening.

Samplings began after 15 days of flowering with 5 days intervals from the kernels of each head and the amount of the accumulation of lipid reserves was measured. To measure the oil

and fatty acids content, 500 mg of seed powder was dissolved in the solvents hexane, chloroform-methanol (2:1 v/v) and chloroform-methanol-water (1:2:0.8 v/v) and extraction was performed. The extracts were dried at low pressure and purified according to Belay and Daier (as quoted in De La Roche et al., 1973). Fatty acid composition was measured by gas-liquid chromatography. Data analysis was performed by SAS 9.1 software. Mean comparisons were done with LSD at 5% probability level.

Results

Oil content

The maximum oil content (33.43%) was observed by spraying 0.2 g L⁻¹ of SA at outside region kernels in all samples (Table 2). SA increased the amount of oil 6% in comparison to

Table 4

Comparison of means for the linoleic acid content (%) of different treatments at sampling dates

| treatment | N ₁ * | N ₂ | N ₃ | N ₄ | N ₅ |
|-----------|--------------------|--------------------|--------------------|--------------------|---------------------|
| S1P1 | 0.07 ⁱ | 0.10 ^j | 0.30 ⁱ | 0.60 ^h | 0.800 ^g |
| S1P2 | 3.80 ^f | 8.20 ^f | 11.50 ^f | 16.20 ^e | 18.633 ^e |
| S1P3 | 7.60 ^d | 12.70 ^d | 16.80 ^d | 20.30 ^c | 24.300 ^c |
| S2P1 | 0.10 ^h | 0.30 ^h | 0.50 ^h | 0.70 ^g | 1.033 ^g |
| S2P2 | 6.10 ^e | 10.70 ^e | 14.60 ^e | 17.50 ^d | 20.40 ^d |
| S2P3 | 10.20 ^b | 14.30 ^b | 17.90 ^b | 22.10 ^b | 26.933 ^b |
| S3P1 | 0.20 ^g | 0.40 ^g | 0.60 ^g | 0.80 ^f | 1.266 ^f |
| S3P2 | 9.10 ^c | 12.80 ^c | 17.40 ^c | 20.30 ^c | 24.033 ^c |
| S3P3 | 11.70 ^a | 16.30 ^a | 21.60 ^a | 28.10 ^a | 32.633 ^a |

P1: inner, P2: middle, P3: outer position, S1: 0, S2: 0.1 and S3: 0.2 g L⁻¹ salicylic acid

* N is the samplings with 5 day intervals.

Table 5

Comparison of means for the palmitic acid content (%) of different treatments at sampling dates

| treatment | N ₁ * | N ₂ | N ₃ | N ₄ | N ₅ |
|-----------|--------------------|--------------------|--------------------|--------------------|--------------------|
| S1P1 | 0.0 ^e | 0.0 ^g | 0.0 ^g | 0.043 ^g | 0.076 ^g |
| S1P2 | 1.500 ^b | 2.433 ^d | 3.266 ^d | 4.200 ^d | 5.066 ^d |
| S1P3 | 3.400 ^a | 5.900 ^a | 7.100 ^a | 8.033 ^a | 9.233 ^a |
| S2P1 | 0.0 ^e | 0.0 ^g | 0.0 ^g | 0.033 ^g | 0.046 ^g |
| S2P2 | 1.100 ^c | 1.633 ^e | 2.100 ^e | 2.933 ^e | 3.833 ^e |
| S2P3 | 1.500 ^b | 3.433 ^b | 4.766 ^b | 6.233 ^b | 7.466 ^b |
| S3P1 | 0.0 ^e | 0.0 ^g | 0.0 ^g | 0.0 ^g | 0.050 ^g |
| S3P2 | 0.0 ^e | 0.933 ^f | 1.600 ^f | 2.533 ^f | 2.600 ^f |
| S3P3 | 0.866 ^d | 2.933 ^c | 4.333 ^c | 5.333 ^c | 6.200 ^c |

P1: inner, P2: middle, P3: outer position, S1: 0, S2: 0.1 and S3: 0.2 g L⁻¹ salicylic acid

* N is the samplings with 5 day intervals.

Table 6

Comparison of means for the stearic acid content (%) of different treatments at sampling dates

| treatment | N ₁ * | N ₂ | N ₃ | N ₄ | N ₅ |
|-----------|--------------------|--------------------|--------------------|--------------------|--------------------|
| S1P1 | 0.0 ^f | 0.0 ^g | 0.0 ^g | 0.0 ^g | 0.050 ^g |
| S1P2 | 0.900 ^d | 1.766 ^d | 2.733 ^d | 3.266 ^d | 4.100 ^d |
| S1P3 | 2.500 ^a | 3.600 ^a | 5.100 ^a | 6.400 ^a | 7.533 ^a |
| S2P1 | 0.0 ^f | 0.0 ^g | 0.0 ^g | 0.0 ^g | 0.030 ^g |
| S2P2 | 0.733 ^e | 1.566 ^e | 2.100 ^e | 2.833 ^e | 3.533 ^e |
| S2P3 | 1.933 ^b | 3.00 ^b | 4.666 ^b | 5.766 ^b | 7.100 ^b |
| S3P1 | 0.0 ^f | 0.0 ^g | 0.0 ^g | 0.0 ^g | 0.026 ^g |
| S3P2 | 0.0 ^f | 0.600 ^f | 1.300 ^f | 1.866 ^f | 2.366 ^f |
| S3P3 | 1.233 ^c | 2.733 ^c | 3.700 ^c | 4.900 ^c | 5.833 ^c |

P1: inner, P2: middle, P3: outer position, S1: 0, S2: 0.1 and S3: 0.2 g L⁻¹ salicylic acid

* N is the samplings with 5 day intervals.

the control. None of SA concentrations had any effect on the oil content of central kernels (Table 2).

In the case of fatty acids, oleic and linoleic acids had a similar trend with oil content in different samplings (Tables 3 and 4); however, palmitic and stearic acids showed the opposite trend with oleic and linoleic acids (Tables 5 and 6). The highest percentages of palmitic (9.23) and

stearic acid (7.53) were observed at outer kernels without SA treatment and with increasing the SA concentration from 0 to 0.2 g L⁻¹ there was a reduction in the content of these fatty acids.

Salicylic acid increased the linoleic acid content more than oleic acid. Linoleic acid content increased approximately 8% (from 24.3 to 32.6%), but oleic acid content increased only 3% (from 40.7 to 43.8%) at the final sampling (Table 3 and

4). Therefore, SA was found to promote the edible quality of sunflower oil (by increasing linoleic acid content) and the frying quality (by increasing oleic acid content) of oil, while it was more effective on the edible quality.

Kernel position had significant impact on oil and fatty acids content so that from the periphery toward the center of the head, oil and saturated fatty acids decreased (Tables 2, 5 and 6).

Discussion

In general, SA increased oil percentage and oleic and linoleic acids content (unsaturated fatty acids) and decreased palmitic and stearic acids (saturated fatty acids). In this case with increasing unsaturated fatty acids, quality of oil also increased.

Noreen et al. (2009) reported that high concentrations of SA significantly increased seed oil content and essential unsaturated fatty acids, but it decreased stearic acid significantly. It seems that SA causes flowers to open early in all regions of the head and in this way kernels accumulate more dry matter which finally converts to more oil. Dawood et al. (2012) reported that SA induces flowering, increases flower life, delays aging, and increases cell metabolism.

Spraying of SA at the concentration of 0.2 gL⁻¹ increased kernel weight in all regions of head in sunflower, but was more effective on outer kernels, followed by middle position and then central kernels. SA is effective on dry matter accumulation in sunflower and it can be used to increase the yield of this plant (Sedghi et al., 2013).

Changes in the composition of seed reserves such as lipids, fatty acids, and other metabolites are biochemical changes in seed reservoirs (Chung et al., 1995). Generally, there are two major changes during seed maturation. First, changes in the seed volume and the second, biochemical and physiological changes which interfere in cell division and enlargement. SA affects many physiological and biochemical processes and thereby enhances growth and mobility (transport) of photosynthetic active substances from the source repository to the sinks (Abd El-Wahed et al., 2006; Al-Hakimi and Hamada, 2001). Assimilate direction to the seeds and flowers in different positions related to the

relationship between the sources (leaf and stem carbon stocks, Hall et al., 1990) and sinks and their phloem junctions (Farrar and Gunn, 1996). The increase in oil and protein content can be due to the increase in vegetative growth and the transport of materials in response to SA application (Dawood et al., 2012).

Weiss (1983) reported that during the development of sunflower kernels, the amount of oleic acid decreases while linoleic acid content increases. In addition, SA may be required for auxin and cytokinin synthesis (Metwally et al., 2003; Gharib, 2006). Ibrahim et al. (2001) also reported that cytokinins are likely to be necessary for synthesis of fatty acids, especially unsaturated and long-chain fatty acids compared with saturated fatty acids.

Competition for assimilates and space on the receptacle has been observed among the peripheral kernels and weak central kernels (Keiller and Morgan, 1988; Steer et al., 1988). The number of filled seeds in the center position of head is often low while it significantly reduces the total kernel quality. Low number of filled seeds in the center (i.e., low performance area) is associated with vascular absorption of this area (Durrieu et al., 1985).

Zimmerman and Fick (1973) observed that linoleic acid increases from periphery toward center and oleic acid shows an inverse trend. On the other hand, Baydar and Erbas (2005) in a two-year study reported that in the first year, palmitic, stearic, and linoleic acids decreased while oleic acid increased linearly from peripheral toward central regions, but in the second year there was an increase in palmitic, stearic, and oleic acids and a reduction in linoleic acid content.

Accumulation of fatty acids is strongly influenced by environmental conditions and follows a specific pattern. Certainly, one cannot conclude that with increase in one fatty acid content, others increase or decrease. With kernel growth and maturation all fatty acids are increased but the rate of increase is different for each acid. In other words, genetic differences in oil content and fatty acid composition at development stage needs to find some reasons (Ishikawa et al., 2001).

Kernel oil and fatty acids content in the peripheral and middle regions of head were much

higher than central kernels and the observed results may be due to SA effect (Tables 2 to 6). However, Baydar and Erbas (2005) and Karadogan et al. (1998) reported that the kernels of the middle region had the highest oil content (45.6%).

The average temperature during the growing season was 14-21 °C, relative humidity was 65-70%, and total rainfall was 119 mm (Table 1). In this study the effect of environmental factors should not be ignored. The maximum oil yield achieves in the altitudes less than 1,500 meters. The percentage of unsaturated fatty acids in oilseeds which are grown in high-latitude and temperate regions is higher than oilseeds which are grown in low-latitude and tropical places (Seyed Sharifi, 2007). Anderson et al. (1978) and Harris et al. (1978) reported that the best temperature for oil synthesis is 18-22 °C. This is consistent with our experimental conditions (Table 1).

Miralls et al. (1997) reported that high temperatures and low relative humidity in the growing season caused poor pollination and increased the number of hollow kernels. In general, favorable environmental conditions were presented in this study for growth and synthesis of unsaturated fatty acids and oils. So the results obtained from the use of SA should be considered besides the presence of favorable environmental conditions.

Conclusion

In this experiment, spraying of 0.2 g L⁻¹ SA increased the oil and fatty acid content of sunflowers which subsequently increased edible and frying quality of oil by increasing unsaturated fatty acid content. On the other hand, we observed that remaining kernels around the sampling area even in the central region had greater growth. From the literature and observations of this experiment it is concluded that the presence of more hollow kernels in the center of head may be due to the lack of enough space for growth and late opening of florets in this area. This leads central kernels be weak sinks for absorption of photo assimilates. So, it is suggested that breeders attempt to decrease the density of florets in central region of head and let the kernels fill efficiently.

References

- Abd El-Wahed, M.S.A. A.A. Amin and E.S.M. Rashad.** 2006. 'Physiological effect of some bioregulators on vegetative growth, yield and chemical constituents of yellow maize plants.', *World J Agric Sci*, 2(2): 149-155.
- Al-Hakimi, A.M.A. and A. M. Hamada.** 2001. 'Concentration of salinity stress on wheat plants by grain soaking in ascorbic acid, thiamin or sodium salicylate'. *Biologiae Plantarum*, 44(2): 253-261.
- Anderson, W.K., R.C.G. Smith and J. R. McWilliam.** 1978. 'A systems approach to the adaptation of sunflower to new environments. II. Effects of temperature and radiation on growth and yield. *Field Crops Res*, 1:153-163.
- Anonymous,** 2008. www.FAS.USDA.
- Osman, E.B.A. and M. M. M. Awed.** 2010. 'Response of sunflower (*Helianthus annuus* L.) to phosphorous and nitrogen fertilization under different spacing at New Valley. *Ass Univ Bull Environ Res*, 13(1): 11-19.
- Baydar, H. and S. Erbas.** 2005. 'Influence of seed development and seed position on oil, fatty acids and total tocopherol contents in sunflowers (*Helianthus annuus* L.)'. *Turk J Agric Forest*, 29: 179-186.
- Chandra, A., A. Anand and A. Dubey.** 2007. 'Effect of salicylic acid on morphological and biochemical attributes in cowpea'. *J Environ Biol*, 28(2): 193-196.
- Chung, C.H., Y. J. Yee, D. H. Kim, H. K. Kim and D. S. Chung.** 1995. 'Changes of lipid, protein, RNA and fatty acid composition in developing sesame (*Sesamum indicum* L.) seeds'. *Plant Sci*, 109: 237-243.
- De La Roche, I.A., C. J. Andrews and M. Kates.** 1973. 'Changes in phospholipid composition of a winter wheat cultivar during germination at 2 °C and 24 °C'. *Plant Physiol*, 51: 468-473.
- Durrieu, G., C. Percie du Dert and A. Merrien.** 1985. 'Anatomie du capitule tournesol'. Consequences sur la nutrition des akenes. In: Proc. Int. Sunflower Conf., 11th, Mar del Plata, Argentina.
- Eberhard, S., N. Doubrava, V. Narta, D. Mohnen, A. Southwick, A. Darviell and P.**

- Albershim.** 1989. Pectic cell wall fragments regulate tobacco thin-cell layer explants morphogenesis. *Plant cell*, 1: 747-755.
- Farrar, J.F. and S. Gunn.** 1996. Effects of temperature and atmospheric carbon dioxide on source-sink relations in the context of climate change. In: E. Zamski and A. Schaffer (eds). *Photoassimilate Distribution in Plants and Crops. Source-Sink Relationships*. Marcel Dekker Inc., New York, pp. 389-406.
- Gharib, F.A.** 2006. Effect of salicylic acid on the growth, metabolic activities and oil content of basil and marjoram. *Int J Agric Biol*, 8(4): 485-492.
- Goffner, D., R. Cazalis, C. Percie du Sert, J. Calmes and G. Cavalié.** 1988. 14C photoassimilate partitioning in developing sunflower seeds. *J Exp Bot*, 39: 411-422.
- Guleria, S., S. Sharma and S. K. Munshi.** 2007. Compositional changes in soybean (*Glycine max* L.) seeds influenced by their positions on stem axis. *J Food Sci Technol*, 44: 607- 610.
- Hall, A.J., D. M. Withfield and D. J. Connor.** 1990. Contribution of pre-anthesis assimilates to grain filling in irrigated and water stressed sunflower crops. I. Estimates from a carbon budget. *Field Crops Res*, 24: 273-294.
- Harris, H.C., J. R. McWilliam and W. K. Mason.** 1978. Influence of temperature on oil content and composition of sunflower seed. *Aust J Soil Sci*, 29(6): 1203-1212.
- Hayat, Q., S. Hayat, M. Irfan and A. Ahmad.** 2010. Effect of exogenous salicylic acid under changing environment. *Environ Exp Bot*, 68: 14-25.
- Hussain, M.K., E. Rasul and S. K. Ali.** 2000. Growth analysis of sunflower under drought conditions. *Int J Agric Biol*, 2: 136-140.
- Ibrahim, S.I., S. A. Zaghlool, I. Z. El-Shamy and S. S. Eisa.** 2001. The effect of kinetin or thiodiazuron in combination with ABA or JA on regulating growth, yield and seed lipids composition of sunflower (*Helianthus annuus* L.). *Ann Agric Sci*, 46: 465-483.
- Ishikawa, G., H. Hasegawa, Y. Takagi and T. Tanisaka.** 2001. The accumulation pattern in developing seeds and its relation to fatty acid variation in soybean. *Plant breed*, 120: 417-423.
- Kang, G., C. Wang, G. Sun and Z. Wang.** 2003. Salicylic acid changes activities of H₂O₂ metabolizing enzymes and increases the chilling tolerance of banana seedlings. *Environ Exp Bot*, 50(1): 9-15.
- Karadogan, T., K. Carkei and H. Ozer.** 1998. An investigation on differences in the quality of sunflower seeds in relation to their position in the head. *Helia*, 21: 121-129.
- Keiller, D.R. and D. G. Morgan.** 1988. Effect of pod removal and plant growth regulators on the growth, development and carbon assimilate distribution in oil seed rape. *J Agric Sci*, 111: 357-362.
- Metwally, A., I. Finkemeier, M. Georgi and K. J. Dietz.** 2003. Salicylic acid alleviates the cadmium toxicity in barley seedlings. *Plant Physiol*, 132: 272-281.
- Miralls, O.M., J. A. Valero and F. M. Olalla.** 1997. Growth, development and yield of five sunflower hybrids. *Eur J Agron*, 6: 47-59.
- Dawood, M.G., S. M. Sadak and M. Hozayen.** 2012. Physiological role of salicylic acid in improving performance, yield and some biochemical aspects of sunflower plant grown under newly reclaimed sandy soil. *Aust J Basic App Sci*, 6(4): 82-89.
- Munshi, S.K., S. Sandhu and S. Sharma.** 2007. Lipid composition in fast and slow germinating sunflower (*Helianthus annuus* L.) seeds. *Gen App Plant Physiol*, 33 (3-4): 235-246.
- Munshi, S.K., B. Kaushal and R. K. Bajaj.** 2003. Compositional changes in seed influenced by their position in different whorls of mature sunflower head. *J Food Sci Agric*, 83 (15): 1622-1626.
- Noreen, S., M. Ashraf, M. Hussain and A. Jamil.** 2009. Exogenous application of salicylic acid enhances antioxidative capacity in salt stressed sunflower (*Helianthus annuus* L.) plants. *Pak J Bot*, 41(1): 473-479.
- Rathore, P.S.** 2005. Techniques and managements of field crop production, First ed. Agrobios, India.
- Sedghi, M., H. Khani Basiri and R. Seyed Sharifi.** 2013. Effect of salicylic acid and kernel

- position on the sunflower grain filling'. *Adv Agric Sci Eng Res*, 3(3): 757 - 761.
- Sedghi, M., R. Seyed Sharifi, A. Namvar, T. Khandan-e-Bejandi and P. Molaie.** 2008. 'Responses of Sunflower yield and grain filling period to plant density and weed interference'. *Res J Biol Sci*, 3(9): 1048-1053.
- Seyed Sharifi, R.** 2007. 'Industrial crops'. 1st ed. Iran (In persion).
- Shehata, S.A.M., M. A. Saeed and M. S. Abou El-Nour.** 2000. 'Physiological response of cotton plant to the foliar spray with salicylic acid'. *Ann Agric Sci*, 45(1): 1-18.
- Shehata, S.A.M., S. I. Ibrahim and A. M. Zaghlool.** 2001. 'Physiological response of flag leaf and ears of maize plant induced by foliar application of kinetin (Kin) and salicylic acid (SA)'. *Ann Agric Sci*, 46(2): 435-449.
- Steer, B.T., P. J. Hocking and A. Low.** 1988. 'Dry matter, minerals and carbohydrates in the capitulum of sunflower (*Helianthus annuus* L.): Effects of competition between seeds and defoliation'. *Field Crops Res*, 18: 71-85.
- Taiz, L. and E. Zeiger.** 2006. 'Plant Physiology'. 4th ed. Massachusetts, USA.
- Weiss, E.A.** 1983. 'Oilseeds Crops'. Longman Group Ltd., New York.
- Zaghlool, S.A.M.** 2002. 'The effect of gibberellic acid (GA3), salicylic acid (SA), spermidine (Spd) and methods of application on growth, yield, some chemical constituents and some phytohormones in mungbean (*Vigna radiate* L.)'. *J Agric Sci*, 10(2): 493-504.
- Zimmerman, D.C. and G. N. Fick.** 1973. 'Fatty acid composition of sunflowers (*Helianthus annuus* L.) oil as influenced by seed position'. *J Amer Oil Chem Soc*, 50: 273-275.



اثر اسید سالیسیلیک بر کیفیت روغن خوراکی و ترکیب اسیدهای چرب دانه‌های حاصل از نواحی مختلف طبق آفتابگردان

هاجر خانی باصیری^۱، محمد صدقی*^۲ و رؤف سید شریفی^۲
۱ دانش آموخته رشته علوم و تکنولوژی بذر دانشگاه محقق اردبیلی
۲ استاد دانشگاه محقق اردبیلی

* عهده دارمکاتبات: m_sedghi2@uma.ac.ir

چکیده فارسی

به منظور بررسی اثر اسید سالیسیلیک (SA) بر مقدار و کیفیت روغن در نواحی مختلف طبق آفتابگردان، آزمایشی به صورت فاکتوریل در قالب طرح بلوک کامل تصادفی در دانشگاه محقق اردبیلی اجرا شد. اسید سالیسیلیک در غلظت‌های صفر، ۰/۱ و ۰/۲ گرم در لیتر روی بوته‌های آفتابگردان در ۳ مرحله (دو بار قبل از گلدهی و یک بار پس از گلدهی) محلولپاشی و دانه‌ها از نواحی مختلف طبق (منطقه مرکزی، میانی و بیرونی) برداشت شد. نمونه‌ها ۱۵ روز پس از گلدهی با فاصله زمانی ۵ روزه برداشت و مقدار روغن و چهار اسید چرب مختلف (اسید پالمیتیک، استئاریک، اولئیک و لینولئیک) اندازه‌گیری شد. نتایج نشان داد که SA موجب افزایش مقدار روغن و اسیدهای چرب غیراشباع شد، ولی مقدار اسیدهای چرب اشباع را کاهش داد که نتیجه آن بهبود کیفیت دانه‌های حاصل از گیاه مادری بود. مقدار اسید لینولئیک بیشتر از اسید اولئیک افزایش یافت و محلول‌پاشی SA با غلظت ۰/۲ گرم در لیتر موثرتر بود. موقعیت دانه‌ها اثر معنی‌داری بر مقدار روغن و اسیدهای چرب داشت، به طوری که از سمت بیرون به مرکز طبق مقدار روغن و اسیدهای چرب اشباع کاهش داشت.

کلمات کلیدی: اسید چرب، اسید سالیسیلیک، آفتابگردان، روغن