



Enhancement of valerenic acid production in *Valeriana officinalis* roots by methyl jasmonate -mediated transcriptional changes of sesquiterpene synthase genes

Hengameh Taheri*, Mansour Ghesmati

Department of Agronomy and Plant Breeding, Ramin Agriculture & Natural Resources University of Khouzestan, Iran

Abstract

Valeriana officinalis (valerian), as a nutraceutical herb, is widely used for its sedative and hypnotic properties. It is known that C15 sesquiterpenoid valerenic acid (VA) is the active ingredient responsible for pharmacological effects of *V. officinalis*. To evaluate the effect of methyl jasmonate (MeJA) concentrations (50 and 100 μM) in the modulation of expression patterns of the genes involved in valerenic acid (VA) biosynthesis, transcript abundance of identified sesquiterpene synthase (Sesqui-TPS) genes in the root of *V. officinalis* was monitored by quantitative real time PCR (qRT-PCR) within a 144 h time period. In addition, Valerenic acid contents were measured by high-performance liquid chromatography (HPLC). The highest amount of VA (12.45 mg/g dry weight (DW)) was found at 100 μM MeJA with a 12 fold increase over control culture (1.03 mg/g DW) at exposure time of 72 h. Moreover, MeJA in a concentration dependent manner enhanced transcription rate of VoTPS1 and VoTPS7 genes. Accordingly, exposure to 100 μM MeJA for 24 h can be more effective in the induction of these genes than that observed for 50 μM . Such enhancement was correlated with increased VA accumulation suggesting that these genes may be responsible for the biosynthesis of intermediates involved in the VA-biosynthetic pathway. However, MeJA treatment seemed to have a less significant effect on VoTPS3 expression than VoTPS1 and VoTPS7 genes. This results provide insights for more effective biosynthesis of VA by MeJA-mediated transcriptional changes of putative sesqui-TPS.

Keywords: *Valeriana officinalis*; sesquiterpenes synthase; expression pattern; quantitative real time PCR

Taheri, H. and M. Ghesmati. 2018. 'Enhancement of valerenic acid production in *Valeriana officinalis* roots by methyl jasmonate -mediated transcriptional changes of sesquiterpene synthase genes'. *Iranian Journal of Plant Physiology* 9 (1), 2611-2617.

Introduction

Valeriana officinalis (valerian) is a well-known medicinal plant that has many chemical

constituents in its essential oil extract including monoterpenoids, sesquiterpenoids, iridoid derivatives (known as valepotriates) and alkaloids (Letchamo et al., 2004; Wang et al., 2010). It is known that the C15 sesquiterpenoid valerenic acid (VA) is a pharmacologically-active ingredient

*Corresponding author
E-mail address: taheri@ramin.ac.ir
Received: December, 2017
Accepted: May, 2018

responsible for sedative and hypnotic properties of *Valeriana officinalis* (Bent et al., 2006; Takemoto et al., 2009). Valerian extract increases release of gamma aminobutyric acid (GABA) from brain nerve ending and then inhibits this inhibitory neurotransmitter from being reabsorbed back into nerve cells (Santos et al., 1994; Beaubru and Gray, 2000). Furthermore, VA decreases GABA degradation by its inhibitory effect on an enzyme that destroys GABA and thereby increases GABA availability in the synaptic cleft (Morazzoni and Bombardelli, 1995). Farnesyl diphosphate (FPP, C15) as a precursor for the biosynthesis of numerous structurally diverse sesquiterpenes (e.g. germacrene B/C/D and valerenadiene) is synthesized by the condensation of the 3 isomeric isoprene C5 units, namely, isopentenyl diphosphate (IPP) and dimethylallyl diphosphate (DMAPP) (Hampel et al. 2005). Yeo et al. (2013) revealed seven terpene synthase gene (VoTPSs) candidates. Among them, VoTPS1 and VoTPS7, which were expressed more highly in root tissue, utilized significant amount of FPP rather than geranyl diphosphate (GPP) as substrate. These pieces of information suggest that these terpene synthases may be sesquiterpene synthase (sesqui-TPS). In contrast, VoTPS3 mRNA was present in all tissue types examined (leaves, stems and roots) and used GPP rather than FPP as substrate. VoTPS1 was functionally characterized as a valerenadiene synthase, whereas VoTPS7 encoded germacrene synthase. It is thought that valerenic acid might be derived from valerenadien (Fig. 1). However, biochemical mechanism of their biosynthesis has been remained enigmatic. It was also proposed that germacrene intermediate might contribute to valernadien biosynthesis (Yeo et al., 2013).

Effective strategies such as metabolic engineering, tissue culture, and recruitment of phytohormones as the signaling molecules regulating metabolite pathways have been considered for enhancing the yield of pharmaceutical metabolites (Pandey et al., 2016). MeJA involved in enhancement of bioactive compounds production through the regulations of expression of a series of genes involved in the synthesis of secondary metabolites (Zhang et al., 2004).

In order to increase understanding of how the elicitor modulates VA metabolism in *V. officinalis*, MeJA-mediated dynamic responses of some identified VoTPS genes that probably involved in VA biosynthesis were measured by quantitative real time PCR (qRT-PCR).

Materials and Methods

Plant growth and phytohormone treatment

Seeds of *V. officinalis* were obtained from Pakanbazar Company, Esfahan, Iran. The seeds were surface sterilized with 70 % (v/v) ethanol for a min and rinsed with sterile distilled water. Seeds were then treated with 2.5 % sodium hypochlorite (NaClO) for 10 min, and then rinsed 3 times in distilled water. Surface-sterilized *V. officinalis* seeds were germinated on half-strength Murashige and Skoog (MS) medium (Murashige and Skoog, 1962), supplemented with 30 g l⁻¹ sucrose and solidified with 8 g/l agar. The pH of medium was adjusted to 5.8. The cultures media were maintained under controlled conditions in tissue culture room (16/8 h light/dark cycle at 25±2° C).

MeJA was purchased from Sigma Co. (Sigma Aldrich, Germany). A stock solution of MeJA was prepared by dissolving it in 96% ethanol. Two months after growth of seedlings on half-strength MS media, plants were subcultured on the media containing 50 and 100 μM MeJA. The control cultures were treated with only ethanol. The root sampling was collected at 12, 24, 48, 72, and 144 h after treatment to evaluate terpene synthase genes expression and VA content.




Fig. 1. Proposed pathway for the biosynthesis of the sesquiterpene valerenic acid (Yeo et al., 2013).

Table 1
Primer sequences designed for qRT-PCR

| Target gene | Accession number | sequence (5'-3') | Product size (bp) |
|-----------------|------------------|--|-------------------|
| <i>VoTPS1</i> | JX494699 | F: GTAGGCATGGGAGTAACTACAG R:TCGTGTCCAACCTTATCATCC | 119 |
| <i>VoTPS3</i> | JX494701 | F: CGTGCGTAAAGTCTTACG R: CTACTCACTCCGTGTTCTTTC | 114 |
| <i>VoTPS7</i> | JX494705 | F: GCTAGGCTCGACAATGATAAAG R: TGAACTCTACAACCACTTCTTC | 124 |
| <i>18S rRNA</i> | AJ236003 | F: AGGATGGCAAGCGTATTGG R: TGCATTTCCAGAGTGCTTC | 118 |

Determination of VA content

For preparation of VA extraction, 200 mg of dry powder of roots of *V. officinalis* were extracted three times with 5 ml of 70 % Me OH, sonicated for 10 min and then diluted to a final volume of 15 ml with Me OH. All samples were filtered through a 0.45 μ M micro-filter (Merck, Germany) subjecting to reverse-phase high-performance liquid chromatography (HPLC) analysis as described in previous studies (Tania and Brian, 2004).

HPLC analysis was carried out using a Knauer HPLC system (Germany). VA was separated by reverse-phase isocratic C18 column (250 \times 3 mm) with a pre-column (25 \times 4.6 mm, particle size 5 μ m) and detection by UV was conducted at 225 nm. The mobile-phase consisted of 0.5 % (v/v) phosphoric acid (A) and 0.5 % phosphoric acid:methanol mixture (27:73) (B) delivered at a flow rate of 1 ml min⁻¹ and an injection volume of 20 μ l. VA was provided by Sigma-Aldrich Co. (Germany). The standard solution was prepared by dissolving VA in methanol to obtain solutions with 5 different concentrations in the range of 5 to 25 mg L⁻¹. The calibration curves were constructed based on measuring their respective peak areas. This experiment was carried out with three replicates. Results were expressed as mean values \pm SD. The statistical significance of VA contents was analyzed using a *t* test with Excel software for significant differences between treated and control plants.

RNA extraction, cDNA synthesis, primers design, and qRT-PCR reaction

For expression analysis of terpene synthase genes, 100 mg of roots were collected from treated and control valerian and ground to a powder in liquid nitrogen. Total RNAs were extracted from the roots using GeneAll[®] RiboEx[™] kit (BioFrontier, Korea) based on manufacturer's protocol. RNA quantification was performed by using NanoDrop 2000C (Thermo Fisher Scientific Inc., MA, USA). First strand cDNA synthesis was done from 500 ng of total RNA using Fermentas kit (Revert Aid[™] First Strand cDNA Synthesis Kit) according to the manufacturer's instructions.

Primer pairs of the terpene synthase genes including *VoTPS1*, *VoTPS3* and *VoTPS7* were designed using online IDT Primer Quest[™] Software of Integrated DNA according to the predicted coding DNA sequence (CDS). The primers sequences designed for qRT-PCR amplification of target genes are listed in Table 1.

The quantitative real time PCR (qRT-PCR) was performed using HIFI SYBR[®] Green kit (Iran Master Mix and Step One Plus[®] (ABI, America) machine. PCR conditions were an initial denaturation of 10 min at 95 °C, followed by 40 cycles of denaturation at 95 °C for 15 s, annealing at 59 °C for 20 s and extension at 72 °C for 20 s. Relative fold changes in gene expression were calculated using the 2^{- $\Delta\Delta$ CT} comparative method (Livak and Schmittgen, 2001). The threshold cycle (Ct) for each gene was normalized against the Ct for *18S rRNA* gene (AJ236003) of *V. officinalis*, which was used as the internal reference. Relative




Fig. II. Time-course effects of 50 and 100 μM MeJA on VA production in the root of *V. officinalis* plants; values are mean \pm SD. Data were analyzed using *t* test on VA contents in the treated plants compared to the control ($P < 0.05$).

Expression Software Tool (REST)[®] software (Pfaffl (2001) was used to analyze the obtained data. This experiment was carried out with three biological and two technical repetitions.

Results

Effect of MeJA on valerenic acid production

To evaluate the effects of MeJA concentrations on stimulating VA production 24, 48, 72, and 144 h after elicitation, HPLC analysis of root cultures of valerian was performed. Efficiency of MeJA in promoting VA accumulation at both concentrations are shown in Fig. II. The maximum increments of 4.33 times (9.14 mg/g DW) in VA content in comparison to control culture (2.11 mg/g DW) were obtained after 144 h of exposure to 50 μM MeJA. However, maximum accumulation of VA (12.45 mg/g DW) was recorded at 100 μM MeJA with a 12 fold increase over control culture (1.03 mg/g DW) at 72 h. Accordingly, increasing MeJA concentration from 50 to 100 μM more effectively enhanced VA accumulation at shorter exposure time (72 h).

Effect of MeJA on expression pattern of Sesqui-TPS genes

To examine how dynamic transcriptional responses of some identified *VoTPS* genes

probably involved in VA biosynthesis to MeJA, transcript abundance of *VoTPS1*, *VoTPS3* and *VoTPS7* genes were monitored by qRT-PCR in the root of *V. officinalis* plants (Fig. III). At both MeJA concentrations, the highest level of *VoTPS1* and *VoTPS7* transcripts were detectable 24 h after elicitation. Nevertheless, transcript abundance of these genes increased with increasing MeJA concentration. Compared with the control, mRNA levels of *VoTPS1* and *VoTPS7* were 4 and 11.5-fold higher after exposure to 50 μM MeJA, respectively. However, after exposure to 100 μM MeJA, transcript expression of these genes reached the peak at 24 h (~25 and 68 times more compared to the control plants, respectively). Whereas, transcription rate of *VoTPS3* was mostly unaffected or down-regulated by MeJA at both concentrations.

Discussion

Our study showed that MeJA elicitation at both concentrations (50 and 100 μM) was substantially promoted VA content in the root of *V. officinalis*. However, exposure to 100 μM of MeJA more effectively enhanced VA accumulation than what observed for 50 μM . Moreover, a shorter period (72 h) was required for obtaining the highest increase in VA content than observed




Fig. III. Time-course transcriptional changes of *VoTPS1*, *VoTPS3* and *VoTPS7* induced by 50 and 100 μ M MeJA in the root of *V. officinalis* plants; for normalization, *18S rRNA* gene was used as the endogenous gene. The Y-axis represents relative quantity; * represents a significant difference at $P < 0.05$; ** represents a highly significant difference at $P < 0.01$ and "ns" indicates non-significant difference between the control and treated plants.

in roots of *V. officinalis* elicited with 50 μ M. High

levels of MeJA may trigger further activation of downstream signaling components which results in increased expression of JA target genes and in enhanced accumulation of corresponding protein expression (Endt et al., 2002). Such dose-dependent effect of MeJA on the biosynthesis of secondary metabolites has been supported by the finding of Uddin et al. (2013) and Dini Torkamani et al. (2014).

The mechanism underlying JAs in triggering signal transduction pathways leading to transcriptional reprogramming of secondary metabolism pathway and enhancement of desired metabolites has been studied extensively (Afrin et al., 2015). It is found that JA-ZIM domain (JAZ) acts as an on/off switch in JA signaling. In the absence of the JA and its bioactive derivatives, JAZ recruits a set of transcription factors (TFs) acting in modulation of biosynthesis of secondary metabolites pathways, thereby limiting their activity (Chini et al., 2007). Upon perception of JA signaling, JAZ repressor is targeted for degradation by 26S proteasome-dependent proteolysis and ultimately releases MYC2 (Memelink, 2009). It has been recently revealed that regulators of terpenoid indole alkaloid (TIA) biosynthesis in *Catharanthus roseus* is regulated by a JA-responsive TFs cascade (Yang et al., 2013). These finding suggest that coordinated induction of *VoTPS1* and *VoTPS7* can be conducted by such regulatory network. However, expression pattern of *VoTPS3* was more or less unaffected by MeJA at both concentrations. It may be concluded that *VoTPS3*, as a putative terpene synthase, does not appear to have any significant influence on the biosynthesis of VA.

It is well known that MYC2 as a master switch activates early JA-responsive genes transcription including JAZ. Formation of complex MYC2- JAZ repressor deactivates MYC2 and subsequently weakens JA-dependent responses (Chico et al., 2008). Our data indicated MeJA at both concentrations increased transcripts levels of *VoTPS1* and *VoTPS7* for up to 24 h after elicitation. Further increase in the exposure time decreased transcription rate of both genes probably due to attenuated JA signaling. Herein we also showed that substantial VA content enhancement was associated with simultaneous increased expression of *VoTPS1* and *VoTPS7* suggesting that

intermediate valerenadiene and germacrene B/C/D (encoded by *VoTPS1* and *VoTPS7* respectively) may possibly be involved in the VA biosynthesis and this was also proposed previously (Yeo et al., 2013).

Overall, these results illustrated that induction of *VoTPS1* and *VoTPS7* expression and increase in VA content were more effectively enhanced with increasing concentration of MeJA from 50 to 100 μ M. Therefore, it is concluded that VA biosynthesis is probably regulated at the transcriptional level of *VoTPS1* and *VoTPS7* in roots of *V. officinalis*. However, the molecular mechanism underlying MeJA-mediated transcriptional regulation of these putative terpene synthases needs to be further studied.

Acknowledgements

The authors would like to acknowledge the central laboratory of Ramin Agriculture & Natural Resources University of Khouzestan for providing the necessary laboratory facilities.

References

- Afrin, S., J.J. Huang and Z.Y. Luo, 2015. 'JA-mediated transcriptional regulation of secondary metabolism in medicinal plants'. *Science Bulletin*, 60(12):1062–1072.
- Beaubru, G and G.E. Gray, 2000. 'A review of herbal medicines for psychiatric disorders'. *Psychiatric Services*, 51: 1130-1134.
- Bent, S., A. Padula, D. Moore, M. Patterson and W. Mehling, 2006. 'Valerian for sleep: a systematic review and meta-analysis'. *The American Journal of Medicine*, 119: 1005-1012.
- Chico, J.M., A. Chini, S. Fonseca and R. Solano, 2008. 'JAZ repressors set the rhythm in jasmonate signaling'. *Current Opinion in Plant Biology*, 11(5):486-94.
- Chini, A., S. Fonseca, G. Fernandez, B. Adie, J.M. Chico, O. Lorenzo, G. García-Casado, I. López-Vidriero, F.M. Lozano, M.R. Ponce, J.L. Micol and R. Solano, 2007. 'The JAZ family of repressors is the missing link in jasmonate signalling'. *Nature*, 448:666–671
- DiniTorkamani, MR., M. Jafari, N. Abbaspour and N. Safaie, 2014. 'Enhanced production of

valerenic acid in hairy root culture of *Valeriana officinalis* by elicitation'. *Central*

European Journal of Biology, 9(9): 853–863

Endt, DV., J.W. Kijne and J. Memelink, 2002.

'Transcription factors controlling plant secondary metabolism: what regulates the regulators?'. *Phytochemistry*, 61: 107–114.

Hampel, D., A. Mosandl and M. Wust, 2005.

'Biosynthesis of mono- and sesqui-terpenes in carrot roots and leaves (*Daucus carota* L.): metabolic cross talk of cytosolic mevalonate and plastidial methylerythritol phosphate pathway'. *Phytochemistry*, 66: 305-311.

Letchamo, W., W. Ward, B. Heard and D. Heard,

2004. 'Essential oil of *Valeriana officinalis* L. cultivars and their antimicrobial activity as influenced by harvesting time under commercial organic cultivation'. *Journal of Agricultural and Food Chemistry*, 52: 3915-3919.

Livak, K.J and T.D. Schmittgen, 2001.

'Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method'. *Methods*, 25: 402-408.

Memelink, J., 2009.

'Regulation of gene expression by jasmonate hormones'. *Phytochemistry*, 70: 1560–1570.

Morazzoni, P and E. Bombardelli, 1995.

'*Valeriana officinalis*: traditional use and recent evaluation of activity'. *Fitoterapia*, 66: 99-112.

Murashige, T and F. Skoog, 1962.

'A revised medium for rapid growth and bioassays with tobacco tissue cultures'. *Plant Physiology*, 15: 473-497.

Pandey, S. S., S. Singh, C.S. Babu, K. Shanker,

N.K. Srivastava, A.K. Shukla, 2016. 'Fungal endophytes of *Catharanthus roseus* enhance vindoline content by modulating structural and regulatory genes related to terpenoid indole alkaloid biosynthesis'. *Scientific Reports*, 6:26583.

Pfaffl M.W., 2001. 'A new mathematical for relative quantification in real time RT-PCR'.
Nucleic acids Research, 29:e45.

Santos, MS., F. Ferreira, A.P. Cunha, A.P. Carvalho and T. Macedo, 1994.

'An aqueous extract of valerian influences the transport of GABA in synaptosomes'. *Planta Medica*, 60: 278-279.

Takemoto, H., T. Yagura and M. Ito, 2009.

'Evaluation of volatile components from spikenard. Valerena 4.7(11).diene is a highly active sedative compound'. *Journal of Natural Medicines*, 63: 380-385.

Tania, L and C.F. Brian, 2004.

'In vitro activity of commercial valerian root extracts against human cytochrome P450 3A4'. *Journal of Pharmaceutical Sciences*, 7: 265-273.

Uddin, MR., A.A. Thwe, Y.B. Kim, W.T. Park, S.C. Chae, and S.U. Park, 2013.

'Effects of Jasmonates on Sorgoleone Accumulation and Expression of Genes for Sorgoleone Biosynthesis in Sorghum Roots'. *Journal of Chemical Ecology*, 39(6): 712–722.

Wang, Y., L. Jin, S. Yu, Q. Shi, Y. Gu and H. Kiyota,

2010. 'Chemical constituents of plants from the genus *Valeriana*'. *Mini-Review in Organic Chemistry*, 7: 161-172.

Yang, Z., B. Patra, R. Li, S. Pattanaik and L. Yuan,

2013. 'Promoter analysis reveals cis-regulatory motifs associated with the expression of the WRKY transcription factor CrWRKY1 in *Catharanthus roseus*'. *Planta*, 238(6):1039-49

Yeo, YS., S.E. Nybo, A.G. Chittiboyina, A.D. Weerasooriya, Y.H. Wang, E. Góngora-Castillo, B. Vaillancourt, C.R. Buell, D. DellaPenna, M.D. Celiz et al., 2013.

'Functional identification of valerena-1,10-diene synthase, a terpene synthase catalyzing a unique chemical cascade in the biosynthesis of biologically active sesquiterpenes in *Valeriana officinalis*'. *Journal of Biological Chemistry*, 288: 3163–3173.

Zhang, L., R.X. Ding, Y.R. Chai, M. Bonfill, E. Moyano et al., 2004. 'Engineering tropane biosynthetic pathway in *Hyoscyamus niger* hairy root cultures'. *Proceedings of the National Academy of Sciences*, 101: 6786–6791.



تقویت تولید والرینیک اسید در ریشه *Valeriana officinalis* بوسیله تغییرات رونویسی ژن‌های سزکوئی ترپن سینتاز حاصل از تیمار متیل جاسمونات

هنگامه طاهری* و منصور قسمتی

گروه زراعت و اصلاح نباتات، دانشگاه کشاورزی و منابع طبیعی رامین خوزستان، ایران.

* عهده دارمکاتبات: taheri@ramin.ac.ir

چکیده فارسی

Valeriana officinalis (والرین)، به عنوان یک گیاه دارویی، به دلیل خواص آرامبخشی و خواب آوری به طور گسترده مورد استفاده قرار می‌گیرد. سزکوئی ترپن والرینیک اسید (C15) بعنوان موثرترین ترکیب در خواص دارویی این گیاه شناخته شده است. جهت ارزیابی اثرات متیل جاسمونات (غلظتهای ۵۰ و ۱۰۰ میکرومول) در تغییر الگوی بیان ژن‌های موثر در بیوسنتز والرینیک اسید، فراوانی رونوشت ژن‌های سزکوئی ترپن سینتازهای شناخته شده در ریشه این گیاه بوسیله تکنیک PCR در زمان واقعی در یک دوره زمانی ۱۴۴ ساعته مورد بررسی قرار گرفت. همچنین محتوای والرینیک اسید نیز توسط تکنیک HPLC مورد ارزیابی قرار گرفت. بیشترین میزان والرینیک اسید (۱۲/۴۵ میلی گرم به ازای هر گرم وزن خشک گیاه) در تیمار ۷۲ ساعت بعد از اعمال تیمار ۱۰۰ میکرومول متیل جاسمونات با ۱۲ برابر افزایش نسبت به گیاه شاهد (۱/۰۳ میلی گرم به ازای هر گرم وزن خشک گیاه) مشاهده شد. همچنین متیل جاسمونات در یک رفتار وابسته به غلظت میزان رونویسی ژن‌های *VoTPS1* و *VoTPS7* را تقویت کرد، به این ترتیب که تیمار ۱۰۰ میکرومول (در زمان ۲۴ ساعت) نسبت به تیمار ۵۰ میکرومول در القاء بیان این ژن‌ها بسیار کارآمدتر بود. همبستگی بین تقویت بیان ژن‌ها و محتوای والرینیک اسید نشان می‌دهد که احتمالاً این ژن‌ها در بیوسنتز ترکیبات واسط در مسیر بیوسنتزی والرینیک اسید موثر باشند. در هر صورت بنظر می‌رسد تیمار متیل جاسمونات اثر معنی‌دار کمتری در القاء بیان ژن *VoTPS3* نسبت به ژن‌های *VoTPS1* و *VoTPS7* داشته است. این یافته‌ها بینش ما در مورد بیوسنتز موثرتر والرینیک اسید بوسیله تغییرات رونویسی ژن‌های سزکوئی ترپن سینتاز تحت تیمار متیل جاسمونات را بهبود بخشید.

کلمات کلیدی: *Valeriana officinalis*، سزکوئی ترپن سینتاز، الگوی بیان، PCR در زمان واقعی