





# Sleep Hygiene Practices and Their Relation to Sleep Quality in Medical **Students of Qazvin University of Medical Sciences**

Zohreh Yazdi<sup>1</sup>, Ziba Loukzadeh<sup>2</sup>, Parichehr Moghaddam<sup>1</sup>, Shabnam Jalilolghadr<sup>3\*</sup>

<sup>1</sup>Social Determinants of Health Research Center, Qazvin University of Medical Sciences, Qazvin, Iran <sup>2</sup>Industrial Diseases Research Center, Shahid Sadoughi University of Medical Sciences, Yazd, Iran. <sup>3</sup>Children Growth Research Center, Qazvin University of Medical Sciences, Qazvin, Iran

#### **ARTICLE INFO**

Article Type: Original Article

#### ABSTRACT

Article History: Received: 13 Mar. 2015 Accepted: 20 Oct. 2015 ePublished: 1 Jun. 2016 Keywords: Ver 13. Sleep hygiene practice Medical students Sleep quality

Introduction: Poor quality of sleep is a distressing and worrying condition that can disturb academic performance of medical students. Sleep hygiene practices are one of the important variables that affect sleep quality. The objective of this study was to assess association between sleep hygiene practices and sleep quality of medical students in Qazvin University of Medical Sciences.

Methods: In this descriptive-correlational study, a total of 285 medical students completed a self-administered questionnaire. Demographic data, sleep-wake schedule in weekday and weekend, and sleep duration were collected. Students' sleep quality was assessed by Pittsburg Sleep Quality Index (PSQI). Data were analyzed by SPSS

Results: Overall, 164 (57.5) of students had poor sleep quality. Mean global PSQI score and average score of four subscales were significantly higher in male than female. Regression analysis showed that male students ( $\beta$ =-0.85, P<0.05), students at senior level ( $\beta$ =-0.81, P<0.05), married students ( $\beta$ =-0.45, P<0.05), and those with improper sleep hygiene practices slept worse.

Conclusion: The findings of this study showed that the prevalence of poor sleep quality in medical students is high. Improper sleep hygiene behaviors might be a reason for poor quality of sleep in medical students.

Please cite this paper as: Yazdi Z, Loukzadeh Z, Moghaddam P, Jalilolghadr S. Sleep hygiene practices and their relation to sleep quality in medical students of Qazvin university of medical sciences. J Caring Sci 2016; 5 (2): 153-60. doi:10.15171/jcs.2016.016.

### Introduction

Adequate amount and good quality of sleep are basic human needs. Sleep- wake circadian rhythms are regulated by a pacemaker in brain and controlled by some external factors including light, temperature, and social interaction.<sup>1,2</sup> In addition, pattern of sleep and wakefulness have many variations in different subjects in terms of age, physiologic and psychological characteristics, somatic and disorders, psychiatric and demands of occupation. Furthermore these patterns are affected by social requirements of modern life in recent decades.3,4

Sleep specialists recommend that healthy people generally need about seven hours of sleep per day. But recent surveys in different

countries have indicated that the average of sleep duration is very less than this amount.<sup>5-7</sup> Prior studies performed all over the world have showed that different sleep disorders are associated with psychiatric and/or somatic problems as well as social life disorders.2-6 Sleep deprivation in leads to impaired mood, judgment, ability to learn and maintain information.<sup>1,2</sup> Moreover, it is related to different health problems including cardiovascular disease, obesity, diabetes, metabolic syndrome, and increased mortality.8,9

University students are one of the high risk groups for developing sleep disorders, with a high vulnerability especially in medical students.<sup>1,2</sup> The prevalence of sleep disturbances have been reported high in

\*Corresponding Author: Shabnam Jalilolghadr (MD), email: jalilolghadrshabnam@yahoo.com. This study was approved and funded by deputy of research in Qazvin University of Medical Sciences (Project number: 1003).

<sup>0 3</sup> (CC) © 2016 The Author(s). This work is published by Journal of Caring Sciences as an open access article distributed NC under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by-nc/4.0/). Non-commercial uses of the work are permitted, provided the original work is properly cited.

medical students.<sup>10</sup> The prevalence of poor sleep quality has been estimated about 20 to 60 percent in medical students. In these studies, sleep quality was associated with sex, years of education, perceived adequate amount of sleep, stress and concern about university tasks, and irregular sleep-work hours.<sup>11,12</sup>

Many causes have been suggested for this issue. Some of them include high daily and nightly workload, inadequate time for leisure activities, living conditions, and high level of stress due to intensive curriculum schedule.<sup>8-10</sup> Also, many students live away from their family in dorms. These students are more likely to have improper sleep-wake schedule that is harmful in many cases.<sup>13,14</sup>

Poor academic performance has been reported in medical students who suffer from sleep disorders.<sup>10</sup> Moreover, studies have shown that sleep deprivation increases the risk of medical errors in health care workers.<sup>15,16</sup>

Sleep hygiene practices is one of the important variables that affects sleep quality.<sup>17</sup> inappropriate sleep behaviors are harmful for sleep. These behaviors, which were introduced firstly by Peter Hauri are based on physiology of sleep.<sup>18</sup> Many activities are considered as proper sleep hygiene that promotes good sleep. Some of them include the followings: 1) avoiding caffeine, nicotine, and alcohol near the bed time, 2) avoiding napping as an occasional pattern, 3) maintaining a regular sleep and wake time, 4) keeping bed room comfortable and quiet, and 5) avoiding highly demanding activities in bedroom.<sup>19</sup>

Unfortunately, knowledge of medical students is insufficient about sleep hygiene practices.<sup>20</sup> Also, they are often unaware how poor quality of sleep deteriorates their academic performance and cognitive function.<sup>19,20</sup>

In a survey, knowledge and practice of more than 900 medical students on sleep hygiene practices were assessed. Only 50% of participants answered correct response. Also, students' knowledge on sleep hygiene was positively related to sleep hygiene practices.<sup>21</sup> According to issues mentioned above, the objective of this study was to assess the association between sleep hygiene practices and sleep quality of medical students in Qazvin University of medical sciences, Qazvin, Iran.

### Materials and methods

descriptive-correlational study This was conducted in Qazvin University of medical sciences from March to September, 2012. All medical students were included in the study. None of them had any history of sleep or medical problems. Purpose and methods of the study was clearly explained to students and informed consent was obtained from them. The demographic characteristics of students including age, sex, body mass index (BMI), type of accommodation (living in dormitory, renting houses, and parent-owned houses) and marriage status were assessed by selfadministered questionnaire. All students were also asked about their routine sleep patterns. Information about sleep included total sleep duration, total time in bed, sleep latency, number of awakenings during the night, and naptime during the day. This information was collected on week days and weekends.

To assess the prevalence of good sleep hygiene practices in students, we asked 10 questions. Some questions were as follow: "During the past month, have you taken a nap during daytime occasionally?", "During the past month, have you smoked during two hours before bedtime?", and "Within the past month, have you eaten heavy night meals near the bedtime?" All the questions about sleep hygiene practices of students are presented in table 2. Students answered these questions with yes or no response.

Sleep hygiene questionnaire was designed based on expert's opinion in the field of sleep medicine and using previously published articles about sleep hygiene. Test-retest reliability was done by calculating intraclass coefficient (ICC). For this, fifty questionnaires were completed again by students after 2 weeks. Result from test-retest has shown that ICC was 0.91 and Chronbach's alpha coefficient was 0.78.

Persian version of Pittsburg Sleep Quality Index (PSQI) was used to evaluate subjective sleep quality of participants. PSQI assesses participant's view about her or his quality sleep during one month ago. It constitutes seven subscales including subjective sleep quality, sleep latency, sleep duration, sleep efficiency, sleep disturbances, use of sleep medications, and daytime dysfunction. Each part have four score rating from 0 to 3 which indicates: never, once a week, twice a week, more than three times a week, and respectively. The range of total score is between 0-21 and scores more than 5 shows poor quality of sleep.<sup>22</sup> Reliability and validity of Persian form of PSQI has been assessed in previous published article and its chronbach's alpha was 0.83.23 Data were analyzed by SPSS Ver 13.

Student t-test was used to determine differences in sleep pattern between weekdayweekend, and between male and female participants. Chi-square test was used to assess differences between males and females in frequency of poor sleep hygiene practices. To determine predictive factors (demographic and lifestyle characteristics and sleep hygiene practices) for poor quality of sleep, we used linear regression model. P<0.05 was considered statistically significant.

### Results

Total number of students was 325, in which 285 students (response rate: 87.7) have responded to the questionnaires. Of 285 students who responded to questionnaire, 150 were female (52.6). Mean age of students was 22.8  $\pm$ 1.74 years (ranged from 20 to 27 years).

Mean of Body Mass Index (BMI) was  $21.9 \pm 0.9$ . Fifty six (19.6%) of students were current smokers, and most of the smokers were male (47 students). More than half (53.7%) of the students were at basic science level, and the others were junior (22.8%) and senior (23.5%). Only 32 (10.8%) medical students were married and others were single. A total of 135 (47.3%), 46 (16.1%), and 104 (36.5%) of students were lived in dorms, renting houses, and parent-owned houses, respectively.

Table 1 shows and compares average time of sleep parameters in weekdays and weekends in males and females. As the table shows sleep parameters were statistically different between the weekdays and weekend in term of sex. Table 2 shows the responses of students to 10 questions about sleep hygiene practices. According to the table, male students had poorer sleep hygiene practices than female students in four items (2, 3, 4, and 7). While, sleep hygiene practices of females were worse than males in two items (5 and 6).

Detail results of PSQI questionnaire are shown in table 3. Mean global PSQI score and average score of four subscales among males were significantly higher than females. Overall, 164 (57.5%) of students (52% of female *vs.* 63.7% of male) had poor sleep quality (total score of them was more than 5).

Table 4 shows linear regression model for detecting predictive variables for poor sleep quality. According to the model, males ( $\beta$ =-0.85, P<0.05), students at senior level ( $\beta$ =-0.81, P<0.05), married students ( $\beta$ =-0.45, P<0.05), and students who had six types of improper sleep hygiene practices reported worse sleep.

Sleep variables <sup>*</sup>	Females		Pa	Males		Pb	Pc
	Week day	Weekend		Week day	weekend		
Time in bed	390 (34)	425 (57)	0.002	365 (28)	379 (31)	0.02	0.001
Sleep latency	18.2 (7.1)	23.4 (6.3)	0.04	22.4(6)	21.7 (4)	0.09	0.03
Total sleep time	371 (32)	412 (21)	0.01	361 (27)	375 (53)	0.005	0.07
Number of awakening during night	1.6 (0.3)	0.9 (0.2)	0.06	0.8 (0.2)	0.9 (0.3)	0.08	0.03
Naptime during the day	26.1 (2.4)	22.3 (3.8)	0.003	18.2(5.1)	21.3 (3.5)	0.03	0.07

#### Table1. Comparison of sleep patterns between females and males

\* Minutes, <sup>a</sup>P-value for differences between weekday and weekend variables in females. <sup>b</sup>P-value for differences between weekday and weekend variables in males. <sup>c</sup>P-value for differences between sleep pattern variables between males and females.

Ī

Sleep hygiene practices (SHP) of participants	Fen	nales	Ma	ales	р
	Yes	No	Yes	No	
	N (%)	N (%)	N (%)	N (%)	
Take a nap during daytime occasionally	34 (22.6)	116 (77.4)	46 (34)	89 (66)	0.08
Smoke within 2 hours before bedtime	9 (6)	141 (94)	47 (34.8)	88 (65.2)	< 0.001
Eat heavy night meals near the bedtime	43 (28.6)	107 (71.4)	60 (44.4)	75 (55.6)	0.02
perform dynamic physical activities within 2 hours	23 (15.3)	127 (84.7)	39 (28.9)	96 (71.1)	0.01
before asleep					
Go to bed thirsty or hungry	37 (24.6)	113 (75.4)	25 (18.5)	110 (81.5)	0.004
Think about day's events and worries at bed	75 (50)	75 (50)	39 (28.9)	96 (71.1)	< 0.001
Engage yourself with highly demanded activities near	57 (38)	93 (62)	71 (52.6)	64 (47.4)	0.03
to bedtime, like watching exciting movies					
Use your bed for activities other than sleep	48 (32)	102 (68)	35 (25.9)	100 (74.1)	0.07
Sleep in uncomfortable bedroom (in term of	23 (15.3)	127 (84.7)	31 (22.9)	104 (77.1)	0.19
temperature, light, noise)					
Sleep and wake at the different time during the week	51 (34)	99 (66)	42 (31.1)	93 (68.9)	0.12

T 11	- 0	<b>^</b>		- f - 1	1		1	£ 1	1	
Iani	p Z	( omns	irison i	or sieen.	nvoiene	nractices	netween	temales	ana	males
I GOL		compt	u ison	or sicep	in y gione	practices	bet ween	remaies	unu	marco
				1	20	1				

Table 3. Pittsburgh Sleep Quality Index (PSQI) global and component means and standard deviations

Variables	Female	Males	Р
	Mean (SD) 🧖	Mean (SD)	
Global score of PSQI	5.82 (0.9)	6.64 (1.1)	0.003
Subjective sleep quality	1.27 (0.4)	1.49 (0.5)	0.02
Sleep latency	0.9 (0.3)	0.9 (0.5)	0.09
Sleep duration	1.34 (0.5)	1.51 (0.4)	0.009
Habitual sleep efficiency	0.31 (0.08)	0.36 (0.1)	0.005
Sleep disturbance	0.85 (0.2)	1.22 (0.4)	0.001
Use of sleep medications	0.24 (0.07)	0.14 (0.09)	0.07
Daytime dysfunction	0.91 (0.3)	1.02 (0.5)	0.21

Table 4. Linear regression of the factors predicting sleep quality of medical students

Variables	R	t	Standardized β
Age	0.21	1.3	0.01
Gender	$0.24^{*}$	-1.9	-0.85*
Body mass index	-0.56	1.07	0.23
Type of accommodation	-0.34	0.41	0.32
Years of education	-0.65*	-1.8	-0.81*
Marriage	$-0.78^{*}$	-0.79	-0.45*
Total sleep time	-0.3	1.8	0.21
Take a nap during daytime occasionally	0.28	-0.73	-0.09
Smoke within 2 hours before bedtime	$0.61^{*}$	1.6	$0.86^{*}$
Eat heavy night meals near the bedtime	-0.09	0.46	0.08
perform dynamic physical activities within 2 hours before a	$0.59^{*}$	0.81	$0.65^{*}$
sleep			
Go to bed thirsty or hungry	$0.27^*$	1.9	$0.76^{*}$
Think about day's events and worries at bed	$0.75^{*}$	1.13	$0.45^{*}$
Engage yourself with highly demanded activities near to	$0.63^{*}$	1.7	$0.51^{*}$
bedtime like watching exciting movies			
Use your bed for activities other than sleep	-0.09	-0.53	-0.08
Sleep in uncomfortable bedroom (temperature, light, noise)		1.02	0.26
Sleep and wake at the different time during the week	$0.58^{*}$	2.5	$0.61^*$
P<0.05			

### Discussion

The findings of this study showed that the prevalence of poor sleep quality in medical students is generally, more than half of the students participating in the study suffered from poor quality of sleep. Also, complaint from this problem was more prevalent in male students than females. These results are consistent with most of the studies have been performed in other countries such as Ethiopia<sup>3</sup>, Saudi Arabia,<sup>10</sup> and America.<sup>20,21</sup>

However, some studies reported a lower prevalence of poor sleep quality in students Brazil, 24,25 and such as Palestinian students.<sup>26</sup> The variation between results from different studies may be related to differences in cultural habits, socioeconomic status and university characteristics of different countries. In the other words, in studies that have examined students' sleep quality during exam periods, the situation was worse. In a study conducted on 162 students at Zanjan University of medical sciences in Iran during exam period, 86.4% of students had poor quality of sleep.<sup>27</sup>

Current study also showed that sleep quality of males was worse than females. Results of other studies conducted in this area are inconsistent. Cheng and her assessed sleep quality colleagues of incoming university students. They found that poor sleep quality was significantly associated with female gender.<sup>28</sup> This is also consistent with other studies which have been reported high prevalence of morning fatigue and daytime sleepiness in female than male students.<sup>25-28</sup> This may be due to the fact that females physiologically need more time for sleep.<sup>15,25</sup> It should be noted that this study assessed incoming students (newly signed in) and this leads to discrepancies between the results. In some other studies, no difference was reported between males and females in terms of sleep quality.<sup>12,14</sup> Based on present results, male students had worse sleep hygiene practices than females. It could be a reason for higher prevalence of poor sleep quality in this group.

Other finding of this study showed that the mean sleeping time of participants in the study was less than recommended sleep amounts. In previous studies, association between short sleep duration with various health problems has been reported.<sup>8,9</sup>

Based on the other results, sleep hygiene practices for males was lower than females. Smoking close to bedtime, eating heavy dinner, performing dynamic physical activity, and highly demanded activities before bedtime (like watching exciting movies) were higher among males. Instead, females were more likely to go to bed thirstily and hungry. Furthermore, thinking about daily events and worry thoughts were more prevalent in females. According to the results, improper sleep hygiene behaviors might be a reason for poor quality of sleep in medical students. These results are in line with previous studies.<sup>11-</sup> 14,17,18

Improper behavior of sleep hygiene and poor sleep quality can ultimately lead to poor academic performance and increased medical errors in medical students.<sup>13</sup> A study conducted in Hong Kong indicated that students' awareness of sleep hygiene practices was low. In this study, undesirable sleep hygiene practices were significantly associated with poor quality of sleep.<sup>18</sup>

Good knowledge of students about sleep hygiene practices is not sufficient for improving sleep quality as reported by Brown et al. In fact, implementing and practicing these principles would help to improve sleep quality.<sup>20</sup>

Males, students at senior level, married students, and those with improper sleep hygiene practices had worse sleep. Sleep hygiene practices is shown to be associated with a higher prevalence of insomnia in other studies.<sup>13,14,19</sup>

Current study results showed that study, senior students had worse sleep quality. Poor quality of sleep in senior students can be due to their shift work schedules. Also, it is assumed that junior students may have developed better coping strategies for their university needs.<sup>29</sup> Similar results have been reported in other studies.<sup>13,14</sup>

This study has some limitations too. First the study population were the medical students on Qazvin University of medical sciences and do not cover the all of the Iranian medical students. Secondly, we used self-reported questionnaire for detecting variables, which could have been subject to bias. More subjective instrument like sleep diary, will give more accurate avoid prolonging results. То the questionnaire, authors did not assess some important variables influencing sleep such as stress, anxiety and depression.

## Conclusion

In conclusion, sleep hygiene is one of fundamental factors affecting sleep quality. Current findings recommend conducting further interventional studies for measuring the effects of educational programs on sleep hygiene and academic performance. Moreover, assessing changes of academic performance in students after interventional programs will be helpful.

# Acknowledgments

We would like to thank all medical students in Qazvin University of Medical Sciences participated in this study. The authors wish to thank the staff of the Center for Clinical Researches at Qazvin Children Hospital, affiliated to Qazvin University of Medical Sciences for their help in preparing this paper.

### **Ethical issues**

None to be declared.

# **Conflict of interest**

The authors declare no conflict of interest in this study.

# References

- Dewald JF, Meijer AM, Oort FJ, Kerkhof GA, Bogels SM. The influence of sleep quality, sleep duration and sleepiness on school performance in children and adolescents: A meta-analytic review. Sleep Medicine Reviews 2010; 14 (3): 179-89. doi <u>: 10.1016/j.smrv.2009.10.004</u>
- Vidyashree HM, Patil PP, Moodnur V, Singh D. Evaluation and comparison of sleep quality among medical and yogic students- A questionnaire based study. National Journal of Physiology, Pharmacy and Pharmacology 2013; 3 (1): 71-4. doi: <u>10.5455/njppp.2013.3.71-74</u>
- 3. Lemma S, Gelaye B, Berhane Y, Worku A, Williams MA. Sleep quality and its psychological correlates among university students in Ethiopia: a crosssectional study. BMC Psychiatry 2012; 12: 237. doi:10.1186/1471-244X-12-237
- 4. Augner C. Associations of subjective sleep quality with depression score, anxiety, physical symptoms and sleep onset latency in students. Cent Eur J Public Health 2011; 19 (2): 115-17.
- Smolensky MH, Milia LD, Ohayon MM, Philip P. Sleep disorders, medical conditions, and road accident risk. Accident Analysis and Prevention 2011; 43 (2): 533-48. <u>doi</u>: <u>10.1016/j.aap.</u> <u>2009.12.004</u>
- Anderson C, Horne JA. Do we really want more sleep? A population based study evaluating the strength of desire for more sleep. Sleep Medicine 2008; 9 (2): 184-87. doi: 10.1016/j.sleep.2007.02.006
- Ferrara M, De Gennaro L. How much sleep do we need? Sleep Medicine Review 2001; 5 (2): 155-79. doi: 10.10 53/smrv.2000.0138
- Eller T, Aluoja A, Vasar V, Veldi M. Symptoms of anxiety and depression in Estonia medical students with sleep problems. Depression and Anxiety 2006; 23 (4): 250-56. doi: 10.1002/da.20166

- 9. Owens JA. Sleep loss and fatigue in medical training. Curr Opin Pulm Med 2001; 7: 411-18.
- Abdulghani HM, Alrowais NA, Bin-Saad NS, Al-Subaie NM, Haji AM, Alhagwi AL. Sleep disorder among medical students: relationship to their academic performance. Medical Teacher 2012; 34 (Suppl1): 37-41. doi: 10.3109/0142159X. 2012.656 749.
- Preišegolavičiūtė E, Leskauskas D, Adomaitienė V. Associations of quality of sleep with lifestyle factors and profile of studies among Lithuanian students. Medicina (Kaunas) 2010; 46 (7): 482-9.
- Mesquita G, Reimão R. Quality of sleep among university students: effects of nighttime computer and television use. Arq Neuropsiquitr 2010; 68 (5): 720-25. doi:<u>10.1590/S0004-282X20100005000</u> 09
- Kabrita CS, Hajjar-Muça TA, Duffy JF. Predictors of poor sleep quality among Lebanese university students: association between evening typology, lifestyle behaviors, and sleep habits. Nature and Science of Sleep 2014; 2014 (6): 11-18. doi: 10.2147/NSS.S55538
- Angelone AM, Mattei A, Sbarbati M, Di Orio F. Prevalence and correlates for self-reported sleep problems among nursing students. J Prev Med Hyg 2011; 52 (4): 201-8.
- 15. Yazdi Z, Loukzadeh Z, Abbasi M, Ziaeeha M, Mahmoodi M. Prevalence of insomnia and its relation with academic performance of medical students in Qazvin University of Medical Sciences. The Journal of Medical Education and Development 2013; 6 (1): 19-25. (Persian)
- 16. Kramer M. Sleep loss in resident physicians: the cause of medical errors? Front Neurol 2010; 1 (128): 1-10. doi: 10.3389/fneur.2010.00128. eCollection 2010.

- 17. Bahmman AS, Al-Khairy OK, Al-Taweel AA. Sleep habits and patterns among medical students. Neurosciences (Riyadh) 2005; 10 (2): 159-62.
- Suen LK, Tam WW, Hon KL. Association of sleep hygiene-related factors and sleep quality among university students in Hong Kong. Hong Kong Med J 2010; 16 (3): 180-85.
- Brick CA, Seely DL, Palermo TM. Association between sleep hygiene and sleep quality in medical students. Behav Sleep Med 2010; 8 (2): 113-21. doi: <u>10.</u> <u>1080/15402001003622925</u>.
- Brown FC, Buboltz WC, Soper B. Relationship of sleep hygiene awareness, sleep hygiene practices, and sleep quality in university students. Behavioral Medicine 2002; 28 (1): 33-8. doi:10.108 0/08964280209596396
- 21. Hicks RA, Lucero-Gorman, Bautista J, Hicks GJ. Ethnicity, sleep hygiene knowledge, and sleep hygiene practices. Percept Mot Skills 1999; 88 (3 pt 2): 1095-6.
- 22. Buysse DJ, Reynols CF, Monk TH, Berman SR, Kupfer DJ. The pittsburghh sleep quality index: a new instrument for psychiatric practice and research. Psychiatry Res 1989; 28 (2): 192–213.
- 23. Ebrahimi A, Ghalebandi F, Salehi M, Kafian Tafti AR, Vakili Y, Akhlaghi Farsi E. Sleep parameters and the factors affecting the quality of sleep in patients attending selected clinics of rasoul-e-Akram Hospital. Razi Journal of Medical Sciences 2008; 15 (58) :31-38
- 24. Lima AMA, Varela GCG, Silveira HACS, Parente RDG, Araujo JF. Evening chronotypes experience poor sleep quality when taking classes with early starting times. Sleep Sci 2010; 3 (1): 45–8.
- 25. Medeiros ALD, Mendes DBF, Lima PF, Araujo JF. The relationship between sleep-wake cycle and academic performance in medical students. Biological Rhythm Research 2001; 32

(2): 263–70. doi: <u>10.1076/brhm.32.2.26</u> <u>3.1359</u>

- 26. Sweileh WM, Ali IA, Sawalha AF, Abu-Taha AS, Zyoud SH, Al-Jabi SW. Sleep habits and sleep problems among Palestinian students. Child Adolesc Psychiat Mental Health 2011 15; 5: 25. doi: 10.1186/1753-2000-5-25
- 27. Aghajanloo A, Haririan H, Ghafourifard M, Bagheri H, Ebrahimi SM. Sleep quality of students during final exams in Zanjan University of Medical Science. Modern Care, Scientific Quarterly of Birjabd Nursing and Midwifery Faculty

2012; 8 (4): 230-37. (Persian)

- Cheng SH, Shih CC, Lee IH, Hou Yw, Chen KC, Chen KT, et al. A study on the sleep quality of incoming university students. Psychiatry Res 2012; 197 (3): 270–4. doi: <u>10.1016/j.psychres. 2011. 08.</u> 011
- Niemi PM, <u>Vainiomäki</u> PT. Medical students' distress--quality, continuity and gender differences during a six-year medical programme. Med Teach 2006; 28 (2): 136-41. doi: <u>10.1080/0142159</u> 0600607088

160 / Journal of Caring Sciences, June 2016; 5 (2), 153-160