



Original Article

Effect of Self-Management Program on Self-efficacy and Medication Adherence in Patients with Mechanical Heart Valve: a Randomized Clinical Trial

Leila Javan¹ , Anoshirvan kazemnejad² , Mahin Nomali¹ , Masoumeh Zakerimoghadam^{1*} 

¹Department of Critical Care Nursing, School of Nursing and Midwifery, Tehran University of Medical Sciences, Tehran, Iran

²Department of Biostatistics, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

ARTICLE INFO

Article type: Original article

Article History:

Received: 6 Oct. 2017

Accepted: 22 Oct. 2018

ePublished: 1 Dec. 2019

Keywords:

Heart valves, Self care, Self efficacy, Medication Adherence

*Corresponding Author:

PhD in Nursing. Email:
 mzakerimo@tums.ac.ir

ABSTRACT

Introduction: Patients with mechanical heart valve need anticoagulant therapy to prevent thrombotic events. The treatment interacts with some foods and drugs. The aim was to evaluate the effect of self- management program on self-efficacy and medication adherence in patients with mechanical heart valve.

Methods: This was a randomized controlled trial. eighty eligible patients, with the ability to read and speak in Farsi, aged between 15 to 60, were included in the study from the cardiac surgery clinic in Imam Khomeini hospital affiliated to Tehran University of Medical Sciences (Tehran, Iran) and randomly allocated to intervention and control groups. The participants had no history of psychiatric disorders, had undergone valve replacement surgery at least one year before the study, and were being treated with Warfarin. The intervention was a combination of 2 one- hour self-management education via small groups with 3 to 5 members, self-management educational booklets, and weekly call follow- ups for 8 weeks about 10- 15 minutes. The control group received no intervention. Self- efficacy was the primary outcome and medication adherence, Prothrombin Time (PT), and International Normalized Ratio (INR) were secondary outcomes. Data were analyzed using SPSS13.

Results: Although the mean of self-efficacy and medication adherence, PT, and INR values were not different between the two groups at baseline, they improved significantly following the program.

Conclusion: Self-management program had a positive effect on self-efficacy and medication adherence of patients with mechanical heart valve.

Citation: Javan L, kazemnejad K, Nomali M, Zakerimoghadam M. Effect of self- management program on self-efficacy and medication adherence in patients with mechanical heart valve: a randomized controlled trial. *J Caring Sci* 2019; 8 (4): 207-11. doi:10.15171/jcs.2019.029

Introduction

Valvular heart disease (VHD) is prevalent in developed and developing countries with etiology of degenerative¹ and rheumatic valvular disease by Staphylococci (Group A).^{1,2} In the United States, VHD accounts for 10% to 20%³ and in Iran, it comprised 7.2% of all cardiac surgeries.⁴

From reported complications, thrombosis, thromboembolism, and anticoagulant- related complications account for 75% of all complications⁵ which usually occur six months after surgery and necessitates the post operation follow- ups⁵ Moreover, they are the leading causes of morbidity and mortality⁶ imposing a psychological burden and a negative effect on the quality of life.⁵

Although the long term durability is the main advantage of mechanical heart valves, there is need for Vitamin K antagonists (VKAs) with inherent risk of thromboembolism and bleeding events in contrast.⁷

Warfarin and other VKAs interact with some foods and drugs, which makes the management of oral anticoagulant therapy challenging for both patients and health care providers.⁸ Therefore, the management of oral anticoagulant therapy is a key issue among patients with mechanical heart valve⁷ and the therapy should be individualized by focusing on patient education and monitoring⁹ in order to reduce the risk of bleeding and Thromboembolism events.^{9,10} One of the strategies is self-management program that empowers patients to monitor

and adjust treatment at home.¹¹ Previous studies have shown that self- management program improves the INR control, quality of life,¹² self-efficacy,¹³ medication adherence, and adjustment of treatment¹⁴ and leads to lower risk of all- cause mortality rate,^{15,16} and thromboembolism and major bleeding.¹⁵

Although most patients with a mechanical heart valve may benefit from self-management program,⁷ Iranian patients with mechanical heart valve have a permanent fear and concern about INR fluctuation, its regulation, and correct use of VKAs during their treatment.¹⁷ On the other hand, insufficient support can make life difficult for them.¹⁸ Thus, because the self-management program appears at least better than the routine care,¹¹ and there is crucial need for further randomized controlled trials to assess the efficacy of self-management program in order to consider it as the standard care,¹⁵ and in view of lack of this program in our health care setting so as to overcome the patients` fear and concerns and improve clinical outcomes, the aim of this study was to evaluate the effect of self- management program on self-efficacy and medication adherence of patients with mechanical heart valves.

Materials and methods

This was a parallel randomized controlled trial that has been approved by the institutional review board (IRB) of Tehran University Medical Sciences (TUMS) (No:

130/1225/93/d), and IRCT code IRCT201406154443N13. With Helsinki ethical principles for human research considered during the study period.

The participants were patients aged between 15 to 60 years, who had undergone valve replacement surgery at least one year before the study and were being treated with Warfarin, with the ability to read and speak Farsi, no history of psychiatric disorders. The patients who experienced the drug side effects during the study period, were discharged before completing the two-educational sessions, did not answer the phone calls during the call follow-ups, or were not willing to continue the study were all excluded from the study.

Therefore, 80 eligible patients with mechanical heart valves were included in the study from the cardiac surgery clinic in Imam Khomeini hospital affiliated to TUMS in 2014.

In this trial, the intervention consisted of self-management education, and weekly call follow-ups for 8 continuous weeks, each lasting from 10 to 15 minutes. In order to provide self-management education, the patients were divided into groups of 5 to 7 members and received 2 one-hour self-management education sessions at 2 to 3-day intervals. Self-management education and booklets were about the nature of the disease, medication management, drugs' side effects, and actions needed to be taken to prevent drug side effects, food and drug interactions, having Warfarin card, prevention of actions that lead to complications, physical activity and stress and anxiety reduction methods. The content validity of the educational booklet was evaluated by three nursing and midwifery faculty members of TUMS. While the patients in the control group received no intervention.

The outcomes of this trial were self-efficacy as a primary outcome, and medication adherence, prothrombin time (PT) and international normalized ratio (INR) were the secondary outcomes. In order to including 31 items with five-point Likert from extremely high to extremely low, with the higher scores indicating the higher self-efficacy behavior.^{19,20} The questionnaire was translated into Farsi by forward-backwards translation method and its content validity was approved evaluate the patient's self-efficacy, a researcher-constructed self-efficacy questionnaire was used by 10 faculty members in the school of nursing and midwifery affiliated to TUMS. It had an acceptable reliability coefficient, too (Cronbach's alpha= 0.82). The patients were asked to complete the questionnaire themselves at baseline and at the end of the study. Medication adherence was another study outcome which was evaluated by a medication adherence questionnaire^{21,22} consisting of 22 items with five-point Likert from always to never, where the higher scores were considered as higher medication adherence. 10 nursing and midwifery faculty members approved the content validity and Table 1 shows the demographic and clinical characteristics of the patients in the intervention and control groups. According to this table, no Significant difference for clinical and demographic data were found between the two groups at baseline ($P>0.05$) (Table 1).

Comparison of the mean and standard deviation of the

Cronbach's alpha was 0.85. The medication adherence questionnaire was completed by patient's self-report at baseline and at the end of the study. In addition, PT and INR were measured by BIOLABO kit (France) with therapeutic range of INR between 2 and 3.5 at baseline and at the end of the study.

Demographic and clinical variables including age, sex, education level, job, income, insurance, smoking, family history of cardiovascular diseases (CVDs), consumption of other medications, and post-operation duration were among the study variables which were recorded at baseline for each patients based on their self-report and clinical records.

The collected data were analyzed using SPSS (version 13.0, Chicago, IL, USA). We described demographic and clinical data by descriptive statistics such as frequency, partial frequency percentage, the mean and standard deviation Mean (SD). In order to compare the means between the intervention and control groups, independent-samples t-test and chi-squared test were used to compare the frequency between the two groups. The significance level of statistical tests was less than 0.05.

Results

In this trial, 88 eligible patients were recruited during the enrolment phase and data related to 80 patients were analyzed. The total attrition rate was 10% as it was expected a priori. The most common causes of exclusion from the study were "not answering the call-follow-ups" and "no willingness to continue the study" during the follow-up period. The study process from the enrollment to analysis has been shown in CONSORT 2010 flow diagram (Figure 1).




Figure 1. Flowchart of study

study outcomes between the intervention and control Groups has been shown in Table 2. According to table 2, although the mean and standard deviation of self-efficacy, medication adherence, and PT and INR were not significantly different between the two groups at baseline ($P> 0.05$), they improved significantly in the intervention

Table 1. Demographic and clinical characteristics of patients in the intervention and control groups (n=40)

Demographic and clinical characteristics	Intervention N (%)	Control N (%)	P
Age (Year)			0.36
15-30	7 (17.5)	4 (10)	
31- 45	14 (35)	11 (27.5)	
46- 60	19 (47.5)	25 (62.5)	
Sex			0.62
Male	14 (35)	23 (57.5)	
Female	26 (65)	17 (42.5)	
Education level			0.86
Elementary school	16 (40)	18 (45)	
High school	12 (30)	12 (30)	
Diploma & Academic	12 (30)	10 (25)	
Job			0.81
Yes	22 (55)	21 (52.5)	
No	18 (45)	19 (47.5)	
Income			0.75
Enough	16 (40)	15 (37.5)	
Not enough	24 (60)	25 (62.5)	
Insurance			0.49
Yes	34 (85)	36 (90)	
No	6 (15)	4 (10)	
Smoking			0.36
Yes	5 (12.5)	8 (20)	
No	35 (87.5)	32 (80)	
Family history of CVDs [€]			0.36
Yes	18 (45)	14 (35)	
No	22 (55)	26 (65)	
Consumption of other medications			0.34
Yes	25 (62.5)	29 (72.5)	
No	15 (37.5)	11 (27.5)	
Post-operation duration (Month)			0.44
< 1	11 (27.5)	7 (17.5)	
2-6	14 (35)	13 (32.5)	
7 – 12	15 (37.5)	20 (50)	

[€]CVDs: Cardiovascular Diseases

Compared with the control group following the program (P<0.05).

Table 2. Comparison of mean and standard deviation of study outcomes between the intervention and control group (n=40)

Study outcomes	Intervention Mean(SD)	Control Mean(SD)	P
Self-efficacy			
Before	93.22(12.05)	96.22(14.99)	0.327
After	118.20(12.94)	100.40(16.01)	0.0001
Medication adherence			
Before	60.02(6.26)	79.60(9.04)	0.813
After	94.97(8.55)	80.80(10.21)	0.0001
PT (Second)			
Before	17.78(2.91)	18.21(3.45)	0.530
After	19.94(3.31)	17.56(4.11)	0.008
INR			
Before	1.89(0.49)	1.95(0.62)	0.640
After	2.29(0.67)	1.94(0.45)	0.019

Discussion

In this trial, the effect of self-management program on self-efficacy and medication adherence of patients with mechanical heart valve was evaluated. According to the

result of the present study, the patients` self-efficacy improved significantly following the intervention. Previous studies assessed the effect of the intervention on self- efficacy among patients with multiple sclerosis, care following the intervention, too.^{13,19,23,24} In contrast, in Elzen et al., study, although the mean of self-efficacy score increased following the intervention, this difference was not statistically significant which may be due to the aged study population and different disease conditions.²⁵

Reduce the health care costs.²⁶ the positive effect of self-management program on patients` medication adherence has been shown in our study which was consistent with previous studies where the mean of medication adherence was significantly higher in the intervention group in comparison to the control group.^{22,27}

Education about anticoagulant self-management can result in the patients` satisfaction, quality of life improvement, anticoagulation control, and reduction in thromboembolic events and mortality.²⁸⁻³⁰ in the present study, the mean of PT and INR were significantly higher in the intervention group in comparison to the control group. In Jeon and Park study in 2015, PT and INR were maintained in the therapeutic range, following the intervention among patients with cardiac valve replacement.¹³ In addition, educational intervention plays a key role in controlling INR and its maintenance in normal range³¹ In contrast, Sedri et al., indicated that PT and INR were not significantly different between the two groups of intervention (interactive and non- interactive short message service) and control²² which may be due to the type of intervention and follow- up duration.

Conduction of this research project in a tertiary referral hospital that covers all patients with different health care behaviors was the strength of this study. The main limitation of this study was the presence of patients in educational sessions in order to receive the intervention that made patients discontinue the program. Thus, it is recommended the self-management program be provided through internet or by mobile health applications to improve patients` involvement in their care.

Conclusion

Self-management program had a positive effect on self-efficacy and medication adherence of the patients with mechanical heart valves. Therefore, this intervention can be applied by nurses as a part of the discharge plan in order to reduce the outcomes of patients with mechanical heart valve and improve their self-efficacy and medication adherence.

Acknowledgments

This article was extracted from the MS thesis of Ms. Leila Javan for the field of critical care nursing at Tehran university of medical sciences (Tehran, Iran) .The authors thank IRB and Research Ethics Committee (REC) of TUMS for approving the research projects.

Ethical issues

None to be declared.

Conflict of interest

The authors declare no conflict of interest in this study.

References

1. Jung B, Vahanian A. Epidemiology of valvular heart disease in the adult. *Nat Rev Cardiol* 2011; 8 (3): 162-72. doi: 10.1038/nrcardio.2010.202
2. Shahin Imani M, Asghar Ramezani M, Saeid Hosseini M, Babak Nasiri M, Abbasnezhad M. A multi-center study of the pathologies of valvular heart diseases: 5 year analysis of more than 3400 patients in Iran. *Journal of Cardiovascular and Thoracic Research* 2010; 2 (2): 21-7.(Persian)
3. Maganti K, Rigolin VH, Sarano ME, Bonow RO. Valvular heart disease: diagnosis and management. *Mayo Clin Proc* 2010; 85 (5): 483-500. doi: 10.4065/mcp.2009.0706
4. Karimi A, Ahmadi S, Davoodi S, Marzban M, Movahedi N, Abbasi K, et al. First database report on cardiothoracic surgery in Tehran Heart Center. *Iran J Public Health* 2008 ; 37 (2) :1-8.
5. Kulik A, Rubens FD, Wells PS, Kearon C, Mesana TG, van Berkem J, et al. Early postoperative anticoagulation after mechanical valve replacement: a systematic review. *Ann Thorac Surg* 2006; 81 (2): 770-81. doi: 10.1016/j.athoracsur.2005.07.023
6. Groves P. Surgery of valve disease: late results and late complications. *Heart* 2001; 86 (6): 715-21. doi: 10.1136/heart.86.6.715
7. Christensen TD, Larsen TB. Dabigatran versus warfarin in patients with mechanical heart valves: comment. *J Thromb Haemost* 2014; 12 (3): 424-5. doi: 10.1111/jth.12497
8. Leiria TL, Lopes RD, Williams JB, Katz JN, Kalil RAK, Alexander JH. Antithrombotic therapies in patients with prosthetic heart valves: guidelines translated for the clinician. *J Thromb Thrombolysis* 2011; 31 (4): 514-22. doi: 10.1007/s11239-011-0574-9.
9. Snipelisky D, Kusumoto F. Current strategies to minimize the bleeding risk of warfarin. *J Blood Med* 2013; 4: 89-99.
10. Kaneko T, Aranki SF. Anticoagulation for Prosthetic Valves. *Thrombosis* 2013; 2013: 4. doi: 10.1155 /2013 /346752
11. Christensen TD, Johnsen SP, Hjortdal VE, Hasenkam JM. Self-management of oral anticoagulant therapy: a systematic review and meta-analysis. *Int J Cardiol* 2007; 118 (1): 54-61. doi: 10.1016/j.ijcard.2006.06.018
12. Hamad MAS, van Eekelen E, van Agt T, van Straten AH. Self-management program improves anticoagulation control and quality of life: a prospective randomized study. *Eur J Cardiothorac Surg* 2009; 35(2): 265-9. doi: 10.1016/j.ejcts.2008.10.020
13. Jeon HR, Park JS. Development and application of a self-management program based on Prothrombin INR monitoring for Patients with cardiac valve replacement. *J Korean Acad Nurs* 2015; 45 (4): 554-64. doi: 10.4040 /jkan.2015.45.4.554
14. Pozzi M, Mitchell J, Henaine AM, Hanna N, Safi O, Henaine R. International normalized ratio self-testing and self-management: improving patient outcomes. *Vascular Health and Risk Management* 2016; 12: 387-92. doi: 10.2147/VHRM.S85031
15. Christensen TD, Skjøth F, Nielsen PB, Maegaard M, Grove EL, Larsen TB. Self-management of anticoagulant therapy in mechanical heart valve patients: a matched cohort study. *Ann Thorac Surg* 2016; 101 (4): 1494-9. doi: 10.1016/j.athoracsur.2015.09.084
16. Christensen TD, Skjøth F, Nielsen PB, Maegaard M, Grove EL, Larsen TB. Self-Management of Anticoagulant Therapy in Mechanical Heart Valve Patients: A Matched Cohort Study. *Ann Thorac Med* 2016; 101 (4): 1494-9. doi.org/10.1016
17. Taghadosi M, Memarian R, Ahmadi F. Experiences of warfarin use among cardiac valve-replaced patients in Iran. *Iranian Journal Of Critical Care Nursing* 2014; 6 (4): 207-16. (Persian)
18. Taghadosi M, Memarian R, Ahmadi F. The Experiences of "Difficult Life" in Heart Valve Replaced Patients. *Iran Red Crescent Med J* 2014; 16 (8). doi : 10.5812/ircmj.19147
19. Kaveh Savadkooch O, Zakerimoghdam M, Gheyasvandian S, Kazemnejad A. Effect of self-management program on self-efficacy in hypertensive patients. *Journal of Mazandaran University of Medical Sciences* 2012; 22 (92): 19-28. (Persian)
20. Sol BG, van der Graaf Y, van der Bijl JJ, Goessens NB, Visseren FL. Self-efficacy in patients with clinical manifestations of vascular diseases. *Patient Educ Couns* 2006; 61 (3): 443-8. doi: 10.1016/j.pec.2005.05.011
21. Lambert TJ, Kane J, Kissling W, Parellada E. CERP–Centres of Excellence in Relapse Prevention. An international educational programme to enhance relapse prevention in schizophrenia. *Schizophrenia Research* 2010; 117 (2-3): 295-6. doi: 10.1016/j.schres.2010.02.494
22. Sedri N, Asadi Noughabi A, Zolfaghari M, Haghani H, Tavan A. Comparison of the effect of two types of short message service (interactive and non-interactive) on anticoagulant adherence of patients with prosthetic heart valves. *Journal of Nursing Education* 2014; 2(4): 1-11. (Persian)
23. Kafami F, Mohammadi F, Norouzi K, Rahgozar M. The effect of self-management program on the health status of multiple sclerosis patients. *Journal of Research Development In Nursing & Midwifery* 2012 ; 9 (1) 24 -33. (Persian)
24. Bodenheimer T, Lorig K, Holman H, Grumbach K. Patient self-management of chronic disease in primary care. *Jama* 2002; 288 (19): 2469-75. doi: 10.1001/jama.288.19.2469
25. Elzen H, Slaets JP, Snijders TA, Steverink N. Evaluation of the chronic disease self-management program (CDSMP) among chronically ill older people in the Netherlands. *Soc Sci Med* 2007; 64 (9): 1832-41. doi: 10.1016/j. socscimed.2007.02.008
26. Cutler DM, Everett W. Thinking outside the pillbox—medication adherence as a priority for health care reform. *N Engl J Med* 2010; 362 (17): 1553-5. doi: 10.1056 /NEJMp1002305.
27. Baljani E, Rahimi Zh, Heidari Sh, Azimpour A. The effect of self management interventions on medication adherence and life style in cardiovascular patients. *Journal of Hamadan Nursing and Midwifery Faculty* 2012; 20 (3): 58-68. (Persian)
28. Levi M. Self-management of anticoagulation. *Expert Rev Cardiovasc Ther* 2008; 6 (7): 979-85. doi: 10.1586 /14779072.6.7.979
29. Sawicki PTA structured teaching and self-management

- program for patients receiving oral anticoagulation: a randomized controlled trial. Working Group for the Study of Patient Self-Management of Oral Anticoagulation. *JAMA* 1999; 281 (2): 145-50. doi: 10.1001/jama.281.2.145
30. Garcia-Alamino JM, Ward AM, Alonso-Coello P, Perera R, Bankhead C, Fitzmaurice D, et al. Self-monitoring and self-management of oral anticoagulation. *Cochrane Database Syst Rev* 2010; 128 (1): 246. doi: 10.1590/S1516-31802010000400015
31. Khan TI, Kamali F, Kesteven P, Avery P, Wynne H. The value of education and self-monitoring in the management of warfarin therapy in older patients with unstable control of anticoagulation. *Br J Haematol* 2004; 126 (4): 557-64. doi: 10.1111/j.1365-2141.2004.05074.x