

A one year survey of motor vehicle accidents in Rafsanjan, Iran, in 2014

Esmaili A, PhD¹, Rezaeian M, PhD², Sanji M, MD³, Sheikh Fathollahi M, PhD⁴, Mortavazi SM, MD-MPH⁵, Ghasemi HR, MD-GP⁶, Vazirinejad R, PhD*⁷

1-Associated Prof., Dept. of Social Medicine, Occupational Environmental Research Center, Medical School, Rafsanjan University of Medical Sciences, Rafsanjan, Iran. 2. Professor, PhD of Epidemiology, Dept. of Epidemiology and Biostatistic, Occupational Environmental Research Center, Medical School, Rafsanjan University of Medical Sciences, Rafsanjan, Iran. 3- Assistant Prof., Dept. of Emergency, Rafsanjan University of Medical Sciences, Rafsanjan, Iran. 4- Assistant Prof., Dept. of Social Medicine and Occupational Environment Research Center, Medical School, Rafsanjan University of Medical Sciences, Rafsanjan, Iran. 5- Master of Public Health, Disaster & Emergency Management Center, Rafsanjan University of Medical Sciences, Rafsanjan, Iran. 6-General Practitioner, Medical School, Rafsanjan University of Medical Sciences, Rafsanjan, Iran. 7- Professor, PhD of Epidemiology, Social Determinants of Health Research Centre, Medical School, Rafsanjan University of Medical Science, Rafsanjan, Iran.

Abstract

Received: November 2015, Accepted: December 2015

Background: Injuries and accidents are one of the main problems in the present century. The purpose of this study was the identification and analysis of the characteristics of pre-hospital emergency motor vehicle accidents (MVA) in Rafsanjan, Iran, in 2014.

Materials and Methods: This cross-sectional descriptive study was performed on all patients recorded by pre-hospital emergency during a period of one year (from 21 March 2013 to 20 March 2014). Our data were obtained from a crash record book kept at pre-hospital emergency dispatch. The crash database variables include age and gender of victims, vehicle type, type of collision, response time, location, and crash hotspots, and severity of injury. Accident data have been analyzed in SPSS software.

Results: In 2014, 1878 injured patients were transported to the general hospital by pre-hospital emergency dispatch. The mean age of victims was 29.10 ± 15.65 years, with the age range of 1-95 years. Among these cases, 85.8% were men and 14.2% were women. The majority of the patients (56.7%) were in the age range of 15-29 years, 30-44 years (20.6%), and 46-65 years (10.8%), respectively. Motorcycle accidents were the major cause of injuries (48.6%), and 4 patients (0.2% of all cases) died on the accident scene.

Conclusions: In the present study, motorcycle accidents constitute the majority of cases of MVAs. Therefore, safety training targeted at motorcyclists would be one of the best interventions for the prevention and controlling of trauma due to MVAs. This intervention could be the most efficient among younger drivers.

Keywords: Injuries, Accidents, Motor vehicle, emergency.

Introduction

Injuries are amongst issues of grave concern within societies. They result in the deaths of countless individuals and disabilities in thousands of others. Many significant psychological disorders in families and economical burden on countries and communities are also associated with accidents and their resulting injuries (1-5). It was estimated that approximately 10% of all of injuries lead to death. It is also estimated that

16000 deaths daily worldwide are related to injuries. This figure is equivalent to 5.8 million deaths due to accidents each year, which is more than the total number of deaths due to HIV, tuberculosis, and malaria (3.8 million) (5,6). In many countries, injuries have been reported as the 3rd health problem after

* **Corresponding author:** Reza Vazirinejad, Social Determinants of Health Research Centre, Medical School, Rafsanjan University of Medical Science, Rafsanjan, Iran.
Email: rvazirinejad@yahoo.co.uk

cardiovascular diseases and cancer; furthermore, 90% of deaths from injuries occur in low and middle income countries (5, 6).

Motor vehicle accidents (MVAs) are the most common source of injury and death around the globe. In 2002, more than 1.18 million people died in MVAs, and approximately 15 million are injured every year in MVAs worldwide. About 70% of fatalities occur among young adults aged 15-44 years, and in many countries, MVAs have become the main cause of death among men in this age group. MVAs have become a major health burden in all societies in the 21st century (7-10).

In Iran, a large number of injuries and deaths due to MVAs have been reported in the last few years. Due to the high social and economic cost of road accidents and their physical and psychological effects on individuals and communities, today, traffic accidents is one of the most important topics of scientific research. Similarly, in Rafafsanjān MVAs (as the developing countries) are also social-economically very important (11, 12).

The purpose of this study was the identification and analysis of the characteristics of pre-hospital emergency MVA in Rafsanjan, Iran, during a period of one year (from 21 Mar 2013 to 20 Mar 2014).

Materials and Methods

The present study took place in Rafsanjan District, Kerman Province, 850 km South of Tehran, in the Southeast of Iran. This was a cross-sectional descriptive study. The study sample consisted of all patients referred to emergency wards during this time period. A checklist was designed based on the aim and objective of the study for the purpose of data collection. The checklist is comprised of two sections. The first section contains demographic information (e.g., age and gender). The second section of the checklist contains information about the vehicle type, type of collision, severity of injuries, response time, location and crash hotspots.

Injuries were classified into 3 groups in terms of severity (dead, major injury, and minor injury). Evaluation according to crash severity was performed by pre-hospital emergency technicians. For this reason, all technicians were trained before data collection on the evaluation of the victim's physical condition in terms of anatomic damage.

Data were analyzed using SPSS software (version 18, SPSS Inc., Chicago, IL, USA), and parametric (t-test, correlation test) and non-parametric analysis (chi-square, Fisher's exact test) tests were used where appropriate.

Results

During the year 2014, 1878 injured patients were transferred to general hospitals by pre-hospital emergency dispatch. In other words, 5.14 victims of city street accidents referred to the emergency department of hospitals every day. Of those, 85.8% were men and 14.2% were women (men to women ratio was 6.04:1). The mean age of victims was 29.10 ± 15.65 years, with age range of 1-95 years (min = 19, max = 95 years). The majority of the patients (56.7%) were in the age range of 15-29 years, 30-45 years (20.6%), and 46-65 years (10.8%), respectively. As the results show, the rate of injuries in men was more than six fold higher than that in women, and the majority of injured patients (85.3%) were below the age of 45 years. Data on the victims and their accidents is presented in table 1.

The percentages of victims based on the type of vehicles and their age is presented in table 2. As shown in table 2, the most frequent vehicle accidents were motorcycle accidents (n = 913, 48.6%) and rollovers (n = 396, 21.1%), respectively. Statistical analysis showed that there was a significant association between the age of victims and the vehicle they had been driving ($\chi^2 = 287.38$, df = 16, $P < 0.001$).

More than 50% of victims were in the age range of 15-29 (n = 1065), among which a considerable majority was injured in motorcycle accidents (57.9%, n = 617). Among victims in the age range of older than

65 years of age, the highest rate of accidents had taken place for pedestrians (47.3%).

Table 1: Demographic characteristics of motor vehicle accident victims and data on their accidents

Variables	All motor vehicle accidents	
	N	%
Gender		
Male	1612	85.8
Female	266	14.2
Age at census		
0-14	151	8.0
15-29	1065	56.7
30-45	386	20.6
46-65	202	10.8
> 65	74	3.9
Season		
Spring	461	24.5
Summer	508	27.1
Autumn	515	27.4
Winter	394	21.0
Time of accident		
0-5:59	37	2.0
6-11:59	513	27.3
12-16:59	573	30.5
17-23:59	755	40.2
Vehicle type		
Car	453	24.1
Heavy car	18	1.0
Motorcycle	1407	74.9

Collision types are presented in table 3. As data in table 3 shows, the accidents with the highest frequency were those of a motorcycle with a car (n = 913, 48.6%). It was also observed that in the majority of accidents, one of the vehicles involved was a motorcycle (n =

1076, 56.2%). Frequency distribution of type of vehicle involved in the accidents based on the severity of injury is presented in table 4. There were 4 deceased cases among those injured in MVAs.

Table 2: Association between age and type of vehicle

Type of accident		Age group					Total
		0-14 year	15-29 year	30-45 year	46-65 year	> 65 year	
Car	N	9	52	47	17	6	131
	%	6.0	4.9	12.2	8.4	8.1	7.0
Motorcycle	N	39	617	167	71	19	913
	%	25.8	57.9	43.3	35.1	25.7	48.6
Pedestrians	N	65	85	61	75	35	321
	%	43.0	8.0	15.8	37.1	47.3	17.1
Rollover	N	30	248	78	30	10	396
	%	19.9	23.3	20.2	14.9	13.5	21.1
Other	N	8	63	33	9	4	117
	%	5.3	5.9	8.5	4.5	5.4	6.2
Total	N	151	1065	386	202	74	1878
	%	100	100	100	100	100	100

Table 3: Frequency distribution of accidents based on the type of collision

Type of collision	Frequency	Percentage	Valid percent	Cumulative percentage
Car-Pedestrian	209	11.1	11.1	11.1
Car-Motorcycle	913	48.6	48.6	59.7
Heavy car-Motorcycle	14	0.7	0.7	60.5
Car-Barrier	52	2.8	2.8	63.3
Motorcycle-Barrier	17	0.9	0.9	64.2
Car-car	131	7.0	7.0	71.1
Heavy car-Car	21	1.1	1.1	72.3
Bicycle-Heavy car	12	0.6	0.6	72.9
Bicycle-Barrier	1	0.1	0.1	72.9
Motorcycle-Pedestrian	112	6.0	6.0	78.9
Overturn	396	21.1	21.1	100
Total	1878	100	100	

Moreover, 50% of deceased cases (n = 2) were injured by personal cars and the remaining were injured by motorcycles (n = 2). Of the 1386 victims of accidents due to motorcycles, about 30% were severely injured (n = 410). More than 90% of accidents by bicycles

resulted in mild injuries and less than 10% of individuals riding bicycles were severely injured in the accidents. No significant difference was observed between the severities of accidents by different vehicles (Table 4).

Table 4: Frequency distribution of type of vehicle based on the severity of injuries

Severity		Type					Total
		Car	Heavy car	Motorcycle	Bicycle	Tractor	
Minor	N	298	9	974	19	3	1303
	%	65.5	60.0	70.3	90.5	100	69.4
Severe	N	153	6	410	2	0	571
	%	33.8	40.0	29.6	9.5	0.0	30.4
Death	N	2	0	2	0	0	4
	%	0.4	0.0	0.1	0.0	0.0	0.2
Total	N	453	15	1386	21	3	1878
	%	100	100	100	100	100	100

Discussion

Disabilities and death related to injuries are the most tragic phenomena in the present day. In other words, injuries (intentional and/or unintentional) are among the most common public health problems. As a result, public health has an important role to play in

continuing the research for a better understanding of injuries (3, 6). In this study, we found that men were injured more than women, as injury in men was more than six times that in women. The majority of the victims (56.7%) were in the age group of 15-29 years. Many previous epidemiological

studies declared that the majority of victims were young and middle aged; moreover, there was a higher rate of injuries in men than women (11, 13, 14). The study by Dura et al. showed that the frequency of injuries in women and men were 32% and 68%, respectively. Furthermore, the mean age of victims was 34.8 ± 17.3 years, with an age range of 3 months to 87 years (15).

Another study in Mexico on victims of traffic accidents, reported that 54% of injured individuals had been injured in MVSs (16). In addition, 59.3% of those injured in MVSs were under 29 years of age, and 70.39% were men and 29.61% were women (16). Results obtained in the present study, with respect to sex and age, are in agreement with those reported in other studies (17, 18). There was no difference between the different seasons, although the number of accidents which had occurred in summer and autumn was higher than winter and spring. Also results show that the majority of accidents occur in the 17 hours before midnight. The study by Charkazi et al. found that most accidents occur in the summer and autumn (19), which was in agreement with the results of the present study.

Results presented in table 2 show that most victims were in the age group of 15-29 years and had had an accident with a motorcycle. However, in the age groups of younger than 15 years and older than 65 years, the majority of victims had been a pedestrian at the time of the accident. This indicates the need for education for the age group of 15-29 years in relation to the safe use of motorcycles. The second reason for accidents in the age group of 15-29 years was rollover, indicating the high speed of the vehicle and their inability to control the car. This age group in turn, is deserved to be trained for driving, according to the training instructions issued by government. Charkazi et al. found that the age group of 13-29 years, with 63.9% traffic accidents, had the highest rate of victims (19). Nóbrega et al. analyzed 2570 patients who had had traffic accidents and found that 67.8% were motorcyclists (13).

This result is in accordance with the findings of the current study.

As table 3 shows, the highest percentage of victims were involved in a car collision with a motorcycle. The results showed that motorcyclists in the city are a major factor in accidents; therefore, the use of appropriate interventions for these individuals is essential to reduce injuries. An important intervention is the training of traffic laws and regulations. Car collision with a pedestrian is important and represents pedestrians' use of unsafe passages. Our findings show that the most recorded accidents (75%) had been occurred by motorcycles that illustrates motorcycles are being used frequently in the area of this study. Our study revealed that MVAs constituted the majority of injuries in this district of Kerman Province. Among the MVAs, motorcycle accidents were the most common causes of injuries. Thus, in fact, motorcyclists can be classified as a vulnerable group of the population in this region of Iran.

Palaver et al also found that 67.7% of collisions were with motorcycles, and the most frequent type of collision by vehicles was motorcycle-car on city streets with 42% (20). Spoerri et al. found that only 15.7% of MVAs were motorcycle accidents. Moreover, 16.0% of victims were pedestrians (21). Hayakawa et al. showed that the percentage of fatal vehicle collisions with pedestrians were higher in Japan than in the United States (28.5% vs. 18.2%) (22). According to a study conducted by Dilek Durak et al., trauma was most commonly sustained within the motor vehicle (15).

To reduce this type of accident, it is recommended that pedestrian use overpasses or underpasses. Rollovers are a major cause of traffic accidents and indicate the high speed of drivers. Training regulations and traffic laws to reduce these types of accidents can be a great help.

As shown in table 4, although the number of accidents by heavy cars is small, they result in a high percentage of severe injuries, which is an indication for the necessity of action in this

regard. More than a third of accidents with cars resulted in severe injuries. Hence, it is necessary that appropriate action be taken to enforce traffic laws and guidelines. The research by Hajar et al. shows that 69.96% of victims had non-severe injuries and only 30.96% of victims had severe injuries (16). This finding is in agreement with that of the current research.

Conclusion

The results of this study could help managers and social planners in the health section, especially police department planners to modify and control the generating factors of MVAs in order to reduce MVAs. MVAs, as important causes of injuries, most frequently occurred by motorcycles in the district studied. Therefore, safety training targeted at motorcyclists would be one of the best interventions for the prevention and controlling of trauma due to MVAs. This intervention could be the most efficient among younger drivers.

Acknowledgments

The authors would like to thank and fully acknowledge the participants, in particular our colleagues in Pre-hospital Emergency Ward, for making this study possible. The authors would also like thank Occupational Environment Research Center for their financial support.

Conflict of interest: None declared.

References

- 1- Moshiro C, Heuch I, Astrøm AN, Setel P, Hemed Y, Kvåle G. Injury morbidity in an urban and a rural area in Tanzania: an epidemiological survey. *BMC Public Health* 2005; 5:11.
- 2- Olawale OA, Owoaje ET. Incidence and pattern of injuries among residents of a rural area in South-Western Nigeria: a community-based study. *BMC Public Health* 2007; 7:246.
- 3- Mock C, Quansah R, Krishnan R, Arreola-Risa C, Rivara F. Strengthening the prevention and care of injuries worldwide. *Lancet* 2004; 363(9427):2172-9.
- 4- Wesson HK, Boikhutso N, Bachani AM, Hofman KJ, Hyder AA. The cost of injury and trauma care in low- and middle-income countries: a review of economic evidence. *Health Policy Plan* 2014; 29(6):795-808.
- 5- Zwi AB, Forjuoh S, Murugusampillay S, Odero W, Watts C. Injuries in developing countries: policy response needed now. *Trans R Soc Trop Med Hyg* 1996; 90(6):593-5.
- 6- Norton R, Kobusingye O. Injuries. *N Engl J Med* 2013; 368(18):1723-30.
- 7- Ameratunga S, Hajar M, Norton R. Road-traffic injuries: confronting disparities to address a global-health problem. *Lancet* 2006; 367(9521):1533-40.
- 8- Sharma BR. Road traffic injuries: a major global public health crisis. *Public Health* 2008; 122(12):1399-406.
- 9- Reddy GM, Negandhi H, Singh D, Singh AJ. Extent and determinants of cost of road traffic injuries in an Indian city. *Indian J Med Sci* 2009; 63(12):549-56.
- 10- Mohan D. Road traffic injuries: a stocktaking. *Best Pract Res Clin Rheumatol* 2008; 22(4):725-39.
- 11- Afzali S, Saleh A, Seif Rabiei MA, Taheri K. Frequency of alcohol and substance abuse observed in drivers killed in traffic accidents in Hamadan, Iran. *Arch Iran Med* 2013; 16(4):240-2.
- 12- Majdzadeh R, Khalagi K, Naraghi K, Motevalian A, Eshraghian MR. Determinants of traffic injuries in drivers and motorcyclists involved in an accident. *Accid Anal Prev* 2008; 40(1):17-23.
- 13- Nóbrega LM, Cavalcante GM, Lima MM, Madruga RC, Ramos-Jorge ML, d'Avila S. Prevalence of facial trauma and associated factors in victims of road traffic accidents. *Am J Emerg Med* 2014; 32(11):1382-6.
- 14- Kanaan A, Huertas P, Santiago A, Sánchez JA, Martínez P. Incidence of different health factors and their influence on traffic accidents in the province of Madrid, Spain. *Leg Med (Tokyo)* 2009; 11 Suppl 1:S333-6.
- 15- Durak D, Fedakar R, Türkmen N, Akgöz S, Baduroğlu E. Road traffic collisions in Bursa, Turkey, during 2003, 2004 and 2005. *Injury* 2008; 39(5):547-53.
- 16- Hajar M, Arredondo A, Carrillo C, Solórzano L. Road traffic injuries in an urban area in Mexico. An epidemiological and cost analysis. *Accid Anal Prev* 2004; 36(1):37-42.
- 17- Batstone MD, Monsour FN, Pattel P, Lynham A. The patterns of facial injury suffered by patients in road traffic accidents: a case controlled study. *Int J Surg* 2007; 5(4):250-4.
- 18- Vorko-Jović A, Kern J, Biloglav Z. Risk factors in urban road traffic accidents. *J Safety Res* 2006; 37(1):93-8.
- 19- Charkazi A, Esmaeili A, Garkaz G, Qoreishi Z, Gerey S, Nazari S. Epidemiologic survey

- of road traffic accidents in patients admitted in emergency department of Alejalil Hospital in Aq-Qala city, Golestan province. *Journal of Health Ardabil University of Medical Sciences* 2012; 3(2):42-9.
- 20- Paravar M, Hosseinpour M, Salehi S, Mohammadzadeh M, Shojaee A, Akbari H, et al. Pre-Hospital Trauma Care in Road Traffic Accidents in Kashan, Iran. *Arch Trauma Res* 2013; 1(4):166-71.
- 21- Spoerri A, Egger M, von Elm E; Swiss National Cohort Study. Mortality from road traffic accidents in Switzerland: longitudinal and spatial analyses. *Accid Anal Prev* 2011; 43(1):40-8.
- 22- Hayakawa H, Fischbeck PS, Fischhoff B. Traffic accident statistics and risk perceptions in Japan and the United States. *Accid Anal Prev* 2000; 32(6):827-35.

Archive of SID