



# Antioxidant, Antiglycation and Anti-Hyperlipidemic Effects of *Trigonella foenum* and Cinnamon in Type 2 Diabetic Rats

Seyed Mehrdad Kassaee,<sup>1\*</sup> Mohammad Taghi Goodarzi,<sup>2,3</sup> and Ebrahim Abbasi Oshaghi<sup>2</sup>

<sup>1</sup>Department of Biology, Hamedan Branch, Islamic Azad University, Hamedan, Iran

<sup>2</sup>Department of Clinical Biochemistry, Medical School, Hamadan University of Medical Sciences, Hamadan, Iran

<sup>3</sup>Research Center for Molecular Medicine, Hamadan University of Medical Sciences, Hamadan, Iran

\*Corresponding author: Seyed Mehrdad Kassaee, Department of Biology, Hamedan Branch, Islamic Azad University, Hamedan, Iran. Fax: +98-8138238997, E-mail: kassaee2001@yahoo.com

Received 2016 April 12; Revised 2016 August 18; Accepted 2016 September 08.

## Abstract

**Background:** Diabetes is recognized as a common metabolic disorder, which is treated by different medicines.

**Objectives:** The aim of this study was to study the effect of *Trigonella foenum* and Cinnamon on glycation, some biochemical factors, liver histology and cholesterol 7-alpha hydroxylase activity in type 2 diabetic rats.

**Methods:** Antiglycation and antioxidant ability were determined in vitro. Male Wistar diabetic rats were treated by 2 and 8% of Cinnamon (w/w) and *Trigonella foenum*. Biochemical factors and liver enzymes were measured using enzymatic spectrophotometric methods. Lipase and cholesterol 7-alpha hydroxylase activity were determined. Liver lipid, antioxidant and morphological change were assessed.

**Results:** Different concentrations of these plants (0.032, 0.065, 0.125, 0.25, 0.5 and 1mg/ mL) showed potential antiglycation ability. Cinnamon and *Trigonella foenum* significantly reduced advanced glycosylation end products (AGEs) and fructosamine formation, and also declined protein carbonyl contents and thiol group's oxidation ( $P < 0.001$ ). Cinnamon and *Trigonella foenum* extract in Streptozotocin (STZ)-induced diabetic rat normalized antioxidant capacity and lipid profile ( $P < 0.001$ ). The cholesterol 7-alpha hydroxylase activity significantly increased following these plants administration. Also, the liver histopathological changes were normalized in diabetic rats treated by Cinnamon and *Trigonella foenum* extract.

**Conclusions:** The findings illustrated that Cinnamon and *Trigonella foenum* extract improved hyperlipidemia and hyperglycemic. The hypolipidemic effect was likely by motivation of cholesterol 7-alpha hydroxylase activity. These plants also showed significant anti-diabetic effects in animal model by reducing blood glucose and inhibition of AGEs and fructosamine formation.

**Keywords:** Cinnamon, Cholesterol 7-alpha Hydroxylase, Herbal Medicine, *Trigonella Foenum*, Type 2 Diabetes

## 1. Background

Diabetes is recognized as a common metabolic disorder that is described by high plasma glucose levels, as a result of insulin resistance or lacking of insulin secretion from pancreatic beta cells (1). This disease affected about 382 million people worldwide in the 2013 and its incidence rate is estimated to reach 592 million people by 2035 (2). Many clinical trials and epidemiological experiments support the idea that high blood glucose levels is the major cause of short-term diabetic complications such as diabetic ketoacidosis and hyperosmolar hyperglycaemic state (HHS) and also long-term complications such as nephropathy, cardiovascular disease and retinopathy (1). On the other hand, free radicals are involved in several diseases such as cancer, atherosclerosis and diabetes. Components, which are able to scavenge free radicals, play a potential role in improving these diseases (3, 4). Antioxidants play

a vital role in the protection of body cells from damage induced by free radicals (5). Increase in oxidative stress that is motivated by free radicals has been established in patients with diabetes. Rise of blood glucose and free radicals can start lipid peroxidation that stimulates protein glycation, enzymes inactivation and metabolism alteration, and also plays a major role in the long-term complications of diabetes (1). This non-enzymatic glycation of blood protein leads to formation of advanced glycosylation end products (AGEs), which cause different complications in diabetes (1, 4).

It has also been identified that induced diabetes by Streptozotocin in animal models mainly provokes production of oxygen free radicals and thus destructs pancreatic cells (5). Supplementation with natural antioxidants can be one of the therapeutic approaches for declining hyperglycemia and free radicals in diabetic patients (6, 7). Dur-

ing the past few years, great attention has been paid to the use herbal medicine for the treatment of diabetes (4). Ethno-botanical evidence shows that about 800 plants are applied as traditional remedies for diabetes complications treatment (4).

The hypoglycemic properties of numerous herbal medicines has been assessed and established in humans and animal models. Different extracts of some plants and plant derivatives and their components have been shown to contain significant antioxidant activity and hypoglycemic properties. In this respect, cinnamon and *Trigonella foenum* are traditional herbal medicines, which display hypolipidemic, hypoglycemic and antioxidant properties (6, 8-10). Although Cinnamon and *Trigonella foenum* showed hypolipidemic and hypoglycemic activity, possible mechanism of their action in diabetic animals, especially for the later herbal is unknown.

## 2. Objectives

The current study was planned to investigate the effect of *Trigonella foenum* and Cinnamon on glycation, some biochemical factors, liver histology and lipase and cholesterol 7-alpha hydroxylase activity in type 2 diabetic rats.

## 3. Methods

### 3.1. Preparation of Plant Extracts

*Trigonella foenum* and Cinnamon powder were prepared and then crushed and dried. Next, 20 g of dried powder was mixed with 200 mL of deionized water at room temperature for 48 hours. The residue was filtered, and the filtrate solution was concentrated at 40°C in an incubator for 24 hours. The extract of *Trigonella foenum* and Cinnamon was kept in dark vials at -20°C until analysis.

### 3.2. Glycation of Bovine Serum Albumin and Advanced Glycation End Products (AGE) Assay

Bovine Serum Albumin (BSA) was glycated via treatment with fructose and produced AGEs was measured using spectrophotometric assay according to the method described by Abbasi Oshaghiet al. (11). Briefly, BSA (10 mg/mL) was treated separately with two concentrations of fructose (500 and 200 mM) containing 0.02% sodium azide (in 0.1 M phosphate buffer) and in the absence and precedence of different concentrations of *Trigonella foenum* or Cinnamon (0.25 - 2 mg/ mL). Also, Aminoguanidine (AG) was used as a positive control in this experiment (12). The tubes were incubated at 37°C at dark room for two, three, and four weeks. After that, solutions of every tube were dialyzed against

phosphate buffered saline (PBS) (at 4°C) for 48 hours. Following dialysis, the protein concentration was measured using the Bradford assay. Fluorescence intensity was determined at 335 nm excitation and 460 nm emission (13) using a spectrofluorometer (Jasco FP-6200).

### 3.3. Determination of Fructosamine

The fructosamine formation was determined according to previous published method (13) using nitroblue tetrazolium (NBT). Briefly, 90  $\mu$ L of NBT (0.5 mM) in 0.1 M carbonate buffer (pH 10.4 at 37°C) was mixed with 10  $\mu$ L of a prepared glycated sample (the above step). Afterward, the absorbance of samples was determined at 10- and 15-minute time points at 530 nm using an enzyme linked immunosorbent assay (ELISA) reader (ELX 800, Bio-Tek Inst.).

### 3.4. Determination of Protein Carbonyl Content and Thiol Group

Carbonyl contents of glycated BSA, a marker for protein oxidative damage, were determined, according to a previously published method (13) using 2, 4-Dinitrophenylhydrazine (DNPH). The carbonyl content of each sample was calculated based on the extinction coefficient of DNPH ( $\epsilon = 22,000 \text{ M}^{-1}\text{cm}^{-1}$ ). The results are presented as nmole carbonyl/mg protein. The thiol group was determined based on Ellman's method using DTNB with slight modifications (13). The absorbance of samples was determined at 410 nm using a spectrophotometer (JENWAY 6105 UV/Vis). Finally, the free thiol concentration of samples was determined using L-cysteine (0.015-0.50 mM) as a standard and expressed as nmol/mg protein (13).

### 3.5. Experimental Animals and Design

Male Wistar rats weighting 200 - 220 grams were used in this study. All animals were kept at the animal house of Hamadan University of Medical Sciences (Iran). The rats were kept in standard cages under normal condition of 12/12 hour (light/dark period) with relative humidity of  $56 \pm 3\%$ . After one-week adaptation in an animal house, they were divided randomly to six groups of six rats in each group. For induction of type 2 diabetes, 65 mg/kg Streptozotocin (STZ) was injected to rats after 12 hours of fasting. Nicotinamide at a dose of 110 mg/kg was injected intraperitoneally 15 minutes following STZ injection (11). One week later, glucose levels of animals were determined enzymatically and fasting blood glucose level more than 250 mg/dl was considered diabetic (14). The studied groups were as follow: Group 1 contained normal rats that received standard chow diet (control group). Group 2 was diabetic rats that received standard chow diet. Groups 3 and 4 were diabetic rats that in addition to standard diet received 2% and

8% Cinnamon extract (w/w), respectively. Groups 5 and 6 received 2% and 8% of *Trigonella foenum* extract (w/w). The regimen was continued for 30 days. All processes of the experiments were approved by the animal research ethic committee of Hamadan Azad University (Hamadan, Iran).

### 3.6. Blood and Liver Biochemical Parameters

On day 30, blood samples were collected by heart puncture from each animal. The serum was separated by centrifugation for 10 minutes at  $1500\times g$ . The prepared serum was subsequently used for enzymatic measurement of blood glucose, total cholesterol, high density lipoprotein-cholesterol (HDL-C), triglyceride, alanine aminotransferase (ALT) and aspartate aminotransferase (AST), using commercial kits (Pars Azmun Co. Iran) (15). Very low density lipoprotein cholesterol (VLDL-C) and low density lipoprotein cholesterol (LDL-C) were calculated according to Friedewald's equation (16).

### 3.7. Liver Lipids and Glycogen

Liver lipids were determined according to Butler et al. (17). Briefly, one gram of liver tissue was homogenized in chloroform/methanol. After centrifugation, the solvent was washed using 0.9% NaCl. The solution was centrifuged and chloroform phase was used for cholesterol and triglyceride assays with the same enzymatic kit as applied for serum analysis. Liver glycogen was determined according to Folch et al. (18) methods. Briefly, 500 mg of liver and equal volume of 10% trichloroacetic acid was added and was homogenized after 30 minutes. The resulting solution was centrifuged. The supernatant was obtained and 95% ethanol was added to the samples. After that, sodium chloride was added and resulting precipitate was dissolved in distilled water. The precipitate was dissolved in ethanol and centrifuged again. The supernatant was discarded and the resulting precipitate was dissolved in absolute ethanol and the solution was dispensed in a glass plate. After evaporation of ethanol, the remaining sediment was collected and weighed. Liver glycogen content of liver was reported as mg to gram tissue.

### 3.8. Blood Glucose and Glycosylation End Products Assay

Fasting blood glucose (FBS) was measured enzymatically using a commercial kit (Pars Azmun Co.). For determination of AGEs in the kidney and serum, the samples were diluted 1:50 with PBS. Fluorescence intensity was determined at 335 nm excitation and 460 nm emission using a spectrofluorometer (Jasco FP-6200). Fluorescence intensity was expressed as arbitrary units (AU) (13).

### 3.9. Determination of Oxidative Stress Markers

Malondialdehyde (MDA) level, a marker of lipid peroxidation was measured using the thiobarbituric acid reaction. The results were stated as nmol of MDA/mL of serum. Total antioxidant capacity (TAC) was measured according to a previous published method (19). Thiol group of serum was determined based on Ellman's method using DTNB with slight modifications according to a previous report (13).

### 3.10. Enzymes Assay

Cholesterol 7-alpha hydroxylase level and lipase activity were determined using commercial kits (Eastbio-pharma Co. and Pars Azmun respectively) and according to the manufacturer's instructions.

### 3.11. Histopathological Examination

The livers of animals were fixed in 10% formalin solution and processed by standard methods. Briefly, the samples were embedded in paraffin; sections at  $5\ \mu\text{m}$  was prepared and stained with haematoxylin and eosin (H&E). The stained slides were evaluated under a light microscope.

### 3.12. Statistical Analysis

The data in this study are presented as means  $\pm$  standard error of the mean (SEM). A three-replicate measurement was done in in vitro tests. Statistical analyses were carried out using the SPSS statistical program (SPSS; 16, Inc, USA). One-way analysis of variances (ANOVA) followed by Tukey-test was used for statistical analysis of data. P values of less than 0.05 were regarded as statically significant.

## 4. Results

### 4.1. In Vitro Experiments

*Trigonella foenum* and Cinnamon markedly declined AGEs formation in treated BSA with fructose (Table 1). Cinnamon showed higher effect in this experiment. Interestingly, inhibition of AGEs by *Trigonella foenum* or Cinnamon at a concentration of 1 and 2 mg/mL was more than that of aminoguanidine ( $P < 0.05$ ). Our experiment also established that *Trigonella foenum* and Cinnamon significantly inhibited fructosamine formation ( $P < 0.05$ ). Major rise in the carbonyl content and also thiol groups oxidation were observed when the BSA solution was incubated in presence of 200 and 500 mM fructose with *Trigonella foenum* and Cinnamon (Tables 1-2) ( $P < 0.05$ ).

**Table 1.** Inhibition of Advanced Glycosylation End Products Formation (arbitrary unit) and Fructosamine Levels (mmol/ mg protein) by Cinnamon and *Trigonella foenum* at Three Different Treatment Times<sup>a</sup>

Experimental Groups	AGE levels (arbitrary unit)			Fructosamine levels (mmol/ mg protein)		
	Week 2	Week 3	Week 4	Week 2	Week 3	Week 4
<b>BSA/Fru 500 mM</b>	113.20 ± 6.66	127.24 ± 7.97	137.57 ± 10.15	3.22 ± 0.19	3.27 ± 0.24	3.52 ± 0.37
<b>+Cinnamon 0.25 mg/mL + TF 0.25 mg/mL</b>	50.00 ± 6.21 <sup>b</sup>	54.64 ± 5.23 <sup>b</sup>	68.23 ± 7.13 <sup>b</sup>	2.30 ± 0.11 <sup>b</sup>	2.74 ± 0.03 <sup>b</sup>	3.10 ± 0.04 <sup>b</sup>
	67.12 ± 6.02 <sup>b</sup>	68.89 ± 6.77 <sup>b</sup>	70.11 ± 6.24 <sup>b</sup>	2.55 ± 0.09 <sup>b</sup>	2.88 ± 0.09 <sup>b</sup>	3.02 ± 0.09 <sup>b</sup>
<b>+ Cinnamon 0.5 mg/mL + TF 0.5 mg/mL</b>	48.11 ± 5.55 <sup>b</sup>	51.41 ± 6.01 <sup>b</sup>	67.76 ± 6.09 <sup>b</sup>	2.32 ± 0.09 <sup>b</sup>	2.71 ± 0.08 <sup>b</sup>	3.08 ± 0.06 <sup>b</sup>
	62.17 ± 4.20 <sup>b</sup>	62.17 ± 4.93 <sup>b</sup>	66.38 ± 6.96 <sup>b</sup>	2.46 ± 0.11 <sup>b</sup>	2.80 ± 0.03 <sup>b</sup>	3.00 ± 0.08 <sup>b</sup>
<b>+ Cinnamon 1 mg/mL + TF 1 mg/mL</b>	40.46 ± 6.31 <sup>b</sup>	43.12 ± 5.10 <sup>b</sup>	58.32 ± 4.91 <sup>b</sup>	2.23 ± 0.14 <sup>b</sup>	2.59 ± 0.05 <sup>b</sup>	2.73 ± 0.13 <sup>b</sup>
	66.30 ± 6.87 <sup>b</sup>	60.55 ± 5.19 <sup>b</sup>	60.88 ± 5.55 <sup>b</sup>	2.37 ± 0.12 <sup>b</sup>	2.76 ± 0.03 <sup>b</sup>	2.88 ± 0.15 <sup>b</sup>
<b>+ Cinnamon 2 mg/mL + TF 2 mg/mL</b>	39.89 ± 5.15 <sup>b</sup>	41.31 ± 4.22 <sup>b</sup>	51.17 ± 4.31 <sup>b</sup>	2.20 ± 0.10 <sup>b</sup>	2.07 ± 0.07 <sup>b</sup>	2.25 ± 0.07 <sup>b</sup>
	57.74 ± 6.91 <sup>b</sup>	54.12 ± 4.07 <sup>b</sup>	56.43 ± 5.19 <sup>b</sup>	2.37 ± 0.20 <sup>b</sup>	2.19 ± 0.08 <sup>b</sup>	2.37 ± 0.06 <sup>b</sup>
<b>+ AG 2 mg/mL</b>	45.35 ± 4.11 <sup>b</sup>	47.15 ± 4.49 <sup>b</sup>	50.20 ± 5.13 <sup>b</sup>	2.40 ± 0.08 <sup>b</sup>	2.09 ± 0.14 <sup>b</sup>	2.40 ± 0.14 <sup>b</sup>
<b>BSA/Fru 200 mM</b>	108.30 ± 7.30	133.54 ± 8.18	147.24 ± 5.26	2.71 ± 0.11	2.84 ± 0.15	2.99 ± 0.08
<b>+Cinnamon 0.25 mg/mL + TF 0.25 mg/mL</b>	48.78 ± 5.471 <sup>b</sup>	51.09 ± 6.12 <sup>b</sup>	66.38 ± 6.13 <sup>b</sup>	2.62 ± 0.07 <sup>b</sup>	2.62 ± 0.07 <sup>b</sup>	2.76 ± 0.06 <sup>b</sup>
	65.30 ± 6.40 <sup>b</sup>	66.14 ± 5.11 <sup>b</sup>	67.47 ± 5.33 <sup>b</sup>	2.50 ± 0.09 <sup>b</sup>	2.70 ± 0.12 <sup>b</sup>	2.88 ± 0.01 <sup>b</sup>
<b>+ Cinnamon 0.5 mg/mL + TF 0.5 mg/mL</b>	44.48 ± 5.62 <sup>b</sup>	47.38 ± 5.30 <sup>b</sup>	61.40 ± 5.40 <sup>b</sup>	2.28 ± 0.06 <sup>b</sup>	2.56 ± 0.05 <sup>b</sup>	2.60 ± 0.07 <sup>b</sup>
	65.16 ± 5.22 <sup>b</sup>	66.07 ± 6.20 <sup>b</sup>	63.36 ± 6.28 <sup>b</sup>	2.38 ± 0.09 <sup>b</sup>	2.61 ± 0.06 <sup>b</sup>	2.74 ± 0.08 <sup>b</sup>
<b>+ Cinnamon 1 mg/mL + TF 1 mg/mL</b>	45.16 ± 4.86 <sup>b</sup>	45.81 ± 5.27 <sup>b</sup>	55.54 ± 5.04 <sup>b</sup>	2.16 ± 0.18 <sup>b</sup>	2.48 ± 0.10 <sup>b</sup>	2.51 ± 0.09 <sup>b</sup>
	58.31 ± 5.79 <sup>b</sup>	59.88 ± 5.7 <sup>b</sup>	60.45 ± 5.60 <sup>b</sup>	2.20 ± 0.18 <sup>b</sup>	2.55 ± 0.09 <sup>b</sup>	2.67 ± 0.02 <sup>b</sup>
<b>+ Cinnamon 2 mg/mL + TF 2 mg/mL</b>	40.11 ± 3.10 <sup>b</sup>	44.27 ± 5.10 <sup>b</sup>	48.92 ± 4.71 <sup>b</sup>	2.09 ± 0.08 <sup>b</sup>	2.03 ± 0.05 <sup>b</sup>	2.18 ± 0.07 <sup>b</sup>
	53.81 ± 5.22 <sup>b</sup>	58.78 ± 6.12 <sup>b</sup>	55.33 ± 5.59 <sup>b</sup>	2.12 ± 0.14 <sup>b</sup>	2.30 ± 0.07 <sup>b</sup>	2.22 ± 0.09 <sup>b</sup>
<b>+ AG 2 mg/mL</b>	50.31 ± 3.26 <sup>b</sup>	49.75 ± 5.25 <sup>b</sup>	52.31 ± 5.98 <sup>b</sup>	2.08 ± 0.08 <sup>b</sup>	2.11 ± 0.05 <sup>b</sup>	2.28 ± 0.04 <sup>b</sup>
<b>BSA/PBS</b>	18.10 ± 1.99 <sup>b</sup>	20.17 ± 5.73 <sup>b</sup>	23.66 ± 3.10 <sup>b</sup>	0.35 ± 0.04 <sup>b</sup>	0.19 ± 0.02 <sup>b</sup>	0.26 ± 0.03 <sup>b</sup>

Abbreviations: AG, Aminoguanidine; Alb/Fru, albumin + fructose; BSA, bovine serum albumin; TF, *Trigonella foenum*.

<sup>a</sup>Data represent as mean ± SEM (n = 3).

<sup>b</sup>P < 0.001 compared with BSA/fructose at the same incubation week.

#### 4.2. In Vivo Experiments

Table 3 shows the mean of fasting blood glucose (FBS), triglyceride, cholesterol, LDL-C, HDL-C and VLDL-C levels in different studied groups. The levels of total cholesterol, triglyceride, LDL-C and VLDL-C were significantly reduced (P < 0.05) in *Trigonella foenum* and Cinnamon extract treated animals. *Trigonella foenum* or Cinnamon also significantly normalized liver enzymes (ALT and AST) in treated diabetic animals (Table 3).

The total antioxidant level and serum thiol groups were significantly (P < 0.05) lower in diabetic animals compared with normal rats (Figure 1). In *Trigonella foenum* and Cinnamon-treated diabetic rats total antioxidant level was significantly restored (P < 0.05) (Figure 1). On the other hand, MDA levels were significantly reduced in *Trigonella foenum* and Cinnamon treated rats (P < 0.05, Figure 2).

Advanced Glycosylation End Products formation in the serum of different groups is shown in Figure 3. The change of AGEs formation in diabetic animals, which were treated

with *Trigonella foenum* or Cinnamon was not significant compared with diabetic and control rats. Lipase activity was not changed in all treated groups except in the group that was treated with Cinnamon at a dose of 8 g/100g diet. Cholesterol 7-alpha hydroxylase activity was significantly reduced in *Trigonella foenum* and Cinnamon treated diabetic rats at a dose of 8 g/100g diet (Figure 4).

#### 4.3. Histological Changes

Figure 5 shows the morphological changes in liver of different studied groups. Liver histology in control group showed normal structure including normal cellular construction with portal triad, numerous hepatocytes (a), lining sinusoidal (c) and central vein (b). As shown in Figure 5 accumulation of lipid (d) and foam cell (e) and infiltrations of inflammatory cells (f) in liver were obviously observed in diabetic rats and treated diabetic rats with dose of 2 g/100g diet of *Trigonella foenum* or Cinnamon; while, lipid accumulation was normalized by the dose of 8 g/100g

**Table 2.** The Effect of Cinnamon and *Trigonella foenum* Extracts on Thiol Group (nmol /mg protein) and Carbonyl Content (nmol/mg protein) at Three Different Treatment Times<sup>a</sup>

Experimental Groups	Thiol Group (nmol /mg Protein)			Carbonyl Content (nmol/mg Protein)		
	Week 2	Week 3	Week 4	Week 2	Week 3	Week 4
BSA/Fru 500 mM	1.90 ± 0.08	1.59 ± 0.07	1.20 ± 0.07	2.8 ± 0.06	3.7 ± 0.05	3.9 ± 0.02
+Cinnamon 0.25 mg/mL + TF 0.25 mg/mL	2.21 ± 0.11 <sup>b</sup>	2.35 ± 0.03 <sup>b</sup>	1.90 ± 0.05 <sup>b</sup>	2.18 ± 0.08 <sup>b</sup>	2.51 ± 0.1 <sup>b</sup>	2.99 ± 0.08 <sup>b</sup>
	2.14 ± 0.11 <sup>b</sup>	2.23 ± 0.03 <sup>b</sup>	1.51 ± 0.07 <sup>b</sup>	2.8 ± 0.08	3.31 ± 0.17 <sup>b</sup>	3.49 ± 0.08 <sup>b</sup>
+Cinnamon 0.5 mg/mL + TF 0.5 mg/mL	2.50 ± 0.07 <sup>b</sup>	2.39 ± 0.08 <sup>b</sup>	1.80 ± 0.08 <sup>b</sup>	2.01 ± 0.1 <sup>b</sup>	2.6 ± 0.09 <sup>b</sup>	2.9 ± 0.03 <sup>b</sup>
	2.39 ± 0.07 <sup>b</sup>	2.30 ± 0.07 <sup>b</sup>	1.71 ± 0.10 <sup>b</sup>	2.61 ± 0.11	2.96 ± 0.09 <sup>b</sup>	3.45 ± 0.13 <sup>b</sup>
+ Cinnamon 1 mg/mL + TF 1 mg/mL	2.49 ± 0.08 <sup>b</sup>	2.47 ± 0.05 <sup>b</sup>	1.88 ± 0.06 <sup>b</sup>	1.9 ± 0.1 <sup>b</sup>	2.4 ± 0.03 <sup>b</sup>	2.8 ± 0.08 <sup>b</sup>
	2.40 ± 0.08 <sup>b</sup>	2.38 ± 0.05 <sup>b</sup>	1.87 ± 0.08 <sup>b</sup>	2.2 ± 0.11 <sup>b</sup>	2.88 ± 0.03 <sup>b</sup>	3.1 ± 0.13 <sup>b</sup>
+Cinnamon 2 mg/mL + TF 2 mg/mL	2.66 ± 0.05 <sup>b</sup>	2.51 ± 0.08 <sup>b</sup>	2.14 ± 0.06 <sup>b</sup>	1.9 ± 0.06 <sup>b</sup>	2.2 ± 0.06 <sup>b</sup>	2.7 ± 0.08 <sup>b</sup>
	2.55 ± 0.05 <sup>b</sup>	2.40 ± 0.10 <sup>b</sup>	2.05 ± 0.11 <sup>b</sup>	2.19 ± 0.06 <sup>b</sup>	2.7 ± 0.06 <sup>b</sup>	2.10 ± 0.16 <sup>b</sup>
+AG 2 mg/mL	2.40 ± 0.08 <sup>b</sup>	2.27 ± 0.09 <sup>b</sup>	1.90 ± 0.04 <sup>b</sup>	2.1 ± 0.1 <sup>b</sup>	2.2 ± 0.09 <sup>b</sup>	2.35 ± 0.05 <sup>b</sup>
BSA/Fru200 mM	1.99 ± 0.07	1.64 ± 0.04	1.32 ± 0.05	2.64 ± 0.06	2.74 ± 0.08	2.38 ± 0.02
+Cinnamon 0.25 mg/mL + TF 0.25 mg/mL	1.91 ± 0.11	2.20 ± 0.02 <sup>b</sup>	1.93 ± 0.11 <sup>b</sup>	2.01 ± 0.06	2.03 ± 0.10 <sup>b</sup>	2.95 ± 0.04 <sup>b</sup>
	2.21 ± 0.11 <sup>b</sup>	2.11 ± 0.05 <sup>b</sup>	1.85 ± 0.08 <sup>b</sup>	2.58 ± 0.08	2.13 ± 0.15 <sup>b</sup>	2.90 ± 0.08 <sup>b</sup>
+Cinnamon 0.5 mg/mL + TF 0.5 mg/mL	2.34 ± 0.08 <sup>b</sup>	2.29 ± 0.09 <sup>b</sup>	2.03 ± 0.06	2.00 ± 0.09 <sup>b</sup>	2.14 ± 0.12 <sup>b</sup>	2.97 ± 0.05 <sup>b</sup>
	2.25 ± 0.07 <sup>b</sup>	2.20 ± 0.09 <sup>b</sup>	1.94 ± 0.06 <sup>b</sup>	2.45 ± 0.07	2.22 ± 0.09 <sup>b</sup>	1.91 ± 0.02 <sup>b</sup>
+Cinnamon 1 mg/mL + TF 1 mg/mL	2.59 ± 0.09 <sup>b</sup>	2.29 ± 0.03 <sup>b</sup>	2.15 ± 0.08 <sup>b</sup>	1.88 ± 0.05 <sup>b</sup>	2.10 ± 0.03 <sup>b</sup>	2.8 ± 0.04 <sup>b</sup>
	2.47 ± 0.10 <sup>b</sup>	2.30 ± 0.06 <sup>b</sup>	2.08 ± 0.09 <sup>b</sup>	2.05 ± 0.10 <sup>b</sup>	2.06 ± 0.09 <sup>b</sup>	2.71 ± 0.03 <sup>b</sup>
+Cinnamon 2 mg/mL+ TF 2 mg/mL	2.61 ± 0.08 <sup>b</sup>	2.38 ± 0.05 <sup>b</sup>	2.20 ± 0.05 <sup>b</sup>	1.80 ± 0.07 <sup>b</sup>	1.85 ± 0.09 <sup>b</sup>	2.56 ± 0.13 <sup>b</sup>
	2.56 ± 0.06 <sup>b</sup>	2.41 ± 0.05 <sup>b</sup>	2.19 ± 0.04 <sup>b</sup>	1.91 ± 0.06 <sup>b</sup>	2.05 ± 0.06 <sup>b</sup>	2.65 ± 0.08 <sup>b</sup>
+AG 2 mg/mL	2.65 ± 0.09 <sup>b</sup>	2.32 ± 0.06 <sup>b</sup>	2.06 ± 0.04 <sup>b</sup>	1.56 ± 0.12 <sup>b</sup>	1.49 ± 0.09 <sup>b</sup>	2.38 ± 0.05 <sup>b</sup>

Abbreviations: AG, aminoguanidine; TF, *Trigonella foenum*; Alb/Fru, albumin + fructose; BSA, bovine serum albumin.

<sup>a</sup>Data represent as mean ± SEM (n = 3).

<sup>b</sup>p < 0.001 compared with BSA/fructose at the same incubation week.

diet. Liver of diabetic rats also exhibited cellular abnormalities with area of necrosis, mild fibrosis, cellular degeneration, vascular degeneration and congestion, and also infiltrations of inflammatory cells when compared with control rats. Treatments of diabetic animals with *Trigonella foenum* or Cinnamon at the dose of 8 g/100g diet normalized these changes. These changes were slightly restored with *Trigonella foenum* or Cinnamon at a dose of 2 g/100g diet (Figure 5).

### 5. Discussion

Recent meta-analysis studies indicated the useful properties of natural products in the management of diabetes

and related complications (20). This study showed the effects of Cinnamon on triglyceride, total cholesterol, LDL-C and blood glucose reduction in an animal model of type 2 diabetes.

Vanschoonbeek et al. (6) reported that administration of Cinnamon at a dose of 1.5 g/d did not normalize lipid profile, insulin sensitivity and oral glucose tolerance in postmenopausal patients of type 2 diabetes. Whereas, Khan et al. (7) with administration of 1, 3, or 6 g of Cinnamon per day in 60 people with type 2 diabetes showed that blood glucose levels, total cholesterol, triglyceride and LDL-C were significantly reduced. Differences between the studies of Vanschoonbeek et al. (6) and Khan et al. (7) could be attributed to the different patients (males or females), and the type and dose of medicine. On the other

**Table 3.** Comparison of Biochemical Parameters Among Different Groups<sup>a</sup>

Biochemical Factors	Diabetes	D+ Cinnamon (2g/100g Diet)	D+ Cinnamon (8g/100g Diet)	D + TF (2g/100g Diet)	D + TF (8g/100g Diet)	Control
<b>Serum</b>						
TC, mg/dL	140.1 ± 4.3	121.2 ± 5.2 <sup>b</sup>	111.3 ± 5.9 <sup>c</sup>	124.5 ± 4.6 <sup>b</sup>	119.9 ± 5.2 <sup>b</sup>	70.3 ± 5.0 <sup>d</sup>
TG, mg/dL	101.2 ± 5.4	90.1 ± 4.5	70.5 ± 4.6 <sup>d</sup>	80.8 ± 4.4 <sup>c</sup>	78.2 ± 2.7 <sup>b</sup>	68.4 ± 3.9 <sup>d</sup>
VLDL-C, mg/dL	21.1 ± 2.0	17.9 ± 1.3	15.1 ± 1.6 <sup>c</sup>	16.5 ± 1.1 <sup>c</sup>	16.1 ± 1.6 <sup>a</sup>	14.2 ± 1.8 <sup>d</sup>
HDL-C, mg/dL	34.5 ± 3.7	45.2 ± 3.4 <sup>b</sup>	45.4 ± 2.4 <sup>b</sup>	44.0 ± 4.1 <sup>b</sup>	49.1 ± 3.0 <sup>c</sup>	48.8 ± 9.7 <sup>b</sup>
LDL-C, mg/dL	84.8 ± 2.1	57.9 ± 2.3 <sup>b</sup>	27.1 ± 0.6 <sup>d</sup>	65.4 ± 4.7 <sup>b</sup>	54.1 ± 0.6 <sup>b</sup>	12.7 ± 2.1 <sup>d</sup>
AST U/L	56.7 ± 5.4	41.0 ± 2.9 <sup>b</sup>	39.2 ± 3.8 <sup>b</sup>	42.2 ± 3.6 <sup>b</sup>	44.3 ± 2.8 <sup>b</sup>	42.2 ± 3.6 <sup>b</sup>
ALT, U/L	63.9 ± 4.6	42.8 ± 3.8 <sup>b</sup>	42.2 ± 3.2 <sup>c</sup>	45.5 ± 3.1 <sup>b</sup>	44.6 ± 3.1 <sup>b</sup>	43.4 ± 2.5 <sup>b</sup>
<b>Liver</b>						
TC (mg/g tissue)	8.4 ± 0.11	6.2 ± 0.22 <sup>c</sup>	4.9 ± 0.34 <sup>d</sup>	5.2 ± 0.20 <sup>d</sup>	5.0 ± 0.19 <sup>d</sup>	4.5 ± 0.18 <sup>d</sup>
TG (mg/g tissue)	9.7 ± 0.24	6.6 ± 0.33 <sup>c</sup>	5.1 ± 0.28 <sup>d</sup>	6.1 ± 0.13 <sup>d</sup>	5.9 ± 0.23 <sup>d</sup>	4.9 ± 0.12 <sup>d</sup>
Glycogen(mg/g tissue)	22.8 ± 3.2	43.8 ± 2.5 <sup>d</sup>	55.0 ± 1.5 <sup>d</sup>	37.7 ± 2 <sup>b</sup>	40.0 ± 3.4 <sup>c</sup>	44.5 ± 3.1 <sup>d</sup>

Abbreviations: TC, total cholesterol, TG, triglycerides, VLDL-C, very low-density lipoprotein cholesterol, HDL-C, high-density lipoprotein cholesterol. LDL-C, low density lipoprotein cholesterol and TF, *Trigonella foenum*.

<sup>a</sup>Data represent as mean ± SEM (n = 6).

<sup>b</sup>P < 0.05.

<sup>c</sup>P < 0.01.

<sup>d</sup>P < 0.001 compared with diabetic animals.

hand, Crawford et al. (8) showed that daily supplementation of 1g Cinnamon for three months was safe and led to reduction of blood glucose and HbA1C levels in type 2 diabetics. In our study, diabetic rats, which were treated with high dose of Cinnamon and *Trigonella foenum* (8 g of Cinnamon or *Trigonella foenum* /100g diet) lipid profile, blood glucose and liver enzymes were normalized. In the animals which were treated with 2 g of Cinnamon or *Trigonella foenum* /100g diet, the change in lipid profile and liver enzymes were not significant. The results of previous experiments showed that treatment of diabetic rats with Cinnamon significantly reduced blood glucose, total cholesterol, triglyceride, LDL-C and VLDL-C (7, 9, 21).

The results of the study of Sharma et al. (10) showed that administration of 100 g *Trigonella foenum* powder/day for 21 days in dyslipidemic patients, significantly reduced triglyceride, total cholesterol, LDL-C and VLDL-C levels.

There was a report that showed treatment of type 1 and 2 diabetics patients with *Trigonella foenum* reduced total cholesterol, LDL-C, blood glucose and improved insulin resistance (22, 23). However, the hypoglycemic and hypolipidemic mechanism of this plant is unknown so far. In the recent study, *Trigonella foenum* significantly reduced AGEs and fructosamine formation. This plant also reduced protein carbonyl contents and thiol group oxidation in a dose-dependent manner. Therefore, inhibition of AGEs formation is one of the proposed antidiabetic mechanisms by *Trigonella foenum*. Vijayakumar et al. (24) reported that


*Trigonella foenum* reduced blood lipid by suppression of fat accumulation and also up-regulated of LDL-receptor expression. On the other hand, hypoglycemic and hypolipidemic mechanisms of Cinnamon are not yet completely clear. In this study, Cinnamon extract suppressed in vitro AGEs formation, which is involved in diabetic complications.

A previous study showed that Cinnamon motivated insulin receptor kinase and suppressed dephosphorylation of the insulin receptor, leading to maximal phosphorylation of the insulin receptor (7). Maximal phosphorylation of this receptor leads to increase in insulin activity; consequently normalization of blood lipid and glucose levels. It has been reported that Cinnamon motivated the activity of glycogen synthase enzymes and increased uptake of glucose from blood circulation. All of these properties can lead to increase in insulin sensitivity and glucose reduction (7).

Lee et al. (25) proved that prescription with Cinnamon in hyperlipidemic animals markedly declined total cholesterol concentration via suppression of hepatic hydroxyl methylglutaryl-coenzyme A (HMG-CoA) reductase, the rate-limiting enzyme of cholesterol biosynthesis.

It has been shown that about 80% of the LDL-C uptake is through LDL-receptor that consequently reduces blood cholesterol (7). We recently showed that Cinnamon and *Trigonella foenum* at a high dose significantly increased LDL-receptor gene expression (unpublished data). Several ex-

**Figure 1.** Total Antioxidant (A) and Thiol Group Levels (B) in Different Groups of Diabetic and Control Animals After 30 Days of Treatment




\*P < 0.05, \*\*P < 0.01 and \*\*\*P < 0.001 compared with diabetic rats. Data are presented as means ± SEM. TF: *Trigonella foenum*; D: diabetes.

periments have indicated that increasing the activity of cholesterol 7-alpha hydroxylase could decline the plasma total cholesterol by increasing the conversion of cholesterol to bile acids (26). In our study, cholesterol 7-alpha hydroxylase activity was significantly reduced by Cinnamon and *Trigonella foenum* at a dose of 8 g /100g diet. In this study we established that Cinnamon and *Trigonella foenum* extracts also have potential antioxidant activity, which would lead to further useful effects such as hypoglycemic and liver regeneration properties.


Triglyceride and cholesterol accumulation in liver tissues were noticeably observed in diabetic animals (27). The obtained data indicated an accumulation in lipid content of liver (triglyceride and cholesterol) in diabetic rats that

**Figure 2.** Liver (A) and Heart (B) Malondialdehyde Levels in Different Groups of Diabetic and Control Animals After 30 Days of Treatment



\*P < 0.05 and \*\*P < 0.01 compared with diabetic rats. Data are presented as means ± SEM. TF: *Trigonella foenum*; D: diabetes.

**Figure 3.** Advanced Glycosylation End Products Formation in Serum and Kidney of Different Groups of Diabetic and Control Animals After Thirty Days of Treatment




Data are presented as means ± SEM. TF: *Trigonella foenum*; D: diabetes.

was significantly reduced after treatment with Cinnamon or *Trigonella foenum* extract diet.


It is well known that STZ-induced diabetes in animals led to cytotoxicity and disruption of pancreatic  $\beta$  cell membrane (27). In the treated diabetic groups, morphological changes and liver enzyme were significantly normalized.

**Figure 4.** Cholesterol 7-Alpha Hydroxylase (nmol/mg protein) Level and Lipase Activity (mU/mg protein) in Different Groups of Diabetic and Control Animals After Thirty Days of Treatment



\*P < 0.05 and \*\*P < 0.01 compared with diabetic rats. Data are presented as means ± SEM. TF: *Trigonella foenum*; D: diabetes.

**Figure 5.** The Morphological Changes in Liver of Different Treated Groups



Liver histology in control group showed normal structure including; a: normal cellular construction with portal triad, numerous hepatocytes, b: central vein, c: lining sinusoidal, d: accumulation of lipid.

### 5.1. Conclusion

The results of this study indicated that administration of Cinnamon and *Trigonella foenum* extract in STZ-induced diabetic rats normalized dyslipidemia. These herbal medicines also showed potential antiglycation activity in vitro. Increasing the activity of cholesterol 7-alpha hydroxylase following treatment with these plants is another new possible mechanism for their hypolipidemic properties. Also, the liver antioxidants significantly increased and the histopathological changes were normalized in diabetic animals treated by Cinnamon and

*Trigonella foenum* extract. Hence, these herbal medicines are suggested to be used as anti-diabetic agents; however more studies on human subjects are required.

### Footnotes

**Authors' Contribution:** Seyed Mehrdad Kassae: prepared the draft of manuscript, and performed the antioxidants test; Mohammad Taghi Goodarzi: designed and supervised the experiment, edited the manuscript, and interpreted the results; Ebrahim Abbasi Oshaghi: performed



the glycation test, animal handling and interpreted the results.

**Funding/Support:** The authors would like to acknowledge the Hamedan Branch Islamic Azad University, Hamedan, Iran, for financial support of this study.

## References

- Ahmadi H, Azar ST. Liver disease and diabetes: association, pathophysiology, and management. *Diabetes Res Clin Pract.* 2014;**104**(1):53-62. doi: [10.1016/j.diabres.2014.01.003](https://doi.org/10.1016/j.diabres.2014.01.003). [PubMed: [24485856](https://pubmed.ncbi.nlm.nih.gov/24485856/)].
- Guariguata L, Whiting DR, Hambleton I, Beagley J, Linnenkamp U, Shaw JE. Global estimates of diabetes prevalence for 2013 and projections for 2035. *Diabetes Res Clin Pract.* 2014;**103**(2):137-49. doi: [10.1016/j.diabres.2013.11.002](https://doi.org/10.1016/j.diabres.2013.11.002). [PubMed: [24630390](https://pubmed.ncbi.nlm.nih.gov/24630390/)].
- Medagama AB. The glycaemic outcomes of Cinnamon, a review of the experimental evidence and clinical trials. *Nutr J.* 2015;**14**:108. doi: [10.1186/s12937-015-0098-9](https://doi.org/10.1186/s12937-015-0098-9). [PubMed: [26475130](https://pubmed.ncbi.nlm.nih.gov/26475130/)].
- Patel DK, Prasad SK, Kumar R, Hemalatha S. An overview on antidiabetic medicinal plants having insulin mimetic property. *Asian Pac J Trop Biomed.* 2012;**2**(4):320-30. doi: [10.1016/S2221-1691\(12\)60032-X](https://doi.org/10.1016/S2221-1691(12)60032-X). [PubMed: [23569923](https://pubmed.ncbi.nlm.nih.gov/23569923/)].
- Wang D, Wang J, Liu Y, Zhao Z, Liu Q. Roles of Chinese herbal medicines in ischemic heart diseases (IHD) by regulating oxidative stress. *Int J Cardiol.* 2016;**220**:314-9. doi: [10.1016/j.ijcard.2016.06.161](https://doi.org/10.1016/j.ijcard.2016.06.161). [PubMed: [27390948](https://pubmed.ncbi.nlm.nih.gov/27390948/)].
- Vanschoonbeek K, Thomassen BJ, Senden JM, Wodzig WK, van Loon LJ. Cinnamon supplementation does not improve glycemic control in postmenopausal type 2 diabetes patients. *J Nutr.* 2006;**136**(4):977-80. [PubMed: [16549460](https://pubmed.ncbi.nlm.nih.gov/16549460/)].
- Khan A, Safdar M, Ali Khan MM, Khattak KN, Anderson RA. Cinnamon improves glucose and lipids of people with type 2 diabetes. *Diabetes Care.* 2003;**26**(12):3215-8. [PubMed: [14633804](https://pubmed.ncbi.nlm.nih.gov/14633804/)].
- Crawford P. Effectiveness of cinnamon for lowering hemoglobin A1C in patients with type 2 diabetes: a randomized, controlled trial. *J Am Board Fam Med.* 2009;**22**(5):507-12. doi: [10.3122/jabfm.2009.05.080093](https://doi.org/10.3122/jabfm.2009.05.080093). [PubMed: [19734396](https://pubmed.ncbi.nlm.nih.gov/19734396/)].
- Pham AQ, Kourlas H, Pham DQ. Cinnamon supplementation in patients with type 2 diabetes mellitus. *Pharmacotherapy.* 2007;**27**(4):595-9. doi: [10.1592/phco.27.4.595](https://doi.org/10.1592/phco.27.4.595). [PubMed: [17381386](https://pubmed.ncbi.nlm.nih.gov/17381386/)].
- Kumar P, Kale RK, Baquer NZ. Antihyperglycemic and protective effects of Trigonella foenum graecum seed powder on biochemical alterations in alloxan diabetic rats. *Eur Rev Med Pharmacol Sci.* 2012;**16 Suppl 3**:18-27. [PubMed: [22957414](https://pubmed.ncbi.nlm.nih.gov/22957414/)].
- Abbasi Oshaghi E, Tavilani H, Khodadadi I, Goodarzi MT. Dill tablet: a potential antioxidant and anti-diabetic medicine. *Asian Pac J Trop Biomed.* 2015;**5**:720-7. doi: [10.1016/j.apjtb.2015.06.012](https://doi.org/10.1016/j.apjtb.2015.06.012).
- Oshaghi EA, Khodadadi I, Tavilani H, Goodarzi MT. Aqueous Extract of Anethum Graveolens L. has Potential Antioxidant and Antiglycation Effects. *Iran J Med Sci.* 2016;**41**(4):328-33. [PubMed: [27365555](https://pubmed.ncbi.nlm.nih.gov/27365555/)].
- Caengprasath N, Ngamukote S, Makynen K, Adisakwattana S. The protective effects of pomelo extract (*Citrus grandis* L. Osbeck) against fructose-mediated protein oxidation and glycation. *EXCLI J.* 2013;**12**:491-502. [PubMed: [26966424](https://pubmed.ncbi.nlm.nih.gov/26966424/)].
- Javad H, Seyed-Mostafa HZ, Farhad O, Mehdi M, Ebrahim AO, Nader RG. Hepatoprotective effects of hydroalcoholic extract of *Allium hirtifolium* (Persian shallot) in diabetic rat. *J Basic Clin Physiol Pharmacol.* 2012;**23**(2):83-7. doi: [10.1515/jbcp-2012-0017](https://doi.org/10.1515/jbcp-2012-0017).
- Abbasi Oshaghi E, Khodadadi I, Saidijam M, Yadegarazari R, Shabab N, Tavilani H, et al. Lipid Lowering Effects of Hydroalcoholic Extract of *Anethum graveolens* L. and Dill Tablet in High Cholesterol Fed Hamsters. *Cholesterol.* 2015;**2015**:958560. doi: [10.1155/2015/958560](https://doi.org/10.1155/2015/958560). [PubMed: [26823981](https://pubmed.ncbi.nlm.nih.gov/26823981/)].
- Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem.* 1972;**18**(6):499-502. [PubMed: [4337382](https://pubmed.ncbi.nlm.nih.gov/4337382/)].
- Butler NA, Lee EY, Whelan WJ. A protein-bound glycogen component of rat liver. *Carbohydr Res.* 1977;**55**:73-82. [PubMed: [861979](https://pubmed.ncbi.nlm.nih.gov/861979/)].
- Folch J, Lees M, Sloane Stanley GH. A simple method for the isolation and purification of total lipides from animal tissues. *J Biol Chem.* 1957;**226**(1):497-509. [PubMed: [13428781](https://pubmed.ncbi.nlm.nih.gov/13428781/)].
- Okudan N, Belviranli M. Effects of exercise training on hepatic oxidative stress and antioxidant status in aged rats. *Arch Physiol Biochem.* 2016;**1-6**. doi: [10.1080/13813455.2016.1199574](https://doi.org/10.1080/13813455.2016.1199574). [PubMed: [27424521](https://pubmed.ncbi.nlm.nih.gov/27424521/)].
- Aydin Y, Onder E. Herbal self-medication use in Type 2 diabetes mellitus. *Turk J Med Sci.* 2016;**46**(4):1275-6. doi: [10.3906/sag-1507-97](https://doi.org/10.3906/sag-1507-97). [PubMed: [27513437](https://pubmed.ncbi.nlm.nih.gov/27513437/)].
- Sahib AS. Anti-diabetic and antioxidant effect of cinnamon in poorly controlled type-2 diabetic Iraqi patients: A randomized, placebo-controlled clinical trial. *J Intercult Ethnopharmacol.* 2016;**5**(2):108-13. doi: [10.5455/jice.20160217044511](https://doi.org/10.5455/jice.20160217044511).
- Gupta A, Gupta R, Lal B. Effect of Trigonella foenum-graecum (fenugreek) seeds on glycaemic control and insulin resistance in type 2 diabetes mellitus: a double blind placebo controlled study. *J Assoc Physicians India.* 2001;**49**:1057-61. [PubMed: [11868855](https://pubmed.ncbi.nlm.nih.gov/11868855/)].
- Snehlata HS, Payal DR. Fenugreek (*Trigonella foenum-graecum* L.): an overview. *Int J Curr Pharm Rev Res.* 2012;**2**:169-87.
- Vijayakumar MV, Singh S, Chhipa RR, Bhat MK. The hypoglycaemic activity of fenugreek seed extract is mediated through the stimulation of an insulin signalling pathway. *Br J Pharmacol.* 2005;**146**(1):41-8. doi: [10.1038/sj.bjp.0706312](https://doi.org/10.1038/sj.bjp.0706312). [PubMed: [15980869](https://pubmed.ncbi.nlm.nih.gov/15980869/)].
- Cheng SS, Liu JY, Hsui YR, Chang ST. Chemical polymorphism and antifungal activity of essential oils from leaves of different provenances of indigenous cinnamon (*Cinnamomum osmophloeum*). *Biore-sour Technol.* 2006;**97**(2):306-12. doi: [10.1016/j.biortech.2005.02.030](https://doi.org/10.1016/j.biortech.2005.02.030). [PubMed: [16171686](https://pubmed.ncbi.nlm.nih.gov/16171686/)].
- Li T, Chiang JY. Bile acids as metabolic regulators. *Curr Opin Gastroenterol.* 2015;**31**(2):159-65. doi: [10.1097/MOG.0000000000000156](https://doi.org/10.1097/MOG.0000000000000156). [PubMed: [25584736](https://pubmed.ncbi.nlm.nih.gov/25584736/)].
- Gupta D, Raju J, Prakash J, Baquer NZ. Change in the lipid profile, lipogenic and related enzymes in the livers of experimental diabetic rats: effect of insulin and vanadate. *Diabetes Res Clin Pract.* 1999;**46**(1):1-7. [PubMed: [10580609](https://pubmed.ncbi.nlm.nih.gov/10580609/)].