

Original Article

Essential Oil Composition, Total Phenol, Flavonoid, Anthocyanin and Antioxidant Activities in Different Parts of *Artemisia annua* L. in Two Localities (North of Iran)

Masoumeh Mazandarani^{1*}, Zahra Majidi¹, Parastoo Zarghami-Moghaddam², Mehdi Abrodi³, Helen Hemati⁴ and Fatemeh Fathiazad⁵

¹Department of Botany, Gorgan Branch, Islamic Azad University, Gorgan, Iran

²Research Center of Natural Products Safety and Medicinal Plants, North Khorasan University of Medical Science, Bojnurd, Iran

³Young Researchers Club, Gorgan Branch, Islamic Azad University, Gorgan, Iran

⁴Baharan Higher Education Institute of Gorgan, Gorgan, Iran

⁵Department of Pharmacognosy, Faculty of Pharmacy, Tabriz University of Medical Sciences, Tabriz, Iran

Article history: Received: 3 April 2012/Accepted in revised form: 4 July 2012

© 2012 Iranian Society of Medicinal Plants. All rights reserved.

Abstract

In this study, we evaluate the different biological activities of *Artemisia annua* L., locally known as "Moureh", in various altitudes in North of Iran, which has been used as sedative, fever few, anti inflammation, insecticide and anti infection to treat many current diseases. Parts of plants were collected from two different localities (23-1000 m) in Mazandaran province, North of Iran. The most important of secondary metabolites of total phenolics (TP), total flavonoids (TF) and total anthocyanin (TA) content of extracts were investigated by spectrophotometry method and their antioxidant activity were obtained by Total Antioxidant Capacity (TAC), Reducing Power (RP) and 1,1-diphenyl-2-picryl hydrazyl radical scavenging (DPPH). The essential oils were obtained by hydro distilled in a Clevenger-type apparatus for 5h and analyzed by GC/MS. Results indicate that the main components in essential oils were identified as *Artemisia* ketone (25.54- 13.6%), followed by 1,8-cineole (11.98-13.26%), camphor (11.89-13.68%), α -pinene (10.11-9.29%) in AFRACHAL (1000m) and DOLAT ABAD (23m) regions, respectively. TP content had significant variation in different plant parts and regions, ranging from (11.22 to 16.94) mgGAEg⁻¹, TF content (11.62 to 63.74) mgQUE g⁻¹ and quantity of TA (0.03 to 3.59) mgCGEg⁻¹. The highest contents of secondary metabolites were found in aerial parts when compared with the other parts. Amount of antioxidant activity (IC₅₀) in various parts of *A. annua* L. was measured (1.98 to 4.2) in DPPH, (7.07 to 7.46) in TAC and (5.26 to 8.04) in RP methods. In general, the highest contents of activities were found in aerial parts when compared with the other parts, whereas this part with the highest amount of IC₅₀ had the weakest antioxidant activity.

Key words: *Artemisia annua* L., Essential oil, Phenolic content, Antioxidant capacity, Iran

Introduction

Reactive oxygen species including free radicals and non free radicals along with different forms of active oxygen are involved in diverse physicochemical processes and causing more than one hundred disorders in humans such as neurodegenerative disorders, cancer, ischemia, cardiovascular diseases, atherosclerosis, arthritis, AIDS, Alzheimer's, Parkinson's, cataracts and inflammation. Antioxidants prevent diseases by mechanisms such as scavenging free radicals, against oxidative stress and inhibiting lipid peroxidation [31].

Therefore supplementing a food product with antioxidant compounds of plant source may provide a health benefit as well. Secondary metabolites, especially phenol compounds and terpenes in medicine plants have been known to therapeutic activities like antioxidant, antimicrobial, anticancer, anti-inflammatory and etc. in many researches, high correlation was reported between the antioxidant capacity and total phenol and flavonoid contents of medicinal plants [39, 43]. The antioxidant activity of secondary metabolites is due to their redox properties, ability to chelate metals and quenching of singlet oxygen [41]. Flavonoids with chelate metals

* Corresponding author: Department of Botany, Gorgan Branch, Islamic Azad University, Gorgan, Iran

Email Address: Mazandarani.m@gorganiau.ac.ir

such as iron chelating have been recommended for malaria patients [19]. Different studies reported that phenolic compounds have protection against a wide range of diseases such as coronary heart disease, stroke, and certain types of cancers [21, 3] and exhibit biological activities such as: anti-inflammatory, immune-stimulating agents antiallergenic, anti-atherogenic, anti-microbial, antithrombotic, anti stress, anti hyperglycemia and vasodilatory effects [10].

Artemisia annua L. belong to Asteraceae family, comprising with up to 500 species in world and about approximately 34 native *Artemisia* spp. in Iran [44, 46]. Approximately 200 species of *Artemisia* grow in China, where more than 50 of them have been used in traditional medicine of this country [42], which the tea and decoction of *Artemisia annua* L. is famous to an anti-malarial drug plant that has been used for many centuries for treat human fever and malaria [2]. Also in ancient Greece and Roman the essential oil of *Artemisia* species were used in infusion as poison antidote, activates the blood circulation, anti-helminthic and for its abortive qualities indicated in gastric insufficiency [44]. *A. annua* L. is a rich source of biologically active natural products which has been used to treatment of malaria, gastrointestinal helminthosis, hemorrhoid, skin rashes, diarrhea and some other diseases [7-19, 46].

Many secondary metabolites of terpene peroxides such as artemisia ketone, artemisinic alcohol, arteannuin B and myrcene hydroperoxide, over 50 different phenolic compounds belonging to five major groups (flavones, flavonols, coumarins, phenolic acids, and a miscellaneous group) which the main coumarins are coumarin, aesculetin, isofraxidin, scopoletin, scopolin, tomentin, and also other phenolic compounds such as 2,4-dihydroxy-6-methoxy-acetophenone, 5-nonadecyl-3-O-methyletherresorcinol, 2,2,6-trihydroxychromene, and 2,2-dihydroxy-6-methoxychromene have been isolated of this plant [11-14-19,46], the high antioxidant activity of *A. annua* extract is due to its high phenolic content [19]. Artemisinin (sesquiterpene lactone) is a quite expensive compound with very low concentration which the only commercial source available is *A. annua* [2]. From approximately 400 species of artemisias, *A. annua* L. is the main source of artemisinin [20]. Different studies show that Artemisinin-derived drugs are effective against bacteria, fungi, insects and parasites such as *Plasmodium* spp., *Coccidia* spp., *Babesia* spp., *Leishmania* spp. *Neospora caninum* and *Schistosoma* spp. [4-26-28-29-45-49,50] that effective in the health of humans and animals. The aerial parts of *A. annua* L. are highly aromatic and known to give an essential oil of some economic importance, which is rich in monoterpenes, sesquiterpenes, diterpenes and phenyl propanoids [23].

The aim of the present study is to find out how the geographic location influences on the qualitative and quantitative composition of secondary metabolites and essential oil of *Artemisia annua* L. and evaluation of antioxidant activities of different parts of plant, which has been used by the rural healers in North of Iran to prevent and treat of malarial, parasitic, antibacterial, anti inflammation, sedative, dysmenorrhea, diarrhea and infectious disease.

Materials and Methods

Plant materials

The different parts (stem, root and aerial parts) of *Artemisia annua* L. were collected in AFRACHAL (1000 m) and DOLAT ABAD (23 m) regions of Mazandaran province in North of Iran during Sep to Oct 2010. This voucher of specimen was identified and has been deposited at the Herbarium Museum of the Islamic Azad University of Gorgan branch (Golestan province). The plant materials were dried and ground to a fine powder using a laboratory mill, were maintained at room temperature (21–23 °C), and protected from light.

Phytochemical tests

2, 2-Diphenyl-1-picryl hydrazyl (DPPH), butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), methanol and other compounds were purchased from international companies.

Extract preparation

Extract preparation for phytochemical tests

The dried stem, root and aerial parts of plant (5g) were extracted overnight with 100 ml of methanol, in a mechanical shaker at room temperature. Each extract of plant was filtered with Whatman No. 1 filter paper and stored at 4 °C.

Extract preparation for antioxidant activity

A 45 gr of different parts (root and aerial parts) of *Artemisia annua* L. were extracted with 300 ml of methanol solvent in a mechanical shaker at room temperature. Extracts were filtered with Whatman No. 1 filter paper. The filtrates obtained from extract were evaporated into dry at 40°C in a rotary from evaporator and stored at 4 °C [8].

Phytochemical tests

Total phenols determination

Total phenolic content were determined by Folin Ciocalteu method [35]. A 0.5 ml of samples or gallic acid (standard phenolic compound) was mixed with Folin Ciocalteu reagent (5 ml) and aqueous Na₂CO₃ (4 ml, 1 M). The mixtures were allowed to stand for 15 min and the total phenols were determined by colorimeters at 765 nm. Gallic acid was used as a standard for calibration curve. Total phenol values

are expressed in terms of mg equal gallic acid in 1 gr powder dry plant.

Total flavonoids determination

Total flavonoids content was estimated by the Aluminum chloride method, based on the procedure of [35]. Plant extracts (0.5 ml) were separately mixed with 1.5 ml of solvent, 0.1 ml of 10% aluminum chloride, 0.1 ml of 1 M potassium acetate and 2.8 ml of distilled water. It remained at room temperature for 30 min; the absorbance of the reaction mixture was measured at 415 nm with a spectrophotometer, and quercetin was used as a standard for calibration curve. Total flavonoids values are expressed in terms of mg equal quercetin in 1gr powder dry plant.

Total anthocyanin determination

Total anthocyanin content was determined by the pH-differential method described by Giusti [22], using 2 buffer systems: potassium chloride buffer, pH 1 (1.86 g KCl in 1 L of distilled water, pH value adjusted to 1.0 with concentrated HCl), and sodium acetate buffer, pH 4.5 (54.43 g CH₃CO₂Na·3H₂O in 1 L of distilled water, pH value adjusted to 4.5 with concentrated HCl). The sample diluted with corresponding buffer and the absorbance was measured at 510 and 700 nm. Total anthocyanins were calculated as cyanidin-3-glucoside according to the following equation:

$$TAC = \left(\frac{A \times MW \times DF \times 100}{MA} \right)$$

$A = (A_{510} - A_{700})_{pH1} - (A_{510} - A_{700})_{pH4.5}$
 MW: 449.2 g/mol for cyanidin-3-glucoside
 DF = dilution factor; MA: 26900

Antioxidant Activity Tests

Reducing Power assay

This assay is based on Arabshahi-Delouee method. First, The dried extract (12.5–1000 µg) in 1 ml of the corresponding solvent was combined with 2.5 ml of phosphate buffer (0.2 M, pH 6.6) and 2.5 ml of potassium ferricyanide (K₃Fe(CN)₆; 10 g l⁻¹), after the mixture was incubated at 50°C for 30 min. Then, 2.5 ml of trichloroacetic acid (100 g l⁻¹) were added and the mixture centrifuged at 1650g for 10 min. Then, 2.5 ml of the supernatant solution was mixed with 2.5 ml of distilled water and 0.5 ml of FeCl₃ (1 g l⁻¹), and the samples absorbance was measured at 700 nm [8].

1,1-diphenyl-2-picryl hydrazyl radical scavenging capacity Assay

The ability of the extracts for free radical scavenging was assessed by the method suggested by Arabshahi-Delouee and Urooj [8]. Briefly, 1ml of a 1mM methanolic solution of DPPH was mixed with 3ml of extract solution in methanol (containing 12.5–1000 µg of dried extract). The mixture was then vortexed

vigorously and left for 30 min at room temperature in the dark. The absorbance was measured at 517 nm and activity was expressed as percentage DPPH scavenging relative to control using the following equation:

$$\text{DPPH scavenging activity (\%)} = [(A_{\text{control}} - A_{\text{sample}}) / A_{\text{control}}] \times 100$$

Total Antioxidant Capacity

This experimental procedure was adapted from Arabshahi-Delouee method, which is based on the reduction of Mo (VI) to Mo (V) by the sample and observation of a green phosphate/Mo (V) complex at acidic pH. An aliquot of 0.1 ml of sample solution, containing 12.5–1000µg of dried extract in corresponding solvent, was combined in a tube with 1 ml of reagent solution (0.6 M sulphuric acid, 28 mM sodium phosphate, and 4 mM ammonium molybdate). They were incubated in a thermal block at 95 °C for 90 min. Then we got cold the samples and measured their absorbance at 695 nm. A typical blank solution contained 1 ml of reagent solution and the appropriate volume of the same solvent was used for the sample, and was incubated under the same conditions as the rest of the samples [8].

Isolation of essential oil

The essential oils of aerial parts of plant in blooming were obtained by hydro distilled in a Clevenger-type apparatus for 5h according to the method recommended in the British Pharmacopoeia and was kept refrigerated and protected from direct light until the analysis time.

Gas Chromatography - Mass Spectrometry (GC-MS)

The essential oil was analyzed by GC/MS (6890 N Network GC system and 5975 B Intet MSD). The capillary conditions were as follows; carrier gas, helium with a flow rate of 1.7ml/min; injected 1µL of the essential oil and ionization potential 70ev. The initial temperature of column was 50 °C (held 5min) then heated to 250 °C, then heated to 250 °C and kept constant for 20min. The same condition of temperature programming used for n-alkenes mixture (C7-C31) to calculate the retention indices (RI). The identification of each component was studied by mass spectral data, literature (Adams R, 1995) and NIST computer library. The relative percentage of the oil constituent was calculated.

Statistical analysis

For all assays, data were expressed as means ± S.E. and differences at $P < 0.05$ were considered statistically significant.

Results and Discussion

Total phenolics, flavonoids and anthocyanin

Secondary metabolites contents of different parts of *Artemisia annua* L. in various regions show in Table

1, comparison of the results indicated that the TP content of various parts of *A. annua* L. had significant variation, ranging from 11.22 ± 0.74 to 16.94 ± 0.52 mgGAEg⁻¹, TF content 11.62 ± 0.59 to 63.74 ± 1.01 mgQUE g⁻¹ and quantity of TA 0.03 ± 0.002 to 3.59 ± 0.4 mgCGEg⁻¹. The highest contents of secondary metabolites of TP (15.58 and 16.94 mgEGAg⁻¹), TF (57.28 and 63.74 mgEQUg⁻¹) and TA (3.59 and 3.34 mgECGg⁻¹) were found in aerial parts when compared with the other parts, whereas the lowest contents TP (11.22 mgGAEg⁻¹) and TF (11.62 mgQUE g⁻¹) were in stem extract of AFRACHAL region (1000m), and the lowest TA detected in the root extract of DOLAT ABAD region (23m) (Table 1 and Fig 1-2,3).

Amount of secondary metabolites (phenolics, flavonoids and anthocyanin) has been associated with increased levels of reactive oxygen species, which are by products of aerobic metabolism or with biotic, abiotic and stresses [6, 17]. Results of Seddik [38] show that high flavonoids and polyphenol contents of *A. herba alba* Asso. were present in the ethyl acetate

phase, while the aqueous phase contains smaller amounts of these compounds.

Also in another studies by Brown [13] indicate that the artemisinin content of *A. annua* has ranging between 0.01% and 1%, depending on variety, and can even be as high as 1.4% in some cultivated strains. Flavonoids are well known for their antioxidant activity due to their redox properties, which can delay or inhibit oxidation of lipids and other molecules by inhibiting the beginning of oxidizing chain reactions [19]. Survey of results showed that was a negative correlation between total phenol and flavonoid contents and antioxidant activity for aerial parts extract of plant. These results were in opposite with the findings of many research groups who reported direct relationships between total phenolic content and antioxidant activity [15-18-27-35-41, 43]. Also in other studies about variety of species (*Sylibum marrianum*, *Lithospermum erythrorhizon*, *Cordia multispicata*, *C. multispicata* and *Tournefortia bicolor*, *Ehretia laevis*, *Cordia myxa* and *Borago officinalis*), showed direct relationships between of them [16-24, 33].

Table 1 Comparison of secondary metabolites of different parts of *Artemisia annua* L. in various regions

Metabolite	(AFRACHAL region) (1000 m)			(DOLAT ABAD region) (23 m)		
	Aerial part	Stem	Root	Aerial part	Stem	Root
Flavonoid (mg QUE g ⁻¹)	63.74 ± 1.01	11.62 ± 0.59	21.76 ± 0.33	57.28 ± 0.25	21.48 ± 0.66	11.88 ± 0.89
Phenol (mg GAE g ⁻¹)	16.94 ± 0.52	11.22 ± 0.74	14.9 ± 0.6	15.58 ± 0.89	13.96 ± 0.86	13.24 ± 0.66
Anthocyanin (mg ECG g ⁻¹)	3.59 ± 0.4	0.26 ± 0.11	0.56 ± 0.09	3.34 ± 0.36	0.31 ± 0.05	0.03 ± 0.002




Fig 1. Total phenol contents of different parts of *Artemisia annua* L.
Sample 1. AFRACHAL region (1000 m), Sample 2. DOLAT ABAD region (23 m)




Fig 2. Total flavonoid contents of various parts of *Artemisia annua* L.
Sample 1. AFRACHAL region (1000 m), Sample 2. DOLAT ABAD region (23 m)




Fig 3. Total anthocyanin contents of different parts of *Artemisia annua* L.
Sample 1. AFRACHAL region (1000m), Sample 2: DOLAT ABAD region (23m)

Antioxidant activity

Fig 4-6 and Table 2 shows antioxidant activity by inhibition of the free radical. Amount of IC₅₀ in various parts of *A. annua* L. was (1.98±0.21 to 4.2±0.63) in DPPH method, (7.07±0.86 to 7.46±0.86) in TAC and (5.26±0.87 to 8.04±0.84) in RP methods. The highest antioxidant activity and radical scavenging effect were observed in root extract with IC₅₀ (1.98 to 2.2) in Dpph, (5.26 to 5.83) in PR and (6.89 to 7.01) in TAC methods, whereas aerial parts with the highest amount of IC₅₀ had the weakest antioxidant activity.

Reactive oxygen species are involved in diverse physicochemical processes in the human body [36] which have main role in the pathogenesis of different diseases [16].

In research by Brisibe [12], total antioxidant capacity reported for the leaves 1,125 µmol TE/g dry matter

and 1,234 µmol TE/g dry matter for the inflorescences and the antioxidant activity values for the stems and roots were similar with 292 and 287 µmol TE/g dry matter, respectively.



Total Antioxidants in Hot-water extracts of *Artemisia monosperma* reported 675.33 µmol Trolox per 100 ml [5].

The traditional tea of *A. annua* is a main source of both antioxidant phenolics (mostly flavonoids) and artemisinin [37, 48].

Another report indicates that the high antioxidant activity of *A. annua* L. extract is most likely due to its high phenolic content and this capacity is stable to boiling [19]. The results of Laciari [30] show that *A. echegarayi* essential oil presented moderate antioxidant activity and its antioxidant activity was lower than that of quercetin, a powerful natural antioxidant.

Table 2 Comparison of IC₅₀ of different parts of *Artemisia annua* L. in various regions

Antioxidant activity	Roots		Aerial parts		BHA	BHT
	(DOLAT ABAD region)	(AFRACHAL region)	(DOLAT ABAD region)	(AFRACHAL region)		
IC ₅₀ TAC	7.01 ± 0.88	6.89 ± 0.88	7.46 ± 0.86	7.07 ± 0.86	3.46 ± 0.88	3.09 ± 0.84
IC ₅₀ RP	5.26 ± 0.87	5.83 ± 0.9	8.04 ± 0.84	7.56 ± 0.88	2.79 ± 0.95	2.6 ± 0.91
IC ₅₀ DPPH	2.2 ± 0.43	1.98 ± 0.21	4.2 ± 0.51	2.4 ± 0.32	-	-

**Fig 4.** Amounts of IC₅₀ in various parts of *Artemisia annua* L. in DPPH method**Fig 5.** Amounts of IC₅₀ in various parts of *Artemisia annua* L. in TAC method




Fig 6. Amounts of IC₅₀ in various parts of *Artemisia annua* L. in RP method

Table 3 The main chemical constituents of the flowering aerial parts of *Artemisia annua* L. essential oil

Compound	AFRACHAL region (1000 m)	DOLAT ABAD region (23 m)
α -pinene	10.11	9.29
Camphene	2.99	2.77
Thujene	1.46	1.89
β -pinene	1.24	0.94
Myrcene	2.57	1.93
1,8-cineole	13.26	11.98
γ -terpinene	0.72	1.12
Artemisia ketons	25.54	13.6
Artemisia alcohol	2.08	2.09
Camphore	11.89	13.68
Pinocarvone	1.03	2.44
Borneol L	0.69	0.84
Naphthalene	5.09	7.01

Essential oil

The essential oil analysis of *Artemisia annua* L. led to the identification of approximately 43 main compounds accounting for the 99.99% of the total essential oil in AFRACHAL (1000m) and 47 compounds accounting for the 93.71% in DOLAT ABAD (23m) regions (Table 3). The main essential oil components were identified as *Artemisia* ketenes (25.54- 13.6%), followed by 1,8-cineole (11.98-13.26%), camphor (11.89-13.68%), α -pinene (10.11-9.29%) in AFRACHAL (1000 m) and DOLAT ABAD (23 m) regions respectively. The chemical composition of the essential oil had been demonstrated in Table 3.

In previous study by Verdian-rizi [47] in Iran, the major constituents (48%), 1,8-cineole (9.39%), camphene (6.98%) and spathulenol (4.69%) were identified in the pre-flowering stage, in the volatile of flowering stage camphor (43.50%), 1,8-cineole (13.90%), spathulenol (3.73%) and *Artemisia* ketone (3.37%) and in the oil obtained from post-flowering stage, camphor (36.75%), 1,8-cineole (12.00%), spathulenol (4.50%) and pinene were the principal components of this oil.

The majority of the compounds identified in the essential oil of *Artemisia annua* growing in Ethiopia reported monoterpenes (57.89%), Sesquiterpenes (36.84%) and phenols comprised (5.55%). Among

the identified monoterpenes 52.17% were monoterpene alcohols, 30.43% - monoterpene hydrocarbons, 13.04% - monoterpene aldehydes and 4.35 % were monoterpene oxides, and camphor was identified as the major component (43.84%) [34].

In other study on the chemistry of *A. annua* L. by Juteau [25] show that the essential oil of *A. annua* L. aerial parts of species growing in France consisted of camphor (44%), germacrene D (16%), *trans*-pinocarveol (11%), β -selinene (9%), β -caryophyllene (8.9%) and Artemisia ketone (3%). In fact essential oil contents show variations in plants of different geographical origin.

The essential oil of *A. annua* aerial parts, that were obtained from plants growing at India consisted of main constituents artemisia ketone (58.8%), 1,8-cineole (10.2%) and camphor (15.8%) while the chief components of the essential oil from aerial parts of plants grown in Kashmir contained artemisia ketone (52.3%), 1,8-cineole (13.1%) and camphor (15.5%) [9].

Accordingly, it seems differences in among of chemical compositions of the *A. annua* L. essential oils from the viewpoint of qualitative and quantitative might depended on different geographical and environmental conditions such as different temperature, soils and altitudes which play an important role on the biological activity of *A. annua* L. essential oil.

Conclusion

The present study indicate the aerial part of *Artemisia annua* L. with highest amount of TP, TF and TA compounds as an important part of this plant, which could provide potential natural sources compounds to prevent and treatment of diseases.

These findings demonstrate the aerial part as the best organ with the most secondary metabolites of plant for future research and also confirmed the interesting of this plant uses by the rural healers to prevent and treat of current infectious disease.

Acknowledgements

The authors are grateful to technical help of laboratory officer of RCMP (Research center of Medicine Plants), Islamic Azad University of Gorgan branch.

References

1. Adams R. Identification of essentials oil components by Gas Chromatography/ Mass Spectroscopy. Allured Publishing Corp. Carol Stream, Illinois. 1995;1-456.
2. Aftab T, A.Khan MM, Idrees M, Naeem M, Hashmi MN. Methyl jasmonate counteracts boron toxicity by preventing oxidative stress and regulating antioxidant enzyme activities and artemisinin biosynthesis in *Artemisia annua* L. Identification of intermediates and enzymes involved in

- Artemisia annua* L. Protoplasma. 2010; doi:10.1007/s0070901002185
3. Aggarwal BB. Targeting inflammatory pathways for chronic diseases by phytochemicals derived from spices, fruits, vegetables, and traditional remedies. Acta Hort. 2009;841:33-46.
4. Allen PC, Lydon J, Danforth HD. Effects of components of *Artemisia annua* on coccidia infections in chickens. Poult. Sci. 1997;76:1156-1163.
5. Al-Soqeer A. Antioxidant Activity and Biological Evaluation of Hot-water Extract of *Artemisia monosperma* and *Capparis spinosa* Against Lead Contamination. Res. J. Bot. 2011;6:11-20.
6. Apel K, Hirt HM. Reactive oxygen species: metabolism, oxidative stress, and signal transduction. Ann. Rev. Plant Biol. 2004;55:373-379.
7. Arab HA, Mardjanmehr SH, Shahbazfar A, Rassouli A, Abdollahi M, Nekouie O. Toxicopathologic Effects of Artemisinin in Broiler Chickens Following a Single Oral Dose: An LD Study. Int. J. Poultry Sci. 2009; 8:808-812.
8. Arabshahi-Delou S, Urooj A. Antioxidant properties of various solvent extracts of mulberry (*Morus indica* L.) leaves. Food Chem. 2007;102:1233-1240.
9. Ateeque A, Misra LN. Terpenoids from *Artemisia annua* and constituents of its essential oil, Int J Plant Biochem. 1994;37:183-186.
10. Balasundram N, Sundram K, Samman S. Phenolic compounds in plants and agri-industrial by products: antioxidant activity, occurrence, and potential uses. Food Chem. 2006;99:191-203.
11. Berteau CM, Freije JR, van der Woude H, Verstappen FWA, Perk L, Marquez V, De Kraker JW, Posthumus MA, Jansen BJM, de Groot A, Franssen MCR, Bouwmeester HJ. The early steps of artemisinin biosynthesis in *Artemisia annua*. Planta Med. 2005;71:40-47.
12. Brisibe EA, Umoren UE, Brisibe F, Magalhes PM, Ferreira JFS, Luthria D, Wuh X, Prior RL. Nutritional characterisation and antioxidant capacity of different tissues of *Artemisia annua* L. Food Chem. 2009;115:1240-1246.
13. Brown GD. The Biosynthesis of Artemisinin (Qinghaosu) and the Phytochemistry of *Artemisia annua* L. (Qinghao). Molecules 2010;15:7603-7698.
14. Brown GD, Liang GY, Sy L. Terpenoids from the seeds of *Artemisia annua*. Phytochem 2003;64:303-323.
15. Cai Y, Luo Q, Sun M, Corke H. Antioxidant activity and phenolic compounds of 112 traditional Chinese medicinal plants associated with anticancer. Life Sci. 2004;74(17):2157-2184.
16. Conforti F, Sosa S, Marrelli M, Menichini F, Statti GA, Uzunov D, Tubaro A, Menichini F, Loggia RD. In vivo anti-inflammatory and in vitro antioxidant activities of Mediterranean dietary plants. J. Ethnopharmacol. 2008;116:144-1451.
17. Dixon RA, Paiva NL. Stress-induced phenylpropanoid metabolism. Plant Cell 1995; 7:1085-1097.
18. Dorman HJD, Bachmayer O, Kosar M, Hiltunen R. Antioxidant properties of aqueous extracts from selected Lamiaceae species grown in Turkey. J. Agri. Food Chem. 2004;52(4):762-770.
19. Ferreira JFS, Luthria DL, Sasaki T, Heyerick A. Flavonoids from *Artemisia annua* L. as Antioxidants and Their Potential Synergism with Artemisinin against Malaria and Cancer. Molecules 2010;15:3135-3170.
20. Ferreira JFS. *Artemisia annua* L.: The hope against malaria and cancer, [Medicinal and Aromatic Plants:

- Production, Business & Applications. Proceedings of the Jan 15-17/2004 meeting. Mountain State University, Beckley, WV. 2004;56-61.
21. George TW, Niwat C, Waroonphan S, Gordon MH, Lovegrove JA, Paterson E. Effect of chronic and acute fruit and vegetable juice consumption on cardiovascular disease risk factor. *Acta Hort.* 2009;841:201-206.
 22. Giusti MM, Wrolstad RE. Anthocyanins, Characterization and measurement with UV-visible Spectroscopy. *Current protocols in food analytical chemistry* (R.E. Wrolstad). Wiley, New York, 2001. F1.2.1-F1.1.13.
 23. Goel D, Singh V, Ali M, Mallavarupu GR, Kumar S. Essential oils of petal, leaf and stem of the antimalarial plant *Artemisia annua*. *J. Nat. Med.* 2007;61:187-191.
 24. Hadaruga DI, Hadaruga NG. Antioxidant Activity of Hepatoprotective Silymarin and *Silybum marianum* L. *Extract Chem. Bull. POLITEHNICA Univ. (Timisoara).* 2009;54(68):2.
 25. Juteau F, Masotti V, Bessiere JM, Dherbomez M, Viano J. Antibacterial and antioxidant activities of *Artemisia annua* essential oil, *Fitoterapia.* 2002;73:532-535.
 26. Kim JT, Park JY, Seo HS, Oh HG, Noh JW, Kim JH, Kim DY, Youn HJ. In vitro antiprotozoal effects of artemisinin on *Neospora caninum*. *Vet. Parasitol.* 2002;103:53-63.
 27. Kırca A, Arslan E. Antioxidant capacity and total phenolic content of selected plants from Turkey. *Int. J. Food Sci. Technol.* 2008;43(11):2038-2046.
 28. Klayman DL. *Artemisia annua*: From weed to respectable antimalarial plant. In *Human Medicinal Agents from Plants*; Kinghorn, A. D., Balandrin, M. F., Eds.; American Chemical Society: Washington, DC. 1993. pp. 242-255.
 29. Kumar S, Gupta AK, Pal Y, Dwivedi SK. In-vivo therapeutic efficacy trial with artemisinin derivative, buparvaquone and imidocarb dipropionate against *Babesia equi* infection in donkeys. *J. Vet. Med. Sci.* 2003;65:1171-1177.
 30. Laciari A, Vaca Ruiz M L, Carrizo Flores R, Saad JR. Antibacterial and antioxidant activities of the essential oil of *Artemisia echegarayi* Hieron. (Asteraceae) *Rev. Argent. Microbiol.* 2009;41(4):226-231.
 31. Mazandarani M, Makari S, Bajian GhR, Zarghami Moghaddam P, Abrodi M. Evaluation of phytochemical and antioxidant activity in different parts of *Heracleum gorganicum* Rech in Golestan province of Iran. *Iranian J. plant physiol.* 2011;2:381-386.
 32. Mittler R. Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sci.* 2002;7:405-410.
 33. Mukarram Shah SM, Khan FA, Hassan Shah SM, Chishti KA, Saifur Shah Pirzada SM, Asif Khan M, Farid A. Evaluation of Phytochemicals and Antimicrobial Activity of White and Blue *Capitulum* and Whole Plant of *Silybum Marianum*. *World Appl. Sci. J.* 2011;12:1139-1144.
 34. Muzemil A. PhD thesis, Addis Ababa University. 2008.
 35. Pourmorad F, Hosseini-mehr SJ, Shahabimajd N. Antioxidant activity, phenol and flavonoid contents of some selected Iranian medicinal plants. *Afr. J. Biotechnol.* 2006;5(11):1142-1145.
 36. Qureshi NN, Kuchekar BS, Logade NA, Haleem MA. Antioxidant and Hepatoprotective activity of *Cordia macleodii* leaves. *Saudi. Pharm. J.* 2009;17:317-332.
 37. Rath K, Taxis K, Walz G, Gleiter C H, Li SM, Heide L. Pharmacokinetic study of Artemisinin after oral intake of a traditional preparation of *Artemisia annua* L. (Annul wormwood). *Am. J. Trop. Med. Hyg.* 2004;70:128-132.
 38. Seddik K, Nadjet I, Daoud BAH, Lekhmici A. Antioxidant and antibacterial activities of extracts from *Artemisia herba alba* Asso. leaves and some phenolic Compounds. *J. Med. Plants Res.* 2002;4:1273-1280.
 39. Silva EM, Souza JNS, Rogez H, Rees JF, Larondelle Y. Antioxidant activities and polyphenolic contents of fifteen selected plant species from the Amazonian region. *Food Chem.* 2007;101:1012-1018.
 40. Snow RW, Guerra CA, Noor AM, Myint HY, Hay SI. The global distribution of clinical episodes of *Plasmodium falciparum* malaria. *Nature* 2005;434:214-217.
 41. Surveswaran S, Cai YZ, Corke H, Sun M. Systematic evaluation of natural phenolic antioxidants from 133 Indian medicinal plants. *Food Chem.* 2007;102(3):938-953.
 42. Tan RX, Zheng WF, Tang HQ. Biologically active substances from the genus *Artemisia*. *Planta Med.* 1998;64:295-302.
 43. Tawaha K, Alali FQ, Gharaibeh M, Mohammad M, Elimat TE. Antioxidant activity and total phenolic content of selected Jordanian plant species. *Food Chem.* 2007;104(4):1372-1378.
 44. Tzenkova R, Kamenarska Z, Draganov A, Atanasov A. *Biotechnol. & Biotechnol.* 2010;10:2478/V1013301000306, 1833-1835.
 45. Utzinger J, Chollet J, Jiqing Y, Jinyan M, Tanner M, Shuhua X. Effect of combined treatment with praziquantel and artemether on *Schistosoma mansoni* in experimentally infected animals. *Acta Trop.* 2001;80:9-18.
 46. Verdian-rizi MR, Sadat-Ebrahimi E, Hadjiakhoondi A, Fazeli MR, Pirali Hamedani M. Chemical Composition and Antimicrobial Activity of *Artemisia annua* L. Essential Oil from Iran. *J. Med. Plants*, 2008a;7:58-62.
 47. Verdian-rizi M. Variation in the essential oil composition of *Artemisia annua* L. of different growth stages cultivated in Iran, *African J. Plant Sci.* 2008b;2:16-18.
 48. Willcox M, Falquet J, Ferreira JFS, Gilbert B, Hsu E, Magalhaes PM, Plaizier-Vercammen J, Sharma VP, Wright CW, Yaode W. *Artemisia annua* as an herbal tea for malaria. *African J. Trad. Complem. Altern. Med.* 2007;4:121-123.
 49. Xiao SH, Guo J, Chollet J, Wu JT, Tanner M, Utzinger J. Effect of artemether on *Schistosoma mansoni*: dose-efficacy relationship, and changes in morphology and histopathology. *Chin. J. Parasitol. Dis.*, 2004;22:148-153.
 50. Yang DM, Liew FY. Effects of qinghaosu (artemisinin) and its derivatives on experimental cutaneous leishmaniasis. *Parasitology* 1993;106:107.