

Original Article

Autecology, Ethnopharmacology, Phytochemical, Antioxidant and Antimicrobial Activity of *Thymus carmanicus* Jalas. from Golestan Province in North of Iran

Masoumeh Mazandarani

Department of Biology, Gorgan Branch, Islamic Azad University, Gorgan, Iran

Article History: Received: 26 June 2014/Accepted in revised form: 15 February 2015

© 2013 Iranian Society of Medicinal Plants. All rights reserved

Abstract

This research had been carried out to ecological characters, phytochemical, anti bacterial and antioxidant activity of *Thymus carmanicus* Jalas. in Golestan province. In several field observation, the aerial parts of plant in blooming were collected from Deraznoo Mountain in September 2013 (2700 m). Essential oil was obtained by hydro distillation and analyzed by GC/MS. Phytochemical assay included: TP (total phenol) and TF (Total flavonoid) were determined by spectrophotometrically, antioxidant capacity were obtained by TAC, RP and DPPH methods and the antibacterial by using agar well method and the minimum inhibitory concentration (MIC) assay. Results showed that *T. carmanicus* is the most aromatic mountainous herb, which wild growing in cold dry climate (1800-2750 m) in Golestan province, with annual raining 30.5 mm, annual temperature 17.5 °C, in sandy clay loam soil with pH= 7.5 and $E_c=2.1$ dc and have long been used in traditional medicines of this region as a strong tonic, anti inflammation, carminative, anti virus, anti infection, anti fungus, anti ulcer and sedative. The carvacrol (41.4%), thymol (27.2%) and β -caryophyllene were the major constituents of *T. carmanicus* oil, with total phenols (TP= 81.7 \pm 0.3 GAE/ g), total flavonoids (TF= 34.2 \pm 0.8 QUE/g), which had good antioxidant activity (IC₅₀ = 21.8 \pm 0.1 μ g/ml), especially in DPPH method. The maximum activity of the essential oil was observed against *Candida albicans* (35.8 \pm 0.6 mm), *S. epidermidis* (33.1 \pm 0.4 mm), *B. subtilis* (29.5 \pm 0.2 mm), *S. aureus* (28.3 \pm 0.2 mm), *E. coli* (23.1 \pm 0.4 mm), *Enterococcus faecalis* (17.2 \pm 0.8 mm) and *Kelebsiella pneumoniae* (16.5 \pm 0.2mm) with MIC value in the range of 10 - 132 mg/mL. According to these results, the aerial parts of plant have rich source of terpenoides, flavonoid and phenolic compounds and showed good antioxidant and antibacterial activity, which will be confirmed the traditional uses of *T. carmanicus* Jalas. in these regions as an good anti-inflammatory, sedative and anti infection.

Key words: Antioxidant, Autecology, Essential oil, Ethnopharmacology, Golestan province, TF and TP content, Golestan, *Thymus carmanicus* Jalas

Introduction

Global interest in natural antioxidant, antipathogene, anti inflammation and anti mutagen (Poly phenols, terpenoides and flavonoides) in many wild aromatic plants has recently been increased [1]. So present study was carried on phytochemical sources (Essential oil constituents, TF and TP contents), the evaluate of their

antioxidant and antibacterial activity which were collected from 2700m in North of Iran.

Free radicals cause the oxidation of biomolecules, cell injury and death. In recent decades, so the phytochemical of wild aromatic plants have been great interest as the sources of natural anti oxidant and anti microbial products [2]. The genus *Thymus* (Lamiaceae), native to Southern Europe and Asia, which represented by 215 species in the world and 14 species in Iran flore with four endemic species

*Corresponding author: Islamic Azad University, Gorgan Branch, Gorgan, Iran
E-mail Address: Mazandarani.m@gorganiau.ac.ir

[3,4], which has been used as herbal tea, spicy and medicinal condiments. The chemical composition, *in vitro* antioxidant and antibacterial activities of the essential oils and extracts of several *Thymus* species have recently been reported before, which has been traditionally used for many inflammation and infection such as respiratory tract infections, colds, acute and chronic bronchitis, sinusitis and reduced asthma tonic, carminative, anti-infection, antispasmodic, anti-inflammatory, antitussive and expectorant to treatment of many Iranian ailments [5-8].

In several studies, the essential oil composition, antioxidant, antibacterial and antifungal activity of *Thymus carmanicus* Jalas, *Thymus pubescens* Boiss. & Kotschy ex Celak., *Thymus serpyllum* L., *Thymus kotschyanus* Boiss. & Hohen., *Thymus persicus* (Ronniger ex Rech.f.) Jalas and *Thymus revolutus* Celak. from Iran and Turkey has been reported, respectively [5,19-11], which followed the composition and antioxidant activities of *Thymus caespitius* Brot., *Thymus camphoratus* Hoffmanns. & Link and *Thymus mastichina* oils from Portugal have been reported [12].

T. carmanicus is an endemic Iranian species, which has been used in traditional medicine of Golestan province as sedative, anti-infection and anti-inflammation in treatment of rheumatism and skin disorders [3]. Nejad ebrahimi showed that this species is a rich source of antimicrobial and antioxidant activity [5], so the aim of this research was carried out to the essential oil composition, phytochemical, anti-oxidant and antibacterial activity in Deraznoo Mountain (Golestan province).

Material and Methods

Study area

The present study was carried out in Deraznoo Mountain region (2700m), is located in South west of the Golestan province, in of latitude of 36° 37' 24" to 36° 34' 28" and longitude of 54° 35' 26" to 54° 24' 32" with sandy clay loam soils. Its average height is 1100 to 2750 m from sea levels, with dry cool climate, rainfall of about 305 mm/year and a mean temperature -2/8 °C (January to February) and 17/3 °C (July to August).

Ecological requirements and ethno pharmacology

In many field observation, most ecological requirements and traditional pharmaceutical knowledge about *T. carmanicus*, were obtained from rural healers and then all obtained data from questionnaires were compared with the findings in

vivo and *in vitro* experiments in other similar reports. The aerial parts of *T. carmanicus* were collected at full flowering in September 2014 from its wild habitat in Deraznoo Mountain (2700m). Voucher specimen was deposited at the Herbarium of Research Center of Medicinal Plants (RCMP), in Islamic Azad University of Gorgan branch, Gorgan, Iran.

Isolation of essential oil

The plant samples in blooming stage were dried at room temperature for 3 days. The flowers were hydro distilled in a Clevenger-type apparatus for 2 h according to the method recommended in the British Pharmacopoeia (British Pharmacopoeia, 1988). The oil was dried over anhydrous sodium sulphate and deoxygenated under nitrogen gas and the oil obtained was kept refrigerated and protected from direct light until the analysis time.

Gas chromatography-mass spectrometry (GC-MS) analysis

The essential oil was analyzed by GC-MS. The GC-MS analysis was carried out on a Shimadzu GC-MS (QP5050). The capillary conditions were as follows; carrier gas, helium with a flow rate of 1.7 ml/min; injected 0.1 µL of the essential oil and ionization potential 70 eV. The initial temperature of column was 60 °C (held for 1 min) then heated to 280 °C with a rate of 3 °C/min, then heated to 250 °C and kept constant for 4 min. The same condition of temperature programming was used for n-alkenes mixture to calculate the retention indexes (RI). The identification of each component was studied by mass spectral data, literature and National Institute of Standards and Technology (NIST) computer library. The relative percentage of the oil constituent was calculated.

Plant extract preparation for phytochemical and antioxidant tests

One gram of plant parts with 100 ml (methanol 80%) were extracted by maceration. Extracts were filtered with Whatman No. 1 filter paper. The filtrates obtained from extracts were evaporated into dry rotary evaporator at 40 °C and were stored at 4 °C [13].

Chemicals

2,2'-diphenyl-1-picrylhydrazyl (DPPH) and quercetin (2-(3,4-dihydroxyphenyl)-3,5,7-trihydroxy-4H-chromen-4-one) were purchased from Sigma Chemical Co. (St., Louis, USA). Gallic acid, Folin-Ciocalteu reagent, BHT (butylated hydroxytoluene), BHA (butylated hydroxyanisole)

and methanol were purchased from Merck Co. (Germany).

Determination of Total Phenolic Content

It was determined using the Folin-Ciocalteu Reagent. Total phenolic content was estimated by the Folin Ciocalteu method, based on the procedure suggested by Pourmorad *et al.* [13]. Then 0.5 ml of plant extracts or gallic acid (standard phenolic compound) was mixed with Folin Ciocalteu Reagent (5 ml) and aqueous Na₂CO₃ (4 ml, 1 M). The mixtures were allowed to stand for 15 min and the total phenols were determined by colorimetry at 765 nm. Gallic acid was used as a standard for calibration curve. Total phenol values were expressed in terms of mg equal gallic acid in 1 g powder dry plant.

Determination of Total Flavonoid Content

Total flavonoids content were determined by aluminum chloride method. Extract plants (0.5 ml) were separately mixed with 1.5 ml of solvent, 0.1 ml of 10% aluminum chloride, 0.1 ml of 1 M potassium acetate and 2.8 ml of distilled water. They were kept at room temperature for 30 min; the absorbance of the reaction mixture was measured at 415 nm with a spectrophotometer. Total flavonoid values were expressed in terms of mg equal quercetin in one gram powder dry plant [13].

Antioxidant Activity Tests

Reducing Power assay

This assay is based on Arabshahi-Delouee method. First, The dried extract (12.5–1000 µg) in 1 ml of the corresponding solvent was combined with 2.5 ml of phosphate buffer (0.2 M, pH 6.6) and 2.5 ml of potassium ferricyanide (K₃Fe(CN)₆; 10 g l⁻¹), after the mixture was incubated at 50 °C for 30 min. Then, 2.5 ml of trichloroacetic acid (100 g l⁻¹) were added and the mixture centrifuged at 1650g for 10 min. Then, 2.5 ml of the supernatant solution was mixed with 2.5 ml of distilled water and 0.5 ml of FeCl₃ (1 g l⁻¹), and the samples absorbance was measured at 700 nm [14].

1,1-diphenyl-2-picryl hydrazyl radical scavenging capacity Assay

The ability of the extracts for free radical scavenging was assessed by the method suggested by [14]. Briefly, 1 ml of a 1 mM methanolic solution of DPPH was mixed with 3 ml of extract solution in methanol (containing 12.5-1000 µg of dried extract). The mixture was then vortexed vigorously and left for 30 min at room temperature

in the dark. The absorbance was measured at 517 nm and activity was expressed as percentage DPPH scavenging relative to control using the following equation:

$$\text{DPPH scavenging activity (\%)} = \frac{(\text{A}_{\text{control}} - \text{A}_{\text{sample}})}{\text{A}_{\text{control}}} \times 100$$

Total Antioxidant Capacity

This experimental procedure was adapted from Arabshahi-Delouee method, which is based on the reduction of Mo (VI) to Mo (V) by the sample and observation of a green phosphate/Mo (V) complex at acidic pH. An aliquot of 0.1 ml of sample solution, containing 12.5-1000 µg of dried extract in corresponding solvent, was combined in a tube with 1 ml of reagent solution (0.6 M sulphuric acid, 28 mM sodium phosphate, and 4 mM ammonium molybdate). They were incubated in a thermal block at 95 °C for 90 min. Then we got cold the samples and measured their absorbance at 695 nm. A typical blank solution contained 1 ml of reagent solution and the appropriate volume of the same solvent was used for the sample, and was incubated under the same conditions as the rest of the samples [14].

Bacterial Strains

The bacterial strains were obtained from the Microbiology Laboratory, Golestan University of Medical Sciences. The essential oils of plant aerial part was individually tested against two strains of Gram positive and negative bacteria: (*Shigella dysenteriae* (PTCC1188), *Pseudomonas aeruginosa* (PTCC1430), *E. coli* (PTCC1399), *Staphylococcus aureus* (PTCC1431), *Bacillus cereus* (PTCC1015), *Salmonella typhimurium* (ATCC1596), *Staphylococcus epidermidis* (PTCC1114), *Enterococcus faecalis* (PTCC1393), *Kelebsiella pneumoniae* (PTCC1291) and one fungal isolate: *Candida albicans* (PTCC5027).

Antimicrobial activity

At a first screening, the plant extract were tested against the above mentioned bacteria. Minimal inhibitory concentrations (MICs) were determined by the agar serial dilution method at concentration ranging from 0.93 to 60 µg/mL. Two fold serial dilutions were made from essential oil in molten Mueller Hinton agar (Pronadisa- Madrid) cooled to 45-50 °C in a water bath. The essential oil was dispersed in mixture using dimethyl sulfoxide (DMSO). The amount of 0.01 mL of every bacterial suspension, equivalent to McFarland tube No. 0.5 (10⁸ CFU/mL), inoculated on the agar of every well. The culture plates were then incubated

at 37 °C for 24 h. The MIC was defined as the lowest concentration at which no visible growth was observed (15). The Mueller Hinton agar were contained DMSO without essential oil was used a negative control while Gentamycine was used as positive control.

Statistical analysis

For all assays, data were expressed as means \pm S.E. and differences at $P < 0.05$ were considered statistically significant.

Results

In this study, many field observation showed that the *T. carmanicus* in Deraznoo Mountain is the most Mediteranean wild aromatic perennial chamephyte form plant, which grow to 15-20 cm with small wooden branches, growing spearhead with papil green leaves, which need cool dry weather in North slob and enough light in growing stages. Not heavy soil is suitable soil in Deraznoo Mountain, 57 km far from Gorgan (1800- 2750m) with annual raining 30.5ml, annual temperature 17.5 °C and semi dry cold climate in the sandy clay loam soil (clay=23.8%, silt 32% and sand 53%) with $E_c=2.1$ desizimence and $pH=7.5$.

Ethno pharmacological survey showed that dry Leaves of *T. carmanicus* especially in bloomong, is one of the most wild edible aromatic spicy, which have been used in traditional medicines of Mountainous region of Golestan province as a strong tonic, anti inflammation, carminative, anti virus, anti infection, anti fungus, anti ulcer and sedative to treat of Rheumatism, Ulcer, infection, bronchitis, cold, flu, pain, fever, toothache, backache and digestive infection as a single or in combination with other herbs such as below:

For treat of cold and flu: The tea combination of *T. carmanicus* with, *Satureja mutica*, *Mentha longifolia*, the flower of *Sambucus nigra*, *Adiantum cappilus veneris* and *Cynamum zeylanicum* as a good strong tonic, anti inflammation, carminative, anti infection and warm to treat of col, bronchitis

and flu.

Mouth Ulcer and aftus: the gargle of *Thymus* infusion with consumption the Dusin (the mixed of *Nigella sativa* extract in honey).

Finger fungus infection: the extract of *T. carmanicus* with *Mentha longifolia* L. Sedative and anti inflammation: The massage of the oil of *T. carmanicus*, *Perovskia abrotanoides*, *Stachys inflata* Benth. and *Capsicum annuum* L. anti spasm, anti inflammation and sedative to treat of rheumatic pain, backache, sciatic and gout.

Hair tonic: The infusion of *T. carmanicus*, *Urtica dioica* L., *Adiantum cappilus-veneris* L., *S. inflata* and *Myrtus communis* L. in Olea oil as an hair tonic, anti dandruff and anti inflammation and promote of hair growth.

Anti helmentic: the consumption of *T. carmanicus* powder with *Ferula gummosa* Boiss. and *peganum harmala* L. in daily.

Toothache: *T. carmanicus* powder with *Cuminum cyminum* L..

Earach and ear infection: the 1 drope of *T. carmanicus* oil in Garlic extract in the ears
Phytochemical and anti oxidant results are presented in Table 1 and showed that, the content of total phenols (TP= 81.7 ± 0.3 GAE/g), total flavonoids (TF= 34.2 ± 0.8 QUE/g) in flowering aerial parts of *T. carmanicus*, which had good antioxidant activity with $IC_{50} = 21.8 \pm 0.1$ μ g/ml especially in DPPH method, which had higher content of IC_{50} to free radical scavenging.

Essential oil yield of *T. carmanicus* Jalas. was 2.1%. with 31 components were identified, which represented 91.35% of the total detected constituents. The major constituents of the oil were carvacrol (41.4%), thymol (27.2%) and β -caryophyllene (11.2%), followed by other components were presents in amounts less than 2% (Fig. 1, Table 2).

Table 1 The phytochemical analyses of *Thymus carmanicus* Jalas. In Golestan province

Phytochemical	Deraznooregion (2700m)			Total phenole	Total flavonoid
	Antioxidant activity				
	IC50 in RP	IC50 in TAC	IC50 in DPPH		
	67.4 \pm 0.6	41.3 \pm 0.3	21.8 \pm 0.1	81.7 \pm 0.3	34.2 \pm 0.8

Table 2. Essential oil composition of *Thymus carmanicus* Jas. In Golestan province (2700m).

No.	Compound	RI	RT	%	
1	-		-	3.14	0.3
2	α -thujone	926		4.99	0.1
3	β -pinene	935		5.21	0.2
4	Camphene	949		5.66	0.1
5	Sabinene	965		6.43	0.4
6	β -pinane	976		6.58	0.9
7	α -terpinene	1012		7.61	0.2
8	p-cymene	1014		7.88	0.8
9	Limonene	1018		8.08	0.5
10	β -phelandrene	1021		8.11	0.4
11	1,8-cineole	1024		8.22	0.1
12	β -ocimene	1037		8.58	0.3
13	\square -terpinene	1052		9.05	0.9
14	Terpinene 4-01	1213		9.61	0.1
15	Linalylacetat	1224		10.67	0.09
16	Thymol	1237		11.73	27.2
17	Carvacrol	1282		12.46	41.4
18	Bornyl acetate	1284		13.79	0.2
19	Terpinen 4-01	1288		14.12	0.1
20	α -cucubene	1312		14.06	0.2
21	Benzene	1328		16.57	0.04
22	Naphthalene	1364		17.75	0.2
23	Copaene	1385		18.30	0.5
24	β -burbunene	1404		18.80	0.9
25	β -elemene	1412		19.63	0.5
26	β -caryophyllene	1419		21.34	11.2
27	-Bisabolene β	1499		24.13	0.1
28	γ -cadinene	1509		24.46	0.5
29	Caryophyllene oxide	1585		24.90	0.3
30	Naphthalene	1589		25.06	1.397
31	Isodene	1601		25.25	1.255
%Total	91.35				




Fig. 1 Essential oil analysis chromatogram of *Thymus carmanicus* Jalas. In South west of Golestan province (2700m).

Table 3 Antibacterial activity of aerial parts of *Thymus Carmanicus* Jalas. from Golestan province

Microorganisms	region (2020m)		
	inhibition zone (mm) ±SD	MIC (µg/mL)	Gentamycin
<i>Staphylococcus aureus</i>	28.3±0.2	18.4	20.7
<i>Staphylococcus epidermidis</i>	33.1±0.4	12.3	22.3
<i>Bacillus subtilis</i>	29.5±0.2	19.2	16.5
<i>Enterococcus faecalis</i>	17.2±0.8	73.8	9.6
<i>Escherichia coli</i>	23.1±0.4	41.6	17
<i>Pseudomonas aeruginosa</i>	13.9±0.2	80.5	9
<i>Klebsiella pneumonia</i>	16.5±0.2	94.2	na
<i>Salmonella typhimorium</i>	11.2±0.1	132	na
<i>Shigelladisentria</i>	12.1±0.5	102.3	na
<i>Candida albicans</i>	35.8±0.6	10	13.9

As can be seen in Table 3, essential oil have good potential antimicrobial activity against all of microorganisms tested. The MIC values were in the range of 10-132mg mL⁻¹. The maximum activity of the essential oil was observed against pathogenic fungus: *Candida albicans* (35.8±0.6mm), and two groupe of Gram positive and negative bacteria: *S.epidermidis* (33.1± 0.4 mm), *B.subtilis* (29.5± 0.2mm), *S.aureus* (28.3±0.2mm), *E.coli* (23.1±0.4mm), *Enterococcus faecalis* (17.2±0.8mm) and *Kelebsiella pneumoniae* (16.5±0.2mm), but this oil has poor activity on the

growth of *Shigella dysentria* and *Salmonella tyfimorium*with MIC of 102-132 mg/ mL.

Discussion

Briefly, according to results in Table 1,2,3 , indicated that carvacrol and thymol were the major essential oil terpenoides, with good TP= 81.7±0.3 GAE/ g), total flavonoids content (TF= 34.2±0.8 QUE/g) in flowering aerial parts of *T. carmanicus*, which had good antioxidant activity with IC50 =

21.8±0.1 µg/ml especially in DPPH method, which had higher content of IC50 to free radical scavenging and good anti *Candida* and anti bacterial activity (Table 3).

The natural antioxidants is positively reduces risk of developing infectious, inflammatory and cancer diseases. Therefore, there is a growing interesting to natural antioxidants (Terpenoides, phenolic compounds and flavonoids) search in wild aromatic plants with strong radical scavenging activities [1].

Thymus species the most popular aromatic plants throughout the world, which are commonly has been used as a tonic teas, anti infection, carminative, anti-inflammatory, anti oxidant and antispasmodic to treat of cold, flu, whooping cough, bronchitis, inflammation and rheumatism in Iranian folk medicine and many other [8].

The *S. aureus* and *P. aeruginosa* have been applicable in cases of infectious boils, sores, wounds and are considered as the main pathogens causing hospitalized patients' infections [17] and also, the effectiveness of plant essential oil in growth inhibition against *E. coli*, which can cause the diarrhea and dysentery in humans and animals [18,19].

In many literatures, it is described that the antibacterial activity of *Thymus* species can be dependent on variation of their terpenoids, phenols and flavonoids in essential oil or extract, their habitat and their period growth (5). According to similar reports, the essential oil of several *Thymus* species (*T. revolutus*, *T. pubescens*, *T. serpyllum*, *T. kotschyanus*, *T. persicus*, *T. caespititius*, *T. camphorate*, *T. satureioides*, *T. mastichina* and *T. carmanicus*) from Iran, Turkey and another countries (Morroco) have been reported before and were showed differences in quantities of their essential oil components, but their major components were similar and included: thymol, carvacrol, α -pinene, 1,8-cineole, and caryophyllene [5-10].

In similar reports, essential oils rich in phenolic compounds, such as carvacrol and thymol were reported are widely reported to possess high levels of antimicrobial activity. In fact, other constituents, such as camphen, 1,8-cineol, γ -terpinene and terpinene 4-ol have been considered to display relatively good activity due to their possible synergistic or antagonistic effects, anti inflammation, anti infection and sedative to treat

of Cold, flu, arthritis, rheumatic pain and UTI [5-20].

To confirming of our ethnopharmacological survey, (5) were showed that the carvacrol (58.9-68.9%), p-cymene (3.0-8.9%), c-terpinene (4.3-8.0%), thymol (2.4-6.0%) and borneol (2.3-4.0%) were the major compounds in *T. carmanicus* oils which have strong antibacterial activity against seven Gram-positive and Gram-negative bacteria by Different ranges inhibition zone (15-36 mm) and MIC = 0.5–15.0 mg/ml, thus, they represent an inexpensive source of natural antibacterial, antifungal, antiviral, antiparasitic and antioxidant activities substances that exhibited potential for use in pathogenic systems [5,21-23].

In this study, carvacrol and thymol were the major chemotype of *T. carmanicus* essential oil from Golestan province, but in several before studies, the *Thymus* species has varieties in essential oil composition (geraniol, linalool, γ -terpineol, carvacrol, thymol, trans-thujan-4-ol and terpinen-4-ol) which grown in Iran or in other countries [8]. The thyme specie samples in Turkey, Spanish and other countries (thymol, alpha terpinene, p-cymene, alpha pinene, 1,8-cineole, terpenyl acetate, borneol, linalool, beta-pinene, alphaterpineol, carvacrol and camphor were found to be the main component in the previous report [8]. So comparison between our results and the results of other reports showed differences, probably due to the plant varieties or sites, as well as the time of harvesting.

The before similar antifungal and antibacterial activity exhibited by *Thymus* genus essential oil has been demonstrated by several researchers, which reporting that the essential oils are relatively more active against Gram+ve than Gram-ve bacteria [11,16].

However, It was often reported that Gram-negative bacteria were more resistant to the *Thymus* essential oils [24]. our results (Table 1,3) antibacterial and antioxidant activity of essential oils depends on their chemical composition, which is determined by the genotype and influenced by environmental and agronomic conditions [16]. According to before obtained, the thymol, carvacrol, camphor, alpha-pinene, beta-pinene, borneol and 1,8-cineole are mainly responsible for the antimicrobial and antioxidant activity in many aromatic plants oil, such as: *Thymus species*, *Perovskia abrotanoides*, *Achillea millefolium* L., *F. gummosa* and *Ditrichia* species) [17,27].

To confirming of our phytochemical and ethnopharmacological survey of *T. carmanicus*, other studies showed that the thymol and carvacrol are the main phenolic constituent of *Thymus* species oil have the biological activity, which in medicinal field, their essential oils has been used as a good anti-inflammatory and anti infection to treat acute and chronic bronchitis, acute sinusitis and reduced asthma [6-8].

It has been recognized that secondary metabolites (terpenoids, polyphenols and flavonoids) in aromatic wild plants have antioxidant agents through scavenging on human health [25]. So in general, recently, interest has increased considerably in finding naturally occurring antioxidants and anti pathogen for use in foods or medicinal materials to replace synthetic antioxidants and anti bacterial drugs [26].

According to above, the extract of aerial parts of *T. carmanicus* have better capacity against free radical, especially in DPPH test, which can provided and confirmed of the ethnopharmacological survey of this plant in traditional medicine of this region as anti inflammation, sedative and anti infection.

Conclusion

It can be concluded that the phytochemical and essential oil composition of *T. carmanicus* with major constituents (carvacrol and thymol), possess more antibacterial, antioxidant and anti-inflammatory activity. Considering, it can be believed that the present search together with previous studies provide support of ethnopharmacological results of this study as a tonic, anti infection, antibacterial, anti-inflammatory and antioxidant properties of this plant to treat of many infectious and inflammatory current diseases. It can be used as antibacterial supplement towards the development of new therapeutic agents. Additional *in vivo* studies and clinical trials would be needed to justify and further evaluate the potential of this oil as an antimicrobial agent in topical or clinical applications, and further studies are necessary to evaluate the *in vivo* effects of active compounds of this plant. In addition, investigations confirm that wild aromatic plants used as anti infective as a natural source for novel antibiotics. Our results demonstrate that the oil of *T. carmanicus* could become potentials for controlling certain important Gram positive and

negative bacteria which produces many infectious diseases.

Acknowledgement

The author grateful for the technical support of RCMP (Research Center of Medicinal Plants in Islamic Azad University of Gorgan branch and Niak pharmaceutical factory in Gorgan.

References

1. Mazandarani M, Borhani G, Fathiazad F. Phytochemical analysis, antioxidant activity and ecological requirements of *Capparis spinosa* L. in Golestan and Semnan provinces (North of Iran). *J. Med. Plants By-prod.* 2014; 1: 21-26.
2. Zarghami Moghaddam P, Mazandarani M, Zolfaghari MR. Antibacterial and antioxidant activities of root extract of *Onosma dichroanthum* Boiss. in north of Iran. *Afr. J. Microbiol. Res.* 2012;6:1776-1781.
3. Zargari A. *Medicinal Plants*. Tehran, Iran: Tehran University Press, 1990;4:2842
4. Amin G. *Popular medicinal plants of Iran*. Tehran, Iran: Tehran University of Medical Sciences Press. 2005.
5. Nejad Ebrahimi S, Hadian b, Mirjalili MH, Sonboli C, Yousefzadie D. Essential oil composition and antibacterial activity of *Thymus carmanicus* at different phenological stages. *Food Chem* 110 .2008;927-931.
6. Sfaei-Ghomi J, Meshkat alsadat MH, Shamaei SH, Hashem Nejad M, Hasani A. Chemical characterization of bioactive volatile molecules of four *Thymus* species using nanoscale injection method. *Digest J. Nanomater. Biostruct.* 2009; 4:835-841.
7. Hajimehdipour H, Khanavi M, Zahedi H, Abedi Z, Kalantari Khandani N, Adib N, Pirali Hamedani M. Fingerprint Study of *Thymus* spp. by TLC. 2009;31:19-24.
8. Zenasni L, Bakhy KH, Gaboun F, Mousadak R, Benjouad AA, Al Faiz SH. Essential oil composition and biomass productivity of Moroccan endemic *Thymus satureioides* Coss. & Ball. growing in the Agoundis Valley. *J. Med. Plant Res.* 2014;8: 504-512.
9. Karaman S, Digrak M, Ravid U, Ilcim A. Antibacterial and antifungal activity of the essential oils of *Thymus revolutus* Celak. from Turkey. *J. Ethnopharmacol.* 2001;76:183-186.
10. Rasooli I, Mirmostafa S A. Antibacterial properties of *Thymus pubescens* and *Thymus serpyllum* essential oils. *Fitoterapia.* 2002;73:244-250.
11. Rasooli I, Mirmostafa SA. Bacterial susceptibility to and chemical composition of essential oils from *Thymus kotschyanus* and *Thymus persicus*. *J Agric Food Chem.* 2003;51:2200-2205.
12. Miguel G, Simoes M, Figueiredo AC, Barroso JG, Pedro LG, Carvalho L. Composition and antioxidant

- activities of the essential oils of *Thymus caespitius*, *Thymus camphoratus* and *Thymus mastichina*. Food Chemistry.2004;86:183-188.
13. Pourmorad F, Hosseinimehr SJ, Shahabimajd N . Antioxidant activity, phenol and flavonoid contents of some selected Iranian medicinal plants. Afr. J. Biotechnol. 2006;5:1142-1145.
 14. Arabshahi-Delou S, Urooj A. Antioxidant properties of various solvent extracts of mulberry (*Morus indica* L.) leaves. Food Chem. 2007;102:1233-40.
 15. Hammer KA, Carson CF, Riley TV. Antimicrobial activity of essential oils and other plant extracts. J. Appl. Microbiol.1999; 86:985-990.
 16. Imelouane B, Amhamdi H, Wathelet JP, Ankit M, Khedid K, Elbachiri A. Chemical Composition and Antimicrobial Activity of Essential Oil of Thyme (*Thymus vulgaris*) from Eastern Morocco .Int. J. Agric. Biol. 2009;11:205-208.
 17. Mazandarani, M, Osia N, Ghafourian M. Antioxidant activity and ethnopharmacological survey of *Achillea biebersteinii* Afan. in treatment of Dysmenorrhoea in traditional medicine of Golestan Province, Iran. Int. J. Women's Health Reprod. Sci. 2015; 3:107-110.
 18. Mazandarani M, Mirdeilami S Z, Pesarakli M. Essential oil composition and antibacterial activity of *Achillea millefolium* L. from different regions in North east of Iran. J. Med. Plants Res.2013; 7:1063-1069.
 19. Ogbulie JN, Ogueke CC, Okoli IC, Anyanwu BN. Antibacterial activities and toxicological potentials of crude ethanolic extracts of *Euphorbia hirta*. Afr. J. Biotechnol. 2007;6:1544-1548.
 20. Mojab F, Nickavar B. Composition of the essential oils from three species from Labiatae from Iran. Planta Medica. 2006:72.
 21. Rustaiyan A, Masoudi SH, Monfared A, Kamalinejad M, Lajevardi T, Sedaghat S. Volatile constituents of three *Thymus* species grown wild in Iran.Planta Medica. 2000;66:197-198.
 22. Sefidkon F, Dabiri M, Mirmostafa S A. The essential oil of *Thymus persicus* (Ronniger ex Rech. f.) J alas from Iran. J. Essent. Oil Res. 2002b;14:351-352.
 23. Sadjadi S E, Khatamsaz M . Composition of the essential oil of *Thymus daenensis* Celak. ssp. *lancifolius* (Celak.) J alas. J. Essent. Oil Res. 2003;15:34-35.
 24. Knowles J R, Roller S, Murray DB, Naidu A S. Antimicrobial action of carvacrol at different stages of dual-species biofilm development by *Staphylococcus aureus* and *Salmonella entericsero* var. *typhimurium*. Appl. Environ. Microbiol.2005;71:797-803.
 25. Kessler M, Ubeaud G, Jung L. Antiand pro-oxidant activity of rutin and quercetin derivatives.J. Pharma. Pharmacol. 2003;55:131-142
 26. Zheng W, Wang SY. Antioxidant Activity and Phenolic Compounds in Selected Herbs. J. Agric. Food Chem. 2001;49:5165-5170.
 27. Mozaffarian V. A dictionary of iranian plant names. 1996; N(5558):400.