

Characterization of Non-Terpenoids in *Marrubium crassidens* Boiss. Essential Oil

Sanaz Hamedeyazdan¹, Solmaz Asnaashari², Fatemeh Fathiazad^{3*}

¹ Student's Research Committee, Faculty of Pharmacy, Tabriz University of Medical Sciences, Tabriz, Iran.

² Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz, Iran.

³ Department of Pharmacognosy, Faculty of Pharmacy, Tabriz University of Medical Sciences, Tabriz, Iran.

ARTICLE INFO

Article Type:

Research Article

Article History:

Received: 20 April 2013

Revised: 15 May 2013

Accepted: 18 May 2013

ePublished: 20 August 2013

Keywords:

Marrubium crassidens

Lamiaceae

Oil-poor species

GC-MS

Acetophenone

Tolualdehyde

ABSTRACT

Purpose: *Marrubium crassidens*, a plant belonging to the family Lamiaceae, was studied for its volatile components present in the aerial parts of the plant during the flowering stage. **Methods:** The essential oil of the plant obtained through hydrodistillation of the dried plant material was assessed for its chemical composition by GC/MS and GC-FID analyses. **Results:** Twenty-five compounds were identified, which constituted 94.3% of the total oil composition. The major components were identified as, m-tolualdehyde (23.3%), acetophenone (15.8%), nonacosane (13.1%), docosane (7.2%), o-tolualdehyde (4.1%), β -caryophyllene (3.8%) and caryophyllene oxide (3.4%). Non-terpenoids with 75.7% were the most abundant components of the essential oil. **Conclusion:** Overall, *M. crassidens* essential oil revealed to include rather higher proportions of non-terpenoid compounds compared with other species of genus *Marrubium*.

Introduction

The family Lamiaceae with about 220 genera and almost 4000 species worldwide has always been the leader sources for culinary, vegetable and medicinal plants.^{1,2} In this regard, *Marrubium* a genus of about 40 species of flowering plants in this family has been holding a place of value in different cultures and herbal medicine with some known healing attributes.^{3,4} Cytotoxicity, immunomodulating, vasorelaxant, antispasmodic, hypolipidemic, hypoglycemic, and analgesic activities are some of the famous properties for species of genus *Marrubium* that have been reported by several studies.⁵⁻¹³ Although *Marrubium* species are mainly recognized for their non-volatile compounds, like diterpenes, polyphenols, steroids, phenylpropanoid glycosides and flavonoids, they happen to produce small amounts of essential oil and thus they are called as oil-poor species.¹⁴⁻¹⁶ Generally, essential oils from *Marrubium* sp. consist mostly of sesquiterpenes and studies concerning their antimicrobial and antioxidant activities are sparse.¹⁷⁻²³ Nevertheless in our previous study on *M. persicum*, it was described substantially as a species distinct in chemical composition of the essential oil rich in non-terpenoid compounds from other species of *Marrubium*. According to the

chemical diversity of essential oils within a genus indicating both quantitative and qualitative diversity in the composition of the essential oils, searching for the variety of naturally occurring volatile constituents in the essential oil of a plant appears to be of value. So as, based on flora Iranica, *M. crassidens*, endemic to the countries; Armenia, Azerbaijan, Turkey and Iran²⁴ was selected to be further analyzed for its volatile constituents from the aerial parts of the plant that is grown in Azarbaijan province in Iran.

Materials and Methods

Plant material

Aerial parts of *Marrubium crassidens* were collected during the flowering stage from Chichaklou in East Azarbaijan province, in June 2011. A voucher specimen of the plant (Tbz-Fph-719) representing this collection has been deposited at the Herbarium of the Faculty of Pharmacy, Tabriz University of Medical science, Iran.

Essential oil extraction

Air-dried plant material of the aerial parts of *M. crassidens* was subjected to hydrodistillation using a Clevenger-type apparatus (Clevenger, 1928). Since

*Corresponding author: Fatemeh Fathiazad, Department of Pharmacognosy, Faculty of Pharmacy, Tabriz University of Medical Sciences, Tabriz, Iran. Tel: +98 411 3372253, Fax: +98 411 3344798, Email: fathiazad@tbzmed.ac.ir

the oil content was low in quantity, the distillation time was prolonged (3h) and xylene was used as an absorbing medium. The obtained essential oil was stored in sealed glass vial at 4-5 °C prior to analysis.

Gas Chromatography-Mass Spectrometry (GC-MS)

The essential oil was analyzed by GC-MS using a Shimadzu GC-MS-QP 5050A gas chromatograph fitted with a DB1 (methyl phenyl sylonane, 60 m x 0.25 mm i.d., 0.25 µm film thickness) capillary column. The GC was set at the following conditions with helium as the carrier gas; flow rate of 1.3 mL/min; linear velocity: 29.6 cm/s; Split ratio, 1:29; column temperature, 2 min in 60 °C, 50-260 °C at 3 °C/min; injector temperature, 240 °C, and 1 µL of volume injection of the essential oil. The MS operating parameters were as follows: ionization potential, 70 eV; ion source temperature; 270 °C; quadrupole 100 °C, Solvent delay 2 min, scan speed 2000 amu/s and scan range 30-600 amu, EV voltage 3000 volts.

Identification of the compounds

The identification of compounds was based on direct comparison of the retention indices and mass spectral data with those for the standards and by computer matching with the Wiley 229, Nist 107, Nist 21 Library, as well as by comparing the fragmentation patterns of the mass spectra with those reported in the literature.²⁵ For quantification purpose, relative area percentages were obtained by FID without the use of correction factors, where the FID detector condition was set on a duplicate of the same column applying the same operational conditions.

Results and Discussions

The GC-MS analysis of the small amount of essential oil obtained through hydrodistillation led to the identification of 25 different components, representing 94.3% of the total oil constituents. All the identified compounds were arranged in order of their elution from the DB1-MS column and the retention indices whereas their percentage compositions were summarized in Table 1.

Table 1. Chemical constituent of the essential oil from aerial parts of *M. crassidens*.

No.	^a RI	Compounds	Area (%)
1	906	n-Nonane	2.1
2	933	Benzaldehyde	0.5
3	936	α-Pinene	1.8
4	966	1-Octen-3-ol	2.8
5	1004	Decane	0.3
6	1025	Limonene	0.6
7	1037	Acetophenone	15.8
8	1041	m-Tolualdehyde	23.3
9	1053	o-Tolualdehyde	4.1
10	1381	α-Cubebene	0.7
11	1424	β-Caryophyllene	3.8
12	1450	β-Farnesene	1.5
13	1457	α-Humulene	0.4
14	1482	Germacrene D	2.6
15	1488	β-Selinene	0.6
16	1497	Bicyclgermacrene	1.9
17	1520	δ-Cadinene	0.4
18	1571	Spathulenol	0.9
19	1578	Caryophyllene oxide	3.4
20	1834	Hexahydrofarnesyl acetone	1.9
21	1945	Eicosanoic acid	0.7
22	1989	Docosane	7.2
23	2592	Hexacosane ^b	2.0
24	2793	Octacosane ^b	1.9
25	2892	Nonacosane ^b	13.1
Total			94.3
Non-terpenoids			75.7
Monoterpene hydrocarbons			2.4
Sesquiterpene hydrocarbones			11.9
Oxygenated sesquiterpenes			4.3

^a RI is the Retention Index relative to C8–C24 n-alkanes on the DB-1 column.
^b This compound was compared with an authentic sample.

It was established that the essential oil of *M. crassidens* was a complex mixture of mostly non-terpenoids

(75.7%), sesquiterpene hydrocarbons (11.9%), oxygenated sesquiterpenes (4.3%) and monoterpene

hydrocarbons (2.4%). In spite of the absence of oxygenated monoterpenes, something to be expected in study of *Marrubium* species essential oil, our sample was characterized by the presence of alkanes, alkanolic acid derivatives and ketones in great amounts. As regards, m-tolualdehyde with 23.3%, and acetophenone with 15.8% were the two basic constituents present among the non-isoprenoid compounds of the essential oil. Moreover, β -caryophyllene (3.8%), germacrene D (2.6%), and bicyclogermacrene (1.9%) were the leader sesquiterpene hydrocarbons that were present in the *M. crassidens* essential oil. The only identified two monoterpenes in the essential oil were α -pinene, and limonene with the relative percentages of 1.8 and 0.6%, respectively. Besides, caryophyllene oxide with 3.4% and spathulenol with 0.9% were the only components of the oxygenated sesquiterpene compound grouping.

Having reviewed previously reports of the essential oils from genus *Marrubium*, it was established that *Marrubium* species possess rather low amounts of aliphatic and non-terpenoid fractions in essential oils, in contrary to our findings.^{21-23,26} Owing to the fact that the production of terpenoid and non-isoprenoid compounds diverges early in the pathway of anabolic plant secondary compound synthesis,^{27,28} necessitates fundamental studies in advance especially in practical applications of the essential oils in fragrance and flavor industries, as well as in the pharmaceutical and chemical industries. In order to distinguish between the differences observed in the essential oil constituents of *M. crassidens* with other species of genus *Marrubium* additional studies could be of advantageous in determining the intrinsic (genetic, growth stage, etc.) and extrinsic (climatic, seasonal, environmental distillation processes, etc.) conditions affecting the relative biosynthesis pathways of the essential oils.

Conclusion

On the whole, *M. crassidens* in agreement with previous reports for other *Marrubium* species seems to be poor in essential oil content while it shares some similarities and also differences in constituents of the essential oil. Meanwhile, presence of relatively higher amounts of non-terpenoids; acetophenone with different isomers of tolualdehyde in the *M. crassidens* essential oil has been reported for the first time in this study for this plant.

Acknowledgments

The authors would like to thank Drug Applied Research Center of Tabriz University of Medical Sciences for recording the mass spectra. Financial support of this work by the Research Vice-Chancellor of Tabriz University of Medical Sciences is faithfully acknowledged. This article was written based on a data set of PhD. thesis, registered in Tabriz University of Medical Sciences (5/4/6651- NO. 71).

Conflict of Interest

The authors report no conflict of interest in this study.

References

1. Hadley SK, Petry JJ. Medicinal herbs: a primer for primary care. *Hosp Pract* (1995) 1999;34(6):105-6, 109-12, 115-6 passim.
2. Naghibi F, Mosaddegh M, Mohammadi Motamed S, Ghorbani A. Labiatae Family in folk Medicine in Iran: from Ethnobotany to Pharmacology. *Iran J Pharm Res* 2005;2:63-79.
3. Paula De Oliveira A, Santin JR, Lemos M, Klein Junior LC, Couto AG, Meyre Da Silva Bittencourt C, et al. Gastroprotective activity of methanol extract and marrubiin obtained from leaves of *Marrubium vulgare* L. (Lamiaceae). *J Pharm Pharmacol* 2011;63(9):1230-7.
4. Meyre-Silva C, Cechinel-Filho V. A review of the chemical and pharmacological aspects of the genus *marrubium*. *Curr Pharm Des* 2010;16(31):3503-18.
5. Boudjelal A, Henchiri C, Siracusa L, Sari M, Ruberto G. Compositional analysis and in vivo anti-diabetic activity of wild Algerian *Marrubium vulgare* L. infusion. *Fitoterapia* 2012;83(2):286-92.
6. Hamedeyazdan S, Fathiazad F, Sharifi S, Nazemiyeh H. Antiproliferative activity of *Marrubium persicum* extract in the MCF-7 human breast cancer cell line. *Asian Pac J Cancer Prev* 2012;13(11):5843-8.
7. Zaabat N, Hay AE, Michalet S, Darbour N, Bayet C, Skandrani I, et al. Antioxidant and antigenotoxic properties of compounds isolated from *Marrubium deserti* de Noe. *Food Chem Toxicol* 2011;49(12):3328-35.
8. Meyre-Silva C, Yunes RA, Schlemper V, Campos-Buzzi F, Cechinel-Filho V. Analgesic potential of marrubiin derivatives, a bioactive diterpene present in *Marrubium vulgare* (Lamiaceae). *Farmaco* 2005;60(4):321-6.
9. Rigano D, Aviello G, Bruno M, Formisano C, Rosselli S, Capasso R, et al. Antispasmodic effects and structure-activity relationships of labdane diterpenoids from *Marrubium globosum* ssp. *libanoticum*. *J Nat Prod* 2009;72(8):1477-81.
10. El Bardai S, Lyoussi B, Wibo M, Morel N. Pharmacological evidence of hypotensive activity of *Marrubium vulgare* and *Foeniculum vulgare* in spontaneously hypertensive rat. *Clin Exp Hypertens* 2001;23(4):329-43.
11. Karioti A, Skopeliti M, Tsitsilonis O, Heilmann J, Skaltsa H. Cytotoxicity and immunomodulating characteristics of labdane diterpenes from *Marrubium cylleneum* and *Marrubium velutinum*. *Phytochemistry* 2007;68(11):1587-94.
12. Berrougui H, Isabelle M, Cherki M, Khalil A. *Marrubium vulgare* extract inhibits human-LDL oxidation and enhances HDL-mediated cholesterol efflux in THP-1 macrophage. *Life Sci* 2006;80(2):105-12.

13. Argyropoulou A, Samara P, Tsitsilonis O, Skaltsa H. Polar constituents of *Marrubium thessalum* Boiss. & Heldr. (Lamiaceae) and their cytotoxic/cytostatic activity. *Phytother Res* 2012;26(12):1800-6.
14. Calis I, Hosny M, Khalifa T, Ruedi P. Phenylpropanoid glycosides from *Marrubium alysson*. *Phytochemistry* 1992;31(10):3624-6.
15. Karioti A, Skaltsa H, Heilmann J, Sticher O. Acylated flavonoid and phenylethanoid glycosides from *Marrubium velutinum*. *Phytochemistry* 2003;64(2):655-60.
16. Argyropoulou C, Skaltsa H. Identification of essential oil components of *Marrubium thessalum* Boiss. & Heldr., growing wild in Greece. *Nat Prod Res* 2012;26(7):593-9.
17. Semnani KM, Saeedi M, Babanezhad E. The essential oil composition of *Marrubium vulgare* L. from Iran. *J Essent Oil Res* 2008;20(6):488-90.
18. Tajbakhsh M, Khalilzadeh A, Rineh A, Balou J. Essential oils of *Marrubium anisodon* C. Koch and *Marrubium propinquum* Fisch. et C.A. Mey., growing wild in Iran. *J Essent Oil Res* 2008;161-2.
19. Petrovic S, Pavlovic M, Maksimovic Z, Milenkovic M, Couladis M, Tzakouc O, et al. Composition and antimicrobial activity of *Marrubium incanum* Desr. (Lamiaceae) essential oil. *Nat Prod Commun* 2009;4(3):431-4.
20. Laouer H, Yabrir B, Djeridane A, Yousfi M, Beldovini N, Lamamra M. Composition, antioxidant and antimicrobial activities of the essential oil of *Marrubium deserti*. *Nat Prod Commun* 2009;4(8):1133-8.
21. Nagy M, Svajdlenka E. Comparison of essential oils from *Marrubium vulgare* L. and *M. peregrinum* L. *J Essent Oil Res* 1998;10:585-7.
22. Demirci B, Baser KGC, Kirimer N. Composition of the essential oil of *Marrubium bourgaei* ssp. *caricum* P.H. Davis. *J Essent Oil Res* 2004;16:133-4.
23. Javidnia K, Miri R, Soltani M, Khosravi AR. Constituents of the Essential Oil of *Marrubium astracanicum* Jacq. from Iran. *J Essent Oil Res* 2007;19:559-61.
24. Mozaffarian V. A Dictionary of Iranian Plant Names. Tehran, Iran: Farhang Moaser Publication; 2004.
25. Adams RP. Identification of Essential oil Components by Gas Chromatography/ Quadrupole Mass Spectroscopy. Carol Stream, IL: Allured Publ Corp; 2004.
26. Zarai Z, Kadri A, Ben Chobba I, Ben Mansour R, Bekir A, Mejdoub H, et al. The in-vitro evaluation of antibacterial, antifungal and cytotoxic properties of *Marrubium vulgare* L. essential oil grown in Tunisia. *Lipids Health Dis* 2011;10:161.
27. Ganjewala D, Luthra R. Essential oil biosynthesis and regulation in the genus *Cymbopogon*. *Nat Prod Commun* 2010;5(1):163-72.
28. Lawrence BM. Essential oils. Carol Strem, IL: Allured Publishing Corp; 1992.