

Research Article



Dual Preconditioning: A Novel Strategy to Withstand Mesenchymal Stem Cells against Harsh Microenvironments

Hamed Bashiri^{1,2} , Fatemeh Amiri² , Ali Hosseini² , Masoud Hamidi³ , Amaneh Mohammadi Roushandeh³ , Yoshikazu Kuwahara⁴ , Mohammad Ali Jalili² , Mehryar Habibi Roudkenar^{5*} 

¹ Department of Medical Laboratory Sciences, Faculty of Paramedical, Kurdistan University of Medical Sciences, Sanandaj, Iran.

² Department of Medical Laboratory Sciences, School of Paramedicine, Hamadan University of Medical Sciences, Hamadan, Islamic Republic of Iran.

³ Medical Biotechnology Research Center, Paramedicine Faculty, Guilan University of Medical Sciences, Rasht, Iran.

⁴ Division of Radiation Biology and Medicine, Faculty of Medicine, Tohoku Medical and Pharmaceutical University, Sendai, Japan.

⁵ Cardiovascular Disease Research Center, Department of Cardiology, Heshmat Hospital, School of Medicine, Guilan University of Medical Sciences, Rasht, Iran.

Article info

Article History:

Received: 16 January 2018
Revised: 21 May 2018
Accepted: 19 July 2018
ePublished: 29 August 2018

Keywords:

- Hydrogen peroxide
- Mesenchymal Stem Cells
- Serum deprivation
- Simultaneous preconditioning
- Survival
- Harsh microenvironments

Abstract

Purpose: Poor survival rate of mesenchymal stem cells (MSCs) following their transplantation is one of the major challenges in their therapeutic application. Therefore, it is necessary to augment the viability of the MSCs in order to improve their therapeutic efficacy. Several strategies have been used to overcome this problem. Preconditioning of MSCs with oxidative stresses has gained a lot of attention. Therefore, in the present study, we investigated the effects of simultaneous preconditioning of MSCs with hydrogen peroxide and serum deprivation stresses on their survival and resistance to stressful conditions.

Methods: MSCs were isolated from human umbilical cord blood. To perform simultaneous preconditioning, the cells were cultured in DMEM medium containing 1, 2.5 and 5 percent FBS and different concentrations of H₂O₂ (5, 10, 15, 20, 25, 30, 35, 40, 50, 60, 80 and 100 μM) for 24 hrs. Then, the cells were cultured in recovery culture medium. Finally, one group of the cells was exposed to a lethal concentration of H₂O₂ (300μM), and the other cells were cultivated in FBS free DMEM medium as the lethal situation. In addition, the percentage of apoptotic cells was analyzed using Caspase 3 assay kit.

Results: Simultaneous preconditioning of the MSCs with 15μM H₂O₂ plus serum deprivation, 2.5% FBS, significantly increased the resistance of the cells to the toxicity induced following their cultivation in FBS free DMEM medium. It exerted the protective effect on the cells after treating with the lethal dose of H₂O₂ as well.

Conclusion: Simultaneous preconditioning of MSCs with oxidative and serum deprivation stresses enhances their survival against harsh conditions, which might increase the viability and stability of the MSCs following their transplantation.

Introduction

Recently, it has been clear that the mesenchymal stem cells (MSCs) are promising cell source for the treatment of a variety of human diseases^{1,2} including severe aplastic anemia^{3,4} acute graft- versus-host disease,⁵ cardiovascular diseases, acute liver failure⁶ and kidney injuries.^{7,8} Multi-lineage differentiation potential, immune modulatory properties and ability to localize specifically to injured sites have made MSCs as an appropriate alternative.^{2,8} According to animal and clinical studies, MSC transplantation can restore cardiac function probably by myogenesis and angiogenesis after myocardial infarction.⁹⁻¹¹ Although there has been an increasing interest in using these cells in cell therapy but one of the main obstacles of their application is poor survival after transplantation. It has been shown that a majority of

implanted cells die within few days after transplantation.^{12,13} Endogenous and environmental factors¹⁴⁻¹⁸ including inflammatory responses, lack of nutritional factors, hypoxia, reactive oxygen species (ROS) such as superoxide anion (O₂⁻), hydroxyl radical (OH[·]), and hydrogen peroxide (H₂O₂), induce apoptosis and higher cell death either in vitro or in vivo MSCs microenvironments specially in ischemic heart medium.¹⁶ Hence, it is necessary to augment the viability of the MSCs in order to improve their efficacy. Moreover, several strategies including genetic manipulation and injection of growth factors and drugs have been employed to overcome this problem.¹⁹ In this regard, strategies which increase the survival of stem cells have gained significant attention. This indicates a need to understand

*Corresponding author: Mehryar Habibi Roudkenar, Tel: +98 1342536262 Fax: +98 1342565051, Email: roudkenar@gums.ac.ir

©2018 The Authors. This is an Open Access article distributed under the terms of the Creative Commons Attribution (CC BY), which permits unrestricted use, distribution, and reproduction in any medium, as long as the original authors and source are cited. No permission is required from the authors or the publishers.

underlying mechanisms of the decreased viability of the stem cells in the stress conditions.

Recently, preconditioning of MSCs with non-ischemic stresses such as stretch some chemicals, hypoxia, reactive oxygen radicals and oxidative stresses have been considered in the literature.^{20,21} Previous studies have demonstrated that ischemic preconditioning (IPC) and oxidative preconditioning have protective effects on different kinds of cells and stem cells through pathologic condition and could be helpful to treat related diseases.²⁰⁻²⁶ Despite the importance of the effect of preconditioning, there have been no controlled studies on the harsh microenvironment of the injured tissues.

In order to potentiate MSCs against multiple threatening factors, it is necessary to expose them to several stresses conditions, which ensures their resistance against many inappropriate factors.^{21,23,25,26} Therefore, in the present study, the protective effects of co-preconditioning of MSCs with various concentrations of H₂O₂ and low doses of FBS have been evaluated. The main purpose of this study was to investigate the protective effects of simultaneous preconditioning on cell survival and prepare them to face with harsh microenvironments after transplantation.

Materials and Methods

Isolation and expansion of MSCs

The umbilical cord blood sample was collected from women who underwent caesarian section, with informed consent and mixed with citrate phosphate dextrose (CPD) anticoagulant. The sample was diluted in phosphate-buffered saline/ Ethylene diamine tetra acetic acid (PBS/EDTA) at a ratio of 3:1. Then, mononuclear cells were separated by Ficoll-Hypaque density gradient centrifugation (at 435g for 20 min), and seeded cell culture flasks containing low glucose Dulbecco's modified eagle medium (DMEM), antibiotics (0.01% penicillin/streptomycin) and 10% fetal bovine serum (FBS) (All of materials purchased from Gibco, Germany). The cells were incubated in the presence of 95% air and 5% CO₂ at 37°C for 48hrs. Then, the non-adhered erythroid progenitor cells were removed by changing the medium. Medium refreshment was performed two times per week for 14 days prior to further studies. At 80% confluence, cells were detached with 0.25% trypsin-EDTA (Sigma Aldrich, Germany), washed with PBS and re-plated under the same culture conditions. To confirm the identity of the cultured umbilical cord blood MSCs (UCB-MSCs) morphologic features of the cells were evaluated by an inverted microscope and after that, the presence of specific surface markers, CD73, CD90 and CD105²⁷ of MSCs were analyzed by flow cytometry device (Partc PASIII, Germany). The 4th passages of UCB-MSCs were used in the present study.

Preconditioning of MSCs

Preconditioning with H₂O₂

To investigate the possible cytoprotective effect of preconditioning with different concentrations of H₂O₂ on

UCB-MSCs, 10000 cells were cultured in 96-well plate and then incubated with different concentrations of H₂O₂ (5, 10, 15, 20, 25, 30, 35, 40, 50, 60, 80 and 100 μM) for 24 hrs.²⁵ followed by a recovery period of 12 hrs in usual growth medium. The experiment was examined in triplicate. After the recovery period, the culture medium was changed and preconditioned cells were exposed to 300 μM H₂O₂ as the lethal dose for 24 hrs.²⁵

Preconditioning with serum deprivation (SD)

10000 UCB-MSCs were seeded in 96-well plates and preconditioned with 1, 2.5 and 5% FBS for 24 hrs followed by a recovery period of 12hrs in a usual growth medium. All of the concentrations were examined in triplicates. After the recovery period, the medium was decanted and the preconditioned cells were cultured in FBS free low glucose-DMEM for 24hrs as a lethal condition.

Simultaneous preconditioning of MSCs with H₂O₂ and SD

In order to induce more stress condition to the cells and make them stronger against various toxic factors in the harsh microenvironment, the UCB-MSCs were cultured in oxidative stress and SD conditions. Briefly, the UCB-MSCs were cultured in 96-well plates containing low glucose-DMEM supplemented with 2.5% FBS, and simultaneously the cells were treated with 15 μM H₂O₂ solution for 24 hrs, followed by 12 hrs of recovery in a usual growth medium. After the recovery period, the cells separately were cultivated in serum-free condition or lethal dose of 300μM H₂O₂ for 24 hrs.

Evaluation of cell viability

Cell survival of different preconditioned UCB-MSCs and control was assessed by colorimetric method using water-soluble tetrazolium salt-1 (WST-1) as described previously.²¹ Briefly, after treating of the cells with different stress conditions, the WST-1 reagent (Sigma, Germany) was added to culture media at a ratio of 1:10 and mixed gently. The plates were transferred to Co₂ incubator at 37°C for 4 hrs. Using a microplate reader (BioTek, Germany), the optical density (OD) of each well was evaluated at 450 nm.

Assessment of apoptosis

Caspase 3 activity was carried out using Caspase 3 assay kit (Sigma, Germany). According to manufacture protocol, cell lysate of different experimental and control groups was prepared. 5μl of the cell lysate and 5μl of Caspase 3 positive control were added to each well. Then, 10 μl of Caspase 3 substrate, and Caspase 3 inhibitor were added to the wells and incubated for 70-90 minute and finally, the absorbance was read at 405 nm with a microplate reader.

Statistical analysis

At each data point, the mean and standard deviation (SD) were calculated and statistically analyzed using Student's t-test. p<0.05 was considered significant.

Results

UCB-MSCs were fibroblast-like and expressed general surface markers of MSCs

Examination of the cells under an inverted microscope revealed that the isolated UCB-MSCs have fibroblast-like morphology and plastic adherent property (Figure 1A). Flowcytometry analysis was also employed to confirm the identity of UCB-MSCs. The results showed that the isolated cells were positive for CD29, CD105, and CD73 and negative for CD34 and CD45 (Figure1B).




Figure 1. Characterization of umbilical cord blood mesenchymal stem cells (UCB-MSCs) A. Adherent UCB-MSCs displayed fibroblastoid morphology. B. Immunophenotype of isolated MSCs

H₂O₂-preconditioning enhanced cell survival of UCB-MSCs and decreased their apoptosis rate

As described in above, UCB-MSCs were treated by different concentration of H₂O₂. However, only preconditioning with 5, 10, 15 and 20 μM of H₂O₂ significantly increased the survival of UCB-MSCs in comparison with non-preconditioned control groups (without any treatment) following their exposure to lethal concentration (300μM) of H₂O₂ (p< 0.001 for 15 μM of H₂O₂, p< 0.01 for 5 and 10 μM of H₂O₂, and p< 0.05 for 20 μM). However, preconditioning with higher concentrations of H₂O₂ did not protect these cells from oxidative stress-induced cell death (Figure 2A). The results of the Caspase 3 level analysis were presented in Figure 2B. As is shown in Figure 2B, when 5, 10 and 15 μM H₂O₂-preconditioned-UCB-MSCs were exposed to lethal concentration of H₂O₂ (300μM) for 24 hrs obviously exhibited less Caspase 3 level, an index of apoptotic rate, in comparison with the non-preconditioned control cells, (p<0.001 for 15 μM of H₂O₂, p< 0.01 for 5 and 10 μM of H₂O₂). According to these data, 15μM H₂O₂ was considered as optimal dose for further studies.




Figure 2. Evaluation of cell viability and apoptosis of H₂O₂-preconditioned UCB-MSCs following oxidative stress. A. WST-1 assay for analysis of cell viability. Higher survival was detected in the preconditioned cells in comparison with normal non-preconditioned cells (**p< 0.01, ***p< 0.001, *p< 0.05). B. Assessment of Caspase 3 activity level as apoptosis indicator. H₂O₂-preconditioned UCB-MSCs exhibited lower Caspase 3 activity level in comparison with the related control groups (**p< 0.01, ***p< 0.001). Data represents Mean ±SD of two independent experiments

SD preconditioning protected UCB-MSCs from cell death and apoptosis in serum-free medium

As is shown in Figure 3A and 3B, culturing of UCB-MSCs in medium containing 2.5 % FBS following culturing under serum-free medium, as harsh stress-inducing condition, not only led to higher viability percentage in UCB-MSCs but also led to lower Caspase 3 activity level (p < 0.001) in comparison with the non-preconditioning group. P-value was 0.01 and 0.05 for the cells that were exposed to 1% and 5% FBS in comparison with control (non-preconditioned cells) respectively. 2.5 % of FBS was set up as optimized serum deprivation condition and was considered as optimal dose for further studies.

Simultaneous preconditioning of UCB-MSCs with H₂O₂ and serum deprivation conferred more resistance to these valuable cells against harsh condition

For induction of more stresses to UCB-MSCs, they were preconditioned with both 15μM H₂O₂ and 2.5% FBS. Then, they were cultured in 300μM lethal H₂O₂ or serum-free conditions. It is worthy to note that the viability of simultaneously preconditioned cells was higher than control and the groups receiving the single preconditioning modality (p<0.05). In other words according to the Figure 4, simultaneous preconditioning

makes the UCB-MSCs more resistant to the harsh microenvironments.




Figure 3. Assessment of cell viability and apoptosis rate of SD-preconditioned UCB-MSCs under serum free condition. A. WST-1 assay. (***) $p < 0.001$, ** $p < 0.01$, * $p < 0.05$). B. Assessment of Caspase 3 activity level. (***) $p < 0.001$, ** $p < 0.01$, * $p < 0.05$). subjected cells in compared to controls. Data represents Mean \pm SD of two independent experiments




Figure 4. Viability of simultaneous (SD/Oxidative) - preconditioned UCB-MSCs after exposure to lethal oxidative stress and serum free condition. SD/Oxidative group was compared with those groups that were exposed to only SD or oxidative stress (* $p < 0.05$). Data represents Mean \pm SD of two independent experiments

Discussion

Currently, the applications of the MSCs for cell therapy and tissue engineering purposes are under the focus of the investigation.^{1,2} However, many problems were developed in MSCs transplantation which hindered the prosperity of them in cell therapy.^{8,28} Inappropriate environment of tissue containing high amounts of free oxygen radicals, inflammatory cytokines and lack of

nutrition and blood supply are some factors which can threaten the survival of transplanted cells.²⁹ Toma et al reported that less than only 1% of the cells survived few days after their transplantation into the heart of severe combined immune deficiency disorder mice.¹² To solve this problem, several strategies like genetic modification of MSCs^{19,30-32} and injection of growth factors^{33,34} have been suggested in some literature. However, because of the risk of tumor development or low efficacy of these strategies,¹⁹ developing new strategies is necessary. The beneficial effects of preconditioning of the MSCs were first suggested by Murry et al, in 1986.³⁵ Rosova et al reported that hypoxic preconditioning of MSCs led to the improvement of their healing potential.³⁶ Also, Tang et al reported that preconditioning with H₂O₂ resulted in a reduction of apoptosis.³⁴ In their study PC12 cell line, a rat cell line, was used. This group treated the cells with 0, 5, 10, 20, and 30 $\mu\text{mol L}^{-1}$ of H₂O₂. After recovery by 24 hrs cultivation in normal media, these cells were exposed to 20, 30, 50, 100 $\mu\text{mol L}^{-1}$ of H₂O₂ for another 24 hrs. They evaluated apoptosis rate by expression of Bcl2 level, mitochondrial membrane potential, and intracellular ROS³⁴ Considering that the majority of engrafted MSCs may die within the first few days of transplantation, the transient cytoprotective effects of simultaneous preconditioning could be sufficient to protect transplanted cells during the first critical period after transplantation. Enhanced survival of implanted cells might reduce the required number of transplanted cells, which in turn, fewer stem cells may differentiate better.

Gargioli and colleagues exposed mouse perivascular myogenic progenitors to severe oxidative stress (200 and 400 μM of H₂O₂) and studied their survival, self-renewal and myogenic differentiation capacity. They reported that the H₂O₂-treated cells showed higher survival, proliferation and engrafted rate.³⁷

The findings of the present study supported the hypothesis that preconditioning of MSCs might increase their survival and might increase their therapeutic potency for transplantation. Our results indicated that applying of an easy and non-expensive method can protect MSCs against the induced apoptosis by lethal stress conditions.

The molecular mechanism underlying the protective effects of different preconditioning methods on MSCs is not fully understood yet. However, lowering of apoptotic cells by regulation of some anti-apoptotic protein,³⁴ modulation of some important growth factors/cytokines/chemokines and their specific receptors and activation of signaling pathways such as Notch1/Wnt1¹³⁸⁻⁴⁰ are the possible mechanisms.

Pretreatment with sub-lethal oxidative stress induces expression CXCR4 on the MSCs and enhances their survival as well as decreases apoptosis in these valuable cells.²⁵ In the present investigation, we provided new evidence that simultaneous preconditioning of MSCs enhanced their survival rate. However, further and comprehensive studies should be performed to address the limitations of this study including the mechanisms underlying simultaneous preconditioning exert

cytoprotective effects. Moreover, evaluation of apoptosis by other well-known techniques could be the subject of future studies. Finally, to expand the results of current study, *in vivo* study should be conducted to confirm the therapeutic potentialities in animal models.

Conclusion

In conclusion, our study demonstrated that simultaneously preconditioning of MSCs with two different stresses enhanced protective effects. This would be a safe and versatile strategy to increase MCSs-based cell therapy in clinical applications.

Acknowledgments

This study was supported by the Blood Transfusion Research Center, High Institute for Research and Education in Transfusion Medicine, Republic of Iran, grant number, 1413. The authors are thankful to Dr. Mahshid Mohammadipour for her technical assist.

Ethical Issues

It was just *in vitro* study. The umbilical cord blood sample was collected with informed consent to separate MSCs.

Conflict of Interest

The authors have no conflicts of interest.

References

- Murphy MB, Moncivais K, Caplan AI. Mesenchymal stem cells: Environmentally responsive therapeutics for regenerative medicine. *Exp Mol Med* 2013;45:e54. doi: 10.1038/emm.2013.94
- Lotfinegad P, Shamsasenjan K, Movassaghpour A, Majidi J, Baradaran B. Immunomodulatory nature and site specific affinity of mesenchymal stem cells: A hope in cell therapy. *Adv Pharm Bull* 2014;4(1):5-13. doi: 10.5681/apb.2014.002
- Liu Q, Zheng Z. Clinical observation for the treatment of umbilical cord mesenchymal stem cells(UCMSC) on severe aplastic anemia(SAA). *Blood* 2016;128(22):5054.
- Sharma RR, Pollock K, Hubel A, McKenna D. Mesenchymal stem or stromal cells: A review of clinical applications and manufacturing practices. *Transfusion* 2014;54(5):1418-37. doi: 10.1111/trf.12421
- Amorin B, Alegretti AP, Valim V, Pezzi A, Laureano AM, da Silva MA, et al. Mesenchymal stem cell therapy and acute graft-versus-host disease: A review. *Hum Cell* 2014;27(4):137-50. doi: 10.1007/s13577-014-0095-x
- Amiri F, Molaei S, Bahadori M, Nasiri F, Deyhim MR, Jalili MA, et al. Autophagy-modulated human bone marrow-derived mesenchymal stem cells accelerate liver restoration in mouse models of acute liver failure. *Iran Biomed J* 2016;20(3):135-44.
- Roushandeh AM, Bahadori M, Roudkenar MH. Mesenchymal stem cell-based therapy as a new horizon for kidney injuries. *Arch Med Res* 2017;48(2):133-46. doi: 10.1016/j.arcmed.2017.03.007
- Zhaleh F, Amiri F, Mohammadzadeh-Vardin M, Bahadori M, Harati MD, Roudkenar MH, et al. Nuclear factor erythroid-2 related factor 2 overexpressed mesenchymal stem cells transplantation, improves renal function, decreases injuries markers and increases repair markers in glycerol-induced acute kidney injury rats. *Iran J Basic Med Sci* 2016;19(3):323-9.
- Fish KM, Hajjar RJ. Mesenchymal stem cells & endothelial function. *EBioMedicine* 2015;2(5):376-7. doi: 10.1016/j.ebiom.2015.04.015
- Santos Nascimento D, Mosqueira D, Sousa LM, Teixeira M, Filipe M, Resende TP, et al. Human umbilical cord tissue-derived mesenchymal stromal cells attenuate remodeling after myocardial infarction by proangiogenic, antiapoptotic, and endogenous cell-activation mechanisms. *Stem Cell Res Ther* 2014;5(1):5. doi: 10.1186/scrt394
- Moon HH, Joo MK, Mok H, Lee M, Hwang KC, Kim SW, et al. Msc-based vegf gene therapy in rat myocardial infarction model using facial amphipathic bile acid-conjugated polyethyleneimine. *Biomaterials* 2014;35(5):1744-54. doi: 10.1016/j.biomaterials.2013.11.019
- Toma C, Pittenger MF, Cahill KS, Byrne BJ, Kessler PD. Human mesenchymal stem cells differentiate to a cardiomyocyte phenotype in the adult murine heart. *Circulation* 2002;105(1):93-8.
- Sart S, Ma T, Li Y. Preconditioning stem cells for *in vivo* delivery. *Biores Open Access* 2014;3(4):137-49. doi: 10.1089/biores.2014.0012
- Han Q, Fan L, Heng BC, Zigang GE. Apoptosis and metabolism of mesenchymal stem cells during chondrogenic differentiation. *in vitro. Int J Tissu Regen* 2013;4(3):61-4.
- Francois S, Mouiseddine M, Allenet-Lepage B, Voswinkel J, Douay L, Benderitter M, et al. Human mesenchymal stem cells provide protection against radiation-induced liver injury by antioxidative process, vasculature protection, hepatocyte differentiation, and trophic effects. *Biomed Res Int* 2013;2013:151679. doi: 10.1155/2013/151679
- Li Q, Wang Y, Deng Z. Pre-conditioned mesenchymal stem cells: A better way for cell-based therapy. *Stem Cell Res Ther* 2013;4(3):63. doi: 10.1186/scrt213
- Wei H, Li Z, Hu S, Chen X, Cong X. Apoptosis of mesenchymal stem cells induced by hydrogen peroxide concerns both endoplasmic reticulum stress and mitochondrial death pathway through regulation of caspases, p38 and jnk. *J Cell Biochem* 2010;111(4):967-78. doi: 10.1002/jcb.22785
- Xu J, Qian J, Xie X, Lin L, Zou Y, Fu M, et al. High density lipoprotein protects mesenchymal stem cells from oxidative stress-induced apoptosis via activation of the PI3K/Akt pathway and suppression of reactive oxygen species. *Int J Mol Sci* 2012;13(12):17104-20. doi: 10.3390/ijms131217104

19. Amiri F, Jahanian-Najafabadi A, Roudkenar MH. In vitro augmentation of mesenchymal stem cells viability in stressful microenvironments : In vitro augmentation of mesenchymal stem cells viability. *Cell Stress Chaperones* 2015;20(2):237-51. doi: 10.1007/s12192-014-0560-1
20. Kapinya KJ. Ischemic tolerance in the brain. *Acta physiologica Hungarica* 2005;92(1):67-92. doi: 10.1556/APhysiol.92.2005.1.9
21. DeFily DV, Chilian WM. Preconditioning protects coronary arteriolar endothelium from ischemia-reperfusion injury. *Am J Physiol* 1993;265(2 Pt 2):H700-6. doi: 10.1152/ajpheart.1993.265.2.H700
22. Erpicum P, Detry O, Weekers L, Bonvoisin C, Lechanteur C, Briquet A, et al. Mesenchymal stromal cell therapy in conditions of renal ischaemia/reperfusion. *Nephrol Dial Transplant* 2014;29(8):1487-93. doi: 10.1093/ndt/gft538
23. Feng Y, Huang W, Wani M, Yu X, Ashraf M. Ischemic preconditioning potentiates the protective effect of stem cells through secretion of exosomes by targeting mecp2 via mir-22. *PLoS One* 2014;9(2):e88685. doi: 10.1371/journal.pone.0088685
24. Islam CF, Mathie RT, Dinneen MD, Kiely EA, Peters AM, Grace PA. Ischaemia-reperfusion injury in the rat kidney: The effect of preconditioning. *Br J Urol* 1997;79(6):842-7.
25. Li S, Deng Y, Feng J, Ye W. Oxidative preconditioning promotes bone marrow mesenchymal stem cells migration and prevents apoptosis. *Cell Biol Int* 2009;33(3):411-8. doi: 10.1016/j.cellbi.2009.01.012
26. Madrigal M, Rao KS, Riordan NH. A review of therapeutic effects of mesenchymal stem cell secretions and induction of secretory modification by different culture methods. *J Transl Med* 2014;12:260. doi: 10.1186/s12967-014-0260-8
27. Lin CS, Xin ZC, Dai J, Lue TF. Commonly used mesenchymal stem cell markers and tracking labels: Limitations and challenges. *Histol Histopathol* 2013;28(9):1109-16. doi: 10.14670/hh-28.1109
28. Molaei S, Roudkenar MH, Amiri F, Harati MD, Bahadori M, Jaleh F, et al. Down-regulation of the autophagy gene, ATG7, protects bone marrow-derived mesenchymal stem cells from stressful conditions. *Blood Res* 2015;50(2):80-6. doi: 10.5045/br.2015.50.2.80
29. Pagani FD, DerSimonian H, Zawadzka A, Wetzel K, Edge AS, Jacoby DB, et al. Autologous skeletal myoblasts transplanted to ischemia-damaged myocardium in humans. Histological analysis of cell survival and differentiation. *J Am Coll Cardiol* 2003;41(5):879-88.
30. Halabian R, Tehrani HA, Jahanian-Najafabadi A, Habibi Roudkenar M. Lipocalin-2-mediated upregulation of various antioxidants and growth factors protects bone marrow-derived mesenchymal stem cells against unfavorable microenvironments. *Cell Stress Chaperones* 2013;18(6):785-800. doi: 10.1007/s12192-013-0430-2
31. van Velthoven CT, Braccioli L, Willemen HL, Kavelaars A, Heijnen CJ. Therapeutic potential of genetically modified mesenchymal stem cells after neonatal hypoxic-ischemic brain damage. *Mol Ther* 2014;22(3):645-54. doi: 10.1038/mt.2013.260
32. Kiani AA, Kazemi A, Halabian R, Mohammadipour M, Jahanian-Najafabadi A, Roudkenar MH. Hif-1alpha confers resistance to induced stress in bone marrow-derived mesenchymal stem cells. *Arch Med Res* 2013;44(3):185-93. doi: 10.1016/j.arcmed.2013.03.006
33. Kinnaird T, Stabile E, Burnett MS, Lee CW, Barr S, Fuchs S, et al. Marrow-derived stromal cells express genes encoding a broad spectrum of arteriogenic cytokines and promote *in vitro* and *in vivo* arteriogenesis through paracrine mechanisms. *Circ Res* 2004;94(5):678-85. doi: 10.1161/01.res.0000118601.37875.ac
34. Tang XQ, Feng JQ, Chen J, Chen PX, Zhi JL, Cui Y, et al. Protection of oxidative preconditioning against apoptosis induced by H2O2 in PC12 cells: Mechanisms via MMP, ROS, and Bcl-2. *Brain Res* 2005;1057(1-2):57-64. doi: 10.1016/j.brainres.2005.07.072
35. Murry CE, Jennings RB, Reimer KA. Preconditioning with ischemia: A delay of lethal cell injury in ischemic myocardium. *Circulation* 1986;74(5):1124-36.
36. Rosova I, Dao M, Capoccia B, Link D, Nolte JA. Hypoxic preconditioning results in increased motility and improved therapeutic potential of human mesenchymal stem cells. *Stem Cells* 2008;26(8):2173-82. doi: 10.1634/stemcells.2007-1104
37. Gargioli C, Turturici G, Barreca MM, Spinello W, Fuoco C, Testa S, et al. Oxidative stress preconditioning of mouse perivascular myogenic progenitors selects a subpopulation of cells with a distinct survival advantage *in vitro* and *in vivo*. *Cell Death Dis* 2018;9(1):1. doi: 10.1038/s41419-017-0012-9
38. Boopathy AV, Pendergrass KD, Che PL, Yoon YS, Davis ME. Oxidative stress-induced Notch1 signaling promotes cardiogenic gene expression in mesenchymal stem cells. *Stem Cell Res Ther* 2013;4(2):43. doi: 10.1186/scrt190
39. Lu HH, Li YF, Sheng ZQ, Wang Y. Preconditioning of stem cells for the treatment of myocardial infarction. *Chin Med J (Engl)* 2012;125(2):378-84.
40. Zhang J, Chen GH, Wang YW, Zhao J, Duan HF, Liao LM, et al. Hydrogen peroxide preconditioning enhances the therapeutic efficacy of wharton's jelly mesenchymal stem cells after myocardial infarction. *Chin Med J (Engl)* 2012;125(19):3472-8.