

Research Article



## Contribution of Lysosome and Sigma Receptors to Neuroprotective Effects of Memantine Against Beta-Amyloid in the SH-SY5Y Cells

Mojtaba Keshavarz<sup>1\*</sup>, Majid Reza Farrokhi<sup>1</sup>, Elahe Amirinezhad Fard<sup>1</sup>, Mohammad Mehdipour<sup>2</sup>

<sup>1</sup>Shiraz Neuroscience Research Center, Shiraz University of Medical Sciences, Shiraz, Iran.

<sup>2</sup>Department of Neuroscience, School of Advanced Medical Sciences and Technologies, Shiraz University of Medical Sciences, Shiraz, Iran.

### Article info

#### Article History:

Received: 26 Sep. 2019

Revised: 27 Aug. 2019

Accepted: 14 Nov. 2019

published: 11 May 2020

#### Keywords:

- Amyloid beta-peptides
- Lysosomes
- Memantine
- Neuroprotection
- Sigma receptors

### Abstract

**Purpose:** Memantine is an approved drug for the treatment of Alzheimer's disease (AD). Autophagy, lysosome dysfunction, and sigma receptors have possible roles in the pathophysiology of AD. Therefore, we aimed to investigate the contribution of sigma receptors and lysosome inhibition to the neuroprotective effects of memantine against amyloid-beta (A $\beta$ )-induced neurotoxicity in SH-SY5Y cells.

**Methods:** We determined the neuroprotective effects of memantine (2.5  $\mu$ M), dizocilpine (MK801, as a selective N-methyl-D-aspartate (NMDA) receptor antagonist) (5  $\mu$ M) against A $\beta$ 25–35 (2  $\mu$ g/ $\mu$ L)-induced neurotoxicity. We used chloroquine (10, 20, and 40  $\mu$ M) as a lysosome inhibitor and BD-1063 (1, 10, and 30  $\mu$ M) as a selective sigma receptor antagonist. The MTT assay was used to measure the neurotoxicity in the SH-SY5Y cells. Data were analyzed using the one-way ANOVA.

**Results:** Memantine (2.5  $\mu$ M), dizocilpine (5  $\mu$ M), chloroquine (10 and 20  $\mu$ M) and BD-1063 (1, 10 and 30  $\mu$ M) decreased the neurotoxic effects of A $\beta$  on the SH-SY5Y cells. However, chloroquine (40  $\mu$ M) increased the neurotoxic effects of A $\beta$ . Cell viability in the cells treated with memantine + A $\beta$  + chloroquine (10, 20, and 40  $\mu$ M) was significantly lower than the memantine + A $\beta$ -treated group. Moreover, cell viability in the memantine + A $\beta$  group was higher than the memantine + A $\beta$  + BD-1063 (10 and 30  $\mu$ M) groups.

**Conclusion:** The lysosomal and sigma receptors may contribute to the neuroprotective mechanism of memantine and other NMDA receptor antagonists. Moreover, the restoration of lysosomes function and the modulation of sigma receptors are potential targets in the treatment of AD.

### Introduction

Neurodegeneration in several brain regions including neocortex and hippocampus may cause of the cognitive deficits in patients with Alzheimer's disease (AD). Amyloid-beta (A $\beta$ )-induced impairments in the activity of glutamate and autophagy systems may contribute to the neurodegeneration in the central nervous system (CNS).<sup>1,2</sup> Therefore, the restoration of homeostasis in the glutamate and autophagy systems may help block the A $\beta$ -induced neurotoxicity.

Memantine is a non-competitive antagonist of the N-methyl-D-aspartate (NMDA) receptors of glutamate approved for the treatment of AD.<sup>3</sup> Some human studies and animal models have shown its neuroprotective and cognitive-enhancing effects in the AD.<sup>4,5</sup> Other systems alongside with the NMDA receptors may have a role in the therapeutic effects of memantine.<sup>6</sup> By considering the possible roles of autophagy and sigma receptors in

the pathophysiology of the AD,<sup>7,8</sup> these systems may be involved in the mechanism of action of memantine.

Autophagy is a cellular homeostatic process that degrades noxious proteins and damaged organelles.<sup>9</sup> This process is mainly dependent on lysosomes.<sup>10</sup> Lysosomes are the specialized organelles contained hydrolase enzymes such as proteases and lipases.<sup>11</sup> This organelle is closely related to the digestive activity of the autophagy process.<sup>9</sup> Lysosome blockade is among various autophagy dysfunctions contributed to the neurodegeneration.<sup>12</sup> Chloroquine is a lysosomotropic agent which is widely used for the autophagy inhibition in the experimental models.<sup>13</sup> This agent increases the pH of lysosome and inhibits the function of lysosomal enzymes.<sup>14</sup> The A $\beta$ -induced over-activation of the NMDA receptors leads to the autophagy activation and neuronal apoptosis.<sup>15</sup> However, the contribution of lysosome inhibition to the neuroprotective effects of NMDA antagonists like

\*Corresponding Author: Mojtaba Keshavarz, Tel/Fax: +9871 3623 4508, Email: Mkeshavar@sums.ac.ir

© 2020 The Author (s). This is an Open Access article distributed under the terms of the Creative Commons Attribution (CC BY), which permits unrestricted use, distribution, and reproduction in any medium, as long as the original authors and source are cited. No permission is required from the authors or the publishers.

memantine is elusive.

Sigma receptors are widespread in the CNS and they are involved in different brain functions such as learning and memory, neuroprotection and immunomodulation.<sup>16,17</sup> Various experimental models have documented the neuroprotective effects of the sigma receptors in brain injury and neurodegeneration.<sup>18</sup> Sigma receptors regulate several neurotransmitter systems such as glutamatergic, noradrenalinergic, dopaminergic, and cholinergic systems.<sup>19</sup> Although the NMDA receptors can be modulated by the sigma-1 receptors,<sup>19,20</sup> the interaction between these two receptors has remained controversial.<sup>21</sup> Some studies have shown that the activation of the sigma receptor produces neuroprotection by interfering with the intracellular machinery of NMDA receptors.<sup>22</sup> By considering the role of the NMDA receptors in the neurotoxic effects of A $\beta$  peptide<sup>1</sup>, the sigma receptors may have a possible role in the neuroprotective effects of NMDA antagonists like memantine. In this study, we aimed to explore the contribution of sigma receptors and lysosome inhibition to the neuroprotective effects of memantine against A $\beta$ -induced neurotoxicity in the SH-SY5Y cells.

## Materials and Methods

### Materials

The human SH-SY5Y neuroblastoma cells were obtained from the Pasteur Institute (Iran). We purchased Dulbecco's modified Eagle's medium and Ham's nutrient mixture F-12 (DMEM/F12), fetal bovine serum (FBS), and Penicillin-Streptomycin from Gibco® life technologies™ (USA). Moreover, A $\beta$ 25-35, memantine, chloroquine, and BD-1063 (a selective sigma receptor antagonist) were obtained from the Sigma-Aldrich (USA).

### Neuronal cell culture

We used SH-SY5Y neuronal cells to determine the neurotoxic and neuroprotective activities of the administered agents. We seeded the SH-SY5Y cells at the density of  $1 \times 10^5$  cells/well in the 96-well plates. The cell culture medium contained DMEM/F12 (1:1), fetal bovine serum (10 %), penicillin (100 U/mL), and streptomycin (100  $\mu$ g/mL). The cells were kept in a standard condition of the humidified atmosphere of 95% air/5% CO<sub>2</sub> and 37°C.

### Study design and treatments

We used the fibril form of A $\beta$  to conduct the neurotoxicity test. A $\beta$ 25-35 (2  $\mu$ g/ $\mu$ L) was incubated in the water bath for four days at 37°C to induce the aggregation process. Memantine (2.5  $\mu$ M), dizocilpine (MK801, as a selective NMDA antagonist) (5  $\mu$ M), chloroquine (10, 20, 40  $\mu$ M), BD-1063 (1, 10, 30  $\mu$ M), and A $\beta$  (20  $\mu$ M) were dissolved in the sterile water. We selected the concentrations of each agent according to the previous studies and a pilot study. The cells were treated with A $\beta$  with or without

memantine, dizocilpine, chloroquine, and BD1063 for 24h. The procedure was repeated for four samples.

### Cell viability assay

The 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyl tetrazolium bromide (MTT) was used to measure the neuronal viability. In brief, the MTT reagent (5 mg/mL) was poured into the wells containing the treated cells. After 4 hours, the media was removed and the precipitate dissolved in dimethyl sulfoxide (DMSO) (100  $\mu$ L). The absorbance of each well was determined at 570 nm by a microplate reader (Synergy HT, Biotek®) as an index of cell viability.

### Statistical analysis

The results were analyzed using the one-way analysis of variance (ANOVA) test, followed by the LSD test as the post hoc test. *P* values <0.05 was considered statistically significant. We used the SPSS software (version 23) to analyze the data.


## Results and Discussion

### The effects of beta-amyloid, memantine, dizocilpine, chloroquine, and BD-1063 on neuronal survival


In the present study, A $\beta$  at the concentration of 20  $\mu$ M decreased neuronal survival compared to the control-treated group (*P* < 0.001; Figure 1). In contrast, memantine (2.5  $\mu$ M) and dizocilpine (5  $\mu$ M) suppressed the neurotoxic effects of A $\beta$  on the SH-SY5Y cells (*P* < 0.001) (Figure 1). Moreover, chloroquine (10 and 20  $\mu$ M, *P* < 0.001 and *P* < 0.05, respectively) and BD-1063 (1, 10 and 30  $\mu$ M, *P* < 0.001) decreased the neurotoxic effects of A $\beta$  on the SH-SY5Y cells (Figure 1 and Figure 2). However, chloroquine (40  $\mu$ M) increased the neurotoxic effects of A $\beta$  (*P* < 0.001; Figure 1).

The findings of this study showed that the NMDA antagonists such as memantine and dizocilpine inhibited the neurotoxic effects of A $\beta$ . *In vitro* and *in vivo* experiments have shown that memantine reduces the neurotoxic effects of A $\beta$ .<sup>23</sup> This effect might be due to the behavioral effects of memantine.<sup>4</sup> Moreover, Harkany et al<sup>23</sup> have shown that dizocilpine diminished the A $\beta$ -induced neurotoxicity. The over-activation of NMDA receptors may mediate the neurotoxic effects of A $\beta$ , and the inhibition of NMDA receptors suppresses this neurotoxic effect. Thus, the NMDA receptor antagonists have potential roles in the AD amelioration.<sup>24</sup> However, the cellular process of neuroprotection produced by the NMDA antagonists is relatively unknown.

Our study showed that the lower concentrations of chloroquine (10 and 20  $\mu$ M) suppressed the neurotoxic effects of A $\beta$ . In contrast, chloroquine (at a higher concentration: 40  $\mu$ M) increased the neurotoxic effects of A $\beta$ . According to previous studies, chloroquine has produced both neuroprotective or neurotoxic effects. The low concentration of chloroquine had no apparent effect on the lysosomes and produced neuroprotective



**Figure 1.** The effects of N-methyl-D-aspartate antagonists and chloroquine on the beta-amyloid (Aβ)-induced neurotoxicity in the SH-SY5Y cells (N=4). Data were analyzed using the one-way analysis of variance (ANOVA) followed by the LSD test. Cell viability was measured using the 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyl tetrazolium bromide (MTT) test. \*\*  $P < 0.001$  compared to the Aβ-treated group, ††  $P < 0.001$  compared to the Memantine + Aβ-treated group, ††  $P < 0.001$  compared to the MK801+Aβ-treated group, and \*  $P < 0.05$  compared to the Aβ + chloroquine-treated group. Chlor: chloroquine, MK: dizocilpine, Mem: memantine, Aβ: beta-amyloid.



**Figure 2.** The effects of N-methyl-D-Aspartate antagonists and a sigma receptor antagonist on the beta amyloid (Aβ)-induced neurotoxicity in the SH-SY5Y cells (N=4). Data were analyzed using the one-way analysis of variance (ANOVA) followed by the LSD test. Cell viability was measured using the 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyl tetrazolium bromide (MTT) test. \*\*  $P < 0.001$  compared to the Aβ-treated group, †  $P < 0.005$  compared to the Memantine + Aβ-treated group, † and ††  $P < 0.05$  and  $P < 0.001$  compared to the MK801+Aβ-treated group, and †  $P < 0.05$  compared to the Aβ + BD1063-treated group. BD: BD1063, MK: dizocilpine, Mem: memantine, Aβ: beta-amyloid.

effects in the different models.<sup>25,26</sup> In contrast, chloroquine at the concentrations higher than 25 μM decreased primary telencephalic neuronal viability.<sup>27</sup> Another *in vitro* study has also demonstrated the neurotoxic effects of chloroquine at concentrations higher than 25 μM.<sup>28</sup> Accordingly, chloroquine may have detrimental or beneficial effects on neurons.

### Chloroquine effects on the memantine and dizocilpine neuroprotective effects

Cell viability in the SH-SY5Y cells treated with memantine, dizocilpine and chloroquine were different ( $F(12,39) = 47.764, P < 0.001$ ) (Figure 1). Accordingly, cell viability in the cells treated with memantine + Aβ +

chloroquine (10, 20, and 40 μM) was significantly lower than the memantine + Aβ-treated group ( $P < 0.001$ ) (Figure 1). Moreover, cell viability in the memantine + Aβ + chloroquine (20 μM) group was higher compared to the Aβ + chloroquine (20 μM) group ( $P = 0.025$ ) (Figure 1). Our study also showed that cell viability in the dizocilpine + Aβ + chloroquine (10, 20, and 40 μM) was significantly lower than the dizocilpine + Aβ-treated group ( $P < 0.001$ ; Figure 1). Moreover, cells treated with Aβ + chloroquine (40 μM) showed lower cell viability compared to the dizocilpine + Aβ + chloroquine (40 μM)-treated group ( $P = 0.004$ ; Figure 1). The present study showed that cell viability in the memantine + Aβ + chloroquine (20 μM) group was higher than the dizocilpine + Aβ + chloroquine

(20  $\mu$ M) group ( $P=0.006$ ; Figure 1).

The inhibition of lysosomal activity may be related to the neurotoxic effects of chloroquine. Lysosome dysfunction leads to A $\beta$  accumulation and enhances neurotoxicity.<sup>29</sup> Simultaneously, A $\beta$  causes lysosomal dysfunction in the SH-SY5Y cells.<sup>30</sup> Moreover, the neurotoxic effects of A $\beta$  are similar to lysosomotropic agents.<sup>30</sup> On the contrary, the pharmacologic induction of autophagy produces neuroprotective effects against proteinopathies.<sup>31</sup> Chloroquine is a weak base that disrupts the lysosome activity in neurons by the inhibition of autophagosomes inhibition to the lysosome.<sup>32</sup> Therefore, the chloroquine inhibitory effects on lysosome may block the A $\beta$  degradation and increase the A $\beta$ -induced neurotoxicity. In this regard, *in vivo* studies have shown that lysosomal inhibition may lead to the reduction of A $\beta$  degradation and neurodegeneration.<sup>26,33</sup> Therefore, the chloroquine effects on lysosome especially at the higher concentrations overcomes the neuroprotective effects of this agent and potentiates the neurotoxic effects of A $\beta$ .

There are some controversies regarding the memantine effects on the autophagy. In a study conducted by Yoon et al, the authors reported that a high concentration of memantine activated the autophagy and apoptosis in the glioma cell lines.<sup>34</sup> In contrast, memantine has decreased autophagy in a cellular model of the AD.<sup>15</sup> The difference may be due to the dose of memantine and the cells used in these two experiments. However, there are limited data about the memantine effects on the lysosomal activity. According to our study, lysosome and autophagy inhibition decreased the neuroprotective effects of memantine and dizocilpine. Therefore, autophagy activation and the normal lysosomal function may be protective against the A $\beta$ -induced neurotoxicity. In this regard, the autophagy induction has potentiated neuroprotection against A $\beta$ -induced neurotoxicity in the SH-SY5Y cells.<sup>35</sup> Accordingly, the NMDA receptor activation may cause lysosomal defect and magnify the A $\beta$  neurotoxic effects. It is possible to assume that the NMDA antagonists may modulate the lysosome activity and autophagy in the process of neurodegeneration.

#### ***BD-1063 interference with the memantine and dizocilpine neuroprotective effects***

The present study showed that cell viability in the SH-SY5Y cells treated with memantine, dizocilpine, and BD-1063 was different ( $F(12, 39)=14.150$ ,  $P<0.001$ ; Figure 2). In this regard, cell viability in the memantine + A $\beta$  group was higher than the memantine + A $\beta$  + BD-1063 (10 and 30  $\mu$ M) groups ( $P=0.003$  and  $P=0.002$ , respectively; Figure 2). Furthermore, cell viability in the A $\beta$  + BD-1063 (30  $\mu$ M) group was lower than the memantine + A $\beta$  + BD-1063 (30  $\mu$ M) group ( $P=0.020$ ; Figure 2). Our study also showed that cell viability in the dizocilpine + A $\beta$  group was higher compared to the dizocilpine + A $\beta$  + BD-1063 (1, 10, and 30  $\mu$ M) ( $P=0.010$ ,  $P=0.032$ , and  $P<0.001$ ,

respectively) (Figure 2). Furthermore, cell viability in the A $\beta$  + BD-1063 (10  $\mu$ M) group was significantly lower than the dizocilpine + A $\beta$  + BD-1063 (10  $\mu$ M) group ( $P=0.021$ ; Figure 2). Cell viability in the memantine + A $\beta$  + BD-1063 (30  $\mu$ M) group was higher than the dizocilpine + A $\beta$  + BD-1063 (30  $\mu$ M) group ( $P=0.035$ ).

This study showed that a sigma receptor antagonist suppressed the A $\beta$ -induced neurotoxicity. There are some inconsistencies in the literature about the neuroprotective or neurotoxic effects of sigma receptors. Some reports have shown that the sigma receptor agonists have produced neuroprotective effects in the neurodegenerative disorder models.<sup>18</sup> On the other hand, BD1063, as a selective sigma receptor antagonist, protected against methamphetamine-induced neurotoxicity in the rat.<sup>36</sup> *In vitro* and *in vivo* studies have also shown that AC927, another sigma receptor antagonist, prevented the methamphetamine-induced neurotoxicity.<sup>37</sup> Therefore, the inconsistencies may be related to the models and the neurotoxic agent have been used in these experiments.

Our study showed that a sigma receptor antagonist decreased the neuroprotective effects of two NMDA antagonists against A $\beta$ -induced neurotoxicity. Thus, the activation of sigma receptors may potentiate the neuroprotective effects of memantine and dizocilpine. Other studies have shown that the activation of sigma receptors prevented glutamate-mediated neurotoxicity.<sup>38</sup> In contrast to our results, some studies have shown that the sigma receptor agonists have potentiated the glutamate activity by increasing the expression of the NMDA receptors subunits and the plasma membrane level of NMDA receptors.<sup>22</sup> These inconsistencies may be related to the different interactions between NMDA receptors and sigma receptors in the normal and degenerative conditions.

Our study had some limitations. We did not measure the lysosomal function and NMDA receptor trafficking in the treated cells. Moreover, some agents may have non-specific effects. Thus, future studies may help to discover the cellular interaction between the NMDA antagonists and the lysosome and sigma receptor functions in the process of neurodegeneration.

#### **Conclusion**

The lysosomal function and sigma receptors may contribute to the neuroprotective mechanism of memantine and other NMDA receptor antagonists. Moreover, the restoration of lysosomal function and the modulation of sigma receptors may be potential strategies in the treatment of AD. Future studies may help to discover the exact contribution of lysosome and sigma receptor functions to the neuroprotective effects of the NMDA receptors.

#### **Ethical Issues**

Not applicable.



**Conflict of Interest**

Non-declared.

**Acknowledgments**

We gratefully thank the Vice-chancellor for research affairs of the Shiraz University of Medical Science for the financial support. It is important to note that sponsor had no role in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

**References**

1. Harkany T, Abrahám I, Timmerman W, Laskay G, Tóth B, Sasvári M, et al. beta-amyloid neurotoxicity is mediated by a glutamate-triggered excitotoxic cascade in rat nucleus basalis. *Eur J Neurosci* 2000;12(8):2735-45. doi: 10.1046/j.1460-9568.2000.00164.x
2. Zare-Shahabadi A, Masliah E, Johnson GV, Rezaei N. Autophagy in Alzheimer's disease. *Rev Neurosci* 2015;26(4):385-95. doi: 10.1515/revneuro-2014-0076
3. Peskind ER, Potkin SG, Pomara N, Ott BR, Graham SM, Olin JT, et al. Memantine treatment in mild to moderate Alzheimer disease: a 24-week randomized, controlled trial. *Am J Geriatr Psychiatry* 2006;14(8):704-15. doi: 10.1097/01.jgp.0000224350.82719.83
4. Reisberg B, Doody R, Stöffler A, Schmitt F, Ferris S, Möbius HJ. Memantine in moderate-to-severe Alzheimer's disease. *N Engl J Med* 2003;348(14):1333-41. doi: 10.1056/NEJMoa013128
5. Rogawski MA, Wenk GL. The neuropharmacological basis for the use of memantine in the treatment of Alzheimer's disease. *CNS Drug Rev* 2003;9(3):275-308. doi: 10.1111/j.1527-3458.2003.tb00254.x
6. Johnson JW, Kotermanski SE. Mechanism of action of memantine. *Curr Opin Pharmacol* 2006;6(1):61-7. doi: 10.1016/j.coph.2005.09.007
7. Nguyen L, Lucke-Wold BP, Mookerjee SA, Cavendish JZ, Robson MJ, Scandinaro AL, et al. Role of sigma-1 receptors in neurodegenerative diseases. *J Pharmacol Sci* 2015;127(1):17-29. doi: 10.1016/j.jphs.2014.12.005
8. Li Q, Liu Y, Sun M. Autophagy and Alzheimer's disease. *Cell Mol Neurobiol* 2017;37(3):377-88. doi: 10.1007/s10571-016-0386-8
9. Uddin MS, Stachowiak A, Mamun AA, Tzvetkov NT, Takeda S, Atanasov AG, et al. Autophagy and Alzheimer's disease: from molecular mechanisms to therapeutic implications. *Front Aging Neurosci* 2018;10:04. doi: 10.3389/fnagi.2018.00004
10. Levine B, Kroemer G. Autophagy in the pathogenesis of disease. *Cell* 2008;132(1):27-42. doi: 10.1016/j.cell.2007.12.018
11. Xu H, Ren D. Lysosomal physiology. *Annu Rev Physiol* 2015;77:57-80. doi: 10.1146/annurev-physiol-021014-071649
12. Boland B, Nixon RA. Neuronal macroautophagy: from development to degeneration. *Mol Aspects Med* 2006;27(5-6):503-19. doi: 10.1016/j.mam.2006.08.009
13. Shacka JJ, Klocke BJ, Shibata M, Uchiyama Y, Datta G, Schmidt RE, et al. Bafilomycin A1 inhibits chloroquine-induced death of cerebellar granule neurons. *Mol Pharmacol* 2006;69(4):1125-36. doi: 10.1124/mol.105.018408
14. de Duve C. The lysosome turns fifty. *Nat Cell Biol* 2005;7(9):847-9. doi: 10.1038/ncb0905-847
15. Song G, Li Y, Lin L, Cao Y. Anti-autophagic and anti-apoptotic effects of memantine in a SH-SY5Y cell model of Alzheimer's disease via mammalian target of rapamycin-dependent and -independent pathways. *Mol Med Rep* 2015;12(5):7615-22. doi: 10.3892/mmr.2015.4382
16. Maurice T, Su TP. The pharmacology of sigma-1 receptors. *Pharmacol Ther* 2009;124(2):195-206. doi: 10.1016/j.pharmthera.2009.07.001
17. Rousseaux CG, Greene SE. Sigma receptors [sigmaRs]: biology in normal and diseased states. *J Recept Signal Transduct Res* 2016;36(4):327-88. doi: 10.3109/10799893.2015.1015737
18. Ruscher K, Wieloch T. The involvement of the sigma-1 receptor in neurodegeneration and neurorestoration. *J Pharmacol Sci* 2015;127(1):30-5. doi: 10.1016/j.jphs.2014.11.011
19. Cobos EJ, Entrena JM, Nieto FR, Cendán CM, Del Pozo E. Pharmacology and therapeutic potential of sigma(1) receptor ligands. *Curr Neuropharmacol* 2008;6(4):344-66. doi: 10.2174/157015908787386113
20. Kerchner GA, Nicoll RA. Silent synapses and the emergence of a postsynaptic mechanism for LTP. *Nat Rev Neurosci* 2008;9(11):813-25. doi: 10.1038/nrn2501
21. Balasuriya D, Stewart AP, Edwardson JM. The sigma-1 receptor interacts directly with GluN1 but not GluN2A in the GluN1/GluN2A NMDA receptor. *J Neurosci* 2013;33(46):18219-24. doi: 10.1523/jneurosci.3360-13.2013
22. Pabba M, Sibille E. Sigma-1 and N-methyl-D-aspartate receptors: a partnership with beneficial outcomes. *Mol Neuropsychiatry* 2015;1(1):47-51. doi: 10.1159/000376549
23. Harkany T, Mulder J, Sasvári M, Abrahám I, Kónya C, Zarándi M, et al. N-Methyl-D-aspartate receptor antagonist MK-801 and radical scavengers protect cholinergic nucleus basalis neurons against beta-amyloid neurotoxicity. *Neurobiol Dis* 1999;6(2):109-21. doi: 10.1006/nbdi.1998.0230
24. Dong XX, Wang Y, Qin ZH. Molecular mechanisms of excitotoxicity and their relevance to pathogenesis of neurodegenerative diseases. *Acta Pharmacol Sin* 2009;30(4):379-87. doi: 10.1038/aps.2009.24
25. Hirata Y, Yamamoto H, Atta MS, Mahmoud S, Oh-hashii K, Kiuchi K. Chloroquine inhibits glutamate-induced death of a neuronal cell line by reducing reactive oxygen species through sigma-1 receptor. *J Neurochem* 2011;119(4):839-47. doi: 10.1111/j.1471-4159.2011.07464.x
26. Mielke JG, Murphy MP, Maritz J, Bengualid KM, Ivy GO. Chloroquine administration in mice increases beta-amyloid immunoreactivity and attenuates kainate-induced blood-brain barrier dysfunction. *Neurosci Lett* 1997;227(3):169-72. doi: 10.1016/s0304-3940(97)00340-6
27. Zaidi AU, McDonough JS, Klocke BJ, Latham CB, Korsmeyer SJ, Flavell RA, et al. Chloroquine-induced neuronal cell death is p53 and Bcl-2 family-dependent but caspase-independent. *J Neuropathol Exp Neurol* 2001;60(10):937-45. doi: 10.1093/jnen/60.10.937
28. Geng Y, Kohli L, Klocke BJ, Roth KA. Chloroquine-induced autophagic vacuole accumulation and cell death in glioma cells is p53 independent. *Neuro Oncol* 2010;12(5):473-81. doi: 10.1093/neuonc/nop048

29. Khurana V, Elson-Schwab I, Fulga TA, Sharp KA, Loewen CA, Mulkearns E, et al. Lysosomal dysfunction promotes cleavage and neurotoxicity of tau in vivo. *PLoS Genet* 2010;6(7):e1001026. doi: 10.1371/journal.pgen.1001026
30. Yang AJ, Chandswangbhuvana D, Margol L, Glabe CG. Loss of endosomal/lysosomal membrane impermeability is an early event in amyloid Abeta1-42 pathogenesis. *J Neurosci Res* 1998;52(6):691-8. doi: 10.1002/(sici)1097-4547(19980615)52:6<691::aid-jnr8>3.0.co;2-3
31. Thellung S, Scoti B, Corsaro A, Villa V, Nizzari M, Gagliani MC, et al. Pharmacological activation of autophagy favors the clearing of intracellular aggregates of misfolded prion protein peptide to prevent neuronal death. *Cell Death Dis* 2018;9(2):166. doi: 10.1038/s41419-017-0252-8
32. Boya P, González-Polo RA, Casares N, Perfettini JL, Dessen P, Larochette N, et al. Inhibition of macroautophagy triggers apoptosis. *Mol Cell Biol* 2005;25(3):1025-40. doi: 10.1128/mcb.25.3.1025-1040.2005
33. Frautschy SA, Horn DL, Sigel JJ, Harris-White ME, Mendoza JJ, Yang F, et al. Protease inhibitor coinfusion with amyloid beta-protein results in enhanced deposition and toxicity in rat brain. *J Neurosci* 1998;18(20):8311-21.
34. Yoon WS, Yeom MY, Kang ES, Chung YA, Chung DS, Jeun SS. Memantine induces NMDAR1-mediated autophagic cell death in malignant glioma cells. *J Korean Neurosurg Soc* 2017;60(2):130-7. doi: 10.3340/jkns.2016.0101.006
35. Singh AK, Bissoyi A, Kashyap MP, Patra PK, Rizvi SI. Autophagy activation alleviates amyloid-beta-induced oxidative stress, apoptosis and neurotoxicity in human neuroblastoma SH-SY5Y cells. *Neurotox Res* 2017;32(3):351-61. doi: 10.1007/s12640-017-9746-5
36. Shaikh J, Matsumoto RR. Sigma ( $\sigma$ ) receptor antagonist, BD1063, protects against methamphetamine-induced hyperthermia and neurotoxicity in mice. *FASEB J* 2006;20(4):A676. doi: 10.1096/fasebj.20.4.A676-a
37. Matsumoto RR, Shaikh J, Wilson LL, Vedam S, Coop A. Attenuation of methamphetamine-induced effects through the antagonism of sigma ( $\sigma$ ) receptors: evidence from in vivo and in vitro studies. *Eur Neuropsychopharmacol* 2008;18(12):871-81. doi: 10.1016/j.euroneuro.2008.07.006
38. Kume T, Nishikawa H, Taguchi R, Hashino A, Katsuki H, Kaneko S, et al. Antagonism of NMDA receptors by sigma receptor ligands attenuates chemical ischemia-induced neuronal death in vitro. *Eur J Pharmacol* 2002;455(2-3):91-100. doi: 10.1016/s0014-2999(02)02582-7