

ARTHRAOXON P. BEAUV (GRAMINEAE) AND CAREX CARYOPHYLLEA (CYPERACEAE), NEW GENUS AND SPECIES RECORDS FROM IRAN

B. Hamzeh'ee & A. Naqinezhad

Received 10.10.2008. Accepted for publication 05.03.2009

Hamzeh'ee, B. & Naqinezhad A. 2009 06 30: *Arthraxon P. Beauv (Gramineae) and Carex caryophyllea (Cyperaceae)*: New genus and species records from Iran. -*Iran. J. Bot.* 15 (1): 68-71. Tehran.

Arthraxon hispidus (Thunb.) Makino var. *hispidus* (Gramineae) and *Carex caryophyllea* Latourr. (Cyperaceae) are recorded for the first time as new genus and species for the flora of Iran. These taxa have been collected from Kiashahr (Gilan prov.) and Arasbaran Protected Area (Azerbaijan prov.) in N and NW of Iran. Taxonomical characteristics, localities, habitats, geographical distributions and the closest genus and species are discussed here.

Bhnam Hamzeh'ee (correspondence), Research Institute of Forests and Rangelands, P. O. Box 13185-116, Tehran, Iran.— Alireza Naqinezhad, Department of Biology, Faculty of Science, University of Mazandaran, Babolsar, Mazandaran.

گزارش یک جنس (*Arthraxon*) از تیره گندمیان و یک گونه جدید (*Carex caryophyllea*) از تیره جگن برای فلور ایران بهنام حمزه، مربی پژوهش مؤسسه تحقیقات جنگلها و مراتع کشور. علیرضا نقی نژاد، استادیار گروه زیست‌شناسی دانشگاه مازندران.

Carex caryophyllea Latourr. و *Arthraxon hispidus* (Thunb.) Makino var. *hispidus* به عنوان جنس جدید از تیره گندمیان و به عنوان گونه جدید از تیره جگن برای اولین بار از ایران گزارش می گردند. جنس و گونه معرفی شده به ترتیب از استان های گیلان (کیاشهر) و آذربایجان شرقی (منطقه حفاظت شده ارسباران) جمع آوری شده‌اند. ویژگی‌های تاکسونومیک، محل دقیق جمع آوری، رویشگاه و پراکنش جغرافیایی هر یک توضیح داده شده است. همچنین تاکسون‌های فوق با نزدیک‌ترین جنس و گونه مورد مقایسه قرار گرفته است.

Introduction

During floristic studies, a new genus of grass (*Arthraxon hispidus* (Thunb.) Makino var. *hispidus* and a new species of sedge (*Carex caryophyllea* Latourr.) were collected from Kiashahr (Gilan prov.) and Arasbaran Protected Area (Azerbaijan prov.), Iran. The above-mentioned taxa have not been recorded in Flora Iranica area until now (Bor 1970; Kukkonen 1998). They were determined by Grasses of the Soviet Union (Tsvelev 1976) and Flora of Turkey (Mill, 1985 and Nilsson 1985) respectively. *Arthraxon P. Beauv. (Andropogoneae)* includes ca. 30 species in the tropical and subtropical countries of Eurasia and Africa and as adventives plants in many other countries of both hemispheres (Tsvelev 1976) but only two species were recognized in Flora Iranica area (Bor 1970). Introducing of this weed can be highlighted due to its possible distribution into the Hyrcanian forest in Iran like another adventives genus of *Andropogoneae*

(*Microstegium vimineum* (Trin.) A. Camus) which is widespread species in cleared and open places of the forests. A comparison between *Carex caryophyllea* and its much closed species, *C. umbrosa* Host subsp. *huetiana* (Boiss.) Soo is discussed. Including the current new record, 63 taxa of genus *Carex* occur in the flora of Iran by now (Assadi 1988; Akhiani, 1998; Naqinezhad & Ghahreman, 2002; Amini Rad, 2003, 2005a,b, 2006, Naqinezhad et al., 2005; Naqinezhad et al, 2008). All identified specimens are preserved in the TARI and Iran Natural History Museum herbaria.

Arthraxon hispidus (Thunb.) Makino var. **hispidus** - (Fig. 1).

Examined specimens. Gilan province: Astaneh Ashrafieh, Kiashahr, Amir Kiasar, near to Tavakkol Parry, - 25 m, 25.8.2005, A. Naqinezhad, 635-Iran Natural History Museum herbarium, duplicate in TARI.

This plant is mentioned in Flora Iranica (Bor 1970) under the name of *Arthraxon hispidus* but is divided into four subspecies in the Grasses of the Soviet Union (Tsvelev 1976). Based on Flora of Turkey (Mill 1985), this plant is an extremely variable species throughout its immense range and numerous varieties have been described, many of which have been reduced to synonymy. Mill (1985) was in agreement with van Welzen (1981) that these are only minor variants of no taxonomic significance. It seems that the taxonomical characters of our material in Iran have fitness to conclusions of Mill (l. c.), therefore, this plant was named based on Flora of Turkey. *Arthraxon hispidus* var. *hispidus* differs from *Microstegium vimineum* (Trin.) A. Camus by its cordate lanceolate – ovate or broadly lanceolate lamina with bulbous-based spinules and awned sessile spikelets. A short description of the plant is as follows:

Annual, stem ca. 50 cm, nodes pubescent. Leaf blades lanceolate – ovate or broadly lanceolate, slightly cordate at the base with bulbous - based spinules in lower (1/8) 1/3 (-1/2), ciliate in upper part, surfaces pubescent. Inflorescence ca. 3 cm × 2 - 5 mm, with 2-3 branches bearing numerous spikelets. Sessile spikelets only present, ca. 4 mm long, alternate on spicate branches; pedicellate spikelets reduced to minute pedicels. Glumes spinulate along veins; upper lemma 1- veined, with a geniculate awn. Palea absent. Anthers 2.

General distribution. Caucasus, India, China, Japan, Malaysia, C. & E. Africa, Australia, Iran, introduced in Hawaii and eastern C. & N. America.

Carex caryophyllea Latourr.

Examined specimens. Azerbaijan: Arasbaran Protected Area, Ilankosh, 2050 m; Hamzeh'ee & Asri, 81844-TARI; scattered on meadows.

This plant was not recorded from Iran (Kukkonen 1998). Closest locality to the Iranian collection is in Talish, Azerbaijan. This species is considered as two other separated species in Flora of USSR, i.e. *C. scabriscuspis* V. Krecz. and *C. verna* Chaix (Kreczetovicz 1935).

Caespitose. Stem 15-20 cm high, subterete, scabrous above, leafy only at base. Leaves 1.3-2.5 mm broad, keeled, flat, c. 1/3-1/2 of stem length, erect or slightly falcate; ligule ca. 0.2 mm long. Male spike 1, obovoid or oblong-clavate. 8-12 x 3 mm. Female spikes 1 - 3, ovoid to oblong, partly overlapping each other and male spike, 9-11 x 4 mm; lowest bract glumaceous,

setaceous, clasping or with sheath to 5 mm long, nearly equaling its spike or shorter. Female glumes equaling utricles, or slightly shorter, ovate, acute or mucronate-aristate; arista to 0.5 mm long. Utricles obovoid, 2.5-3 mm long, puberulent, shortly stipitate, rather abruptly contracted into a 0.3–0.5 mm long, emarginated to bifid beak. Nut 1.5-1.7 x 1-1.2 mm, annulate, finely papillose.

Carex caryophyllea is similar to *C. umbrosa* Host subsp. *huetiana* (Boiss.) Kük, but laxly caespitose, stem slightly scabrid above and with shortly aristulate glume (Nilsson 1985). This species is also related to *Carex depressa* Link, but there are some distinct differences between them. The occurrence of longer lowest bract, longer utricule and nut and a stem based-originated female spike are most important distinguishing characters of the latter species. It seems that another *Carex* species on the herbarium sheet (81844-TARI) is *Carex depressa* Link subsp. *transilvanica* (Schur) Egor.

General distribution. Pyrenees, Balkans, Caucasus and Turkey.

Acknowledgements

The authors would like to thank Mrs. Habibi the artist in the TARI herbarium, for preparing of the illustrations.

References

- Akhani, H. 1998: Plant biodiversity of Golestan National Park, Iran. -*Stapfia* 53: 1-411.
- Amini Rad, M. 2003: Two new records of Cyperaceae family from Iran. -*Iran J. Bot.* 10 (1): 31–34.
- Amini Rad, M. 2005a: Reinvestigation of *Carex depressa* ssp. *transilvanica* in Iran. -*Rostaniha* 6 (1): 36-37.
- Amini Rad, M. 2005b: New records and interesting species of Cyperaceae family from Iran. -*Iran J. Bot.* 11 (1): 49-53.
- Amini Rad, M. 2006: New taxa records of *Carex* from Iran. -*Rostaniha* 7 (2): 163- 175.
- Assadi, M. 1988. Plants of Arasbaran Protected Area, NW Iran (Part II). -*Iran. J. Bot.* 4 (1): 37-38.
- Bor, N. L. 1970: Graminae in Rechinger, K. H. (ed.), *Flora Iranica*. no. 70— Graz Akademische Druck- und Verlagsanstalt.
- Kreczetovicz, V. I. 1935: *Carex* in B. K. Schischkin (ed.) *Flora of USSR*, vol. 3: 112-464. Izdatel'stvo Akademii Nauk SSSR Moskva, Leningrad
- Kukkonen, H. 1998. Cyperaceae in Rechinger K. H. (ed.) *Flora Iranica* no. 173: 168-293. -Graz: Akademische Druck-u. Verlagsanstalt.



Fig. 1. *Arthraxon hispidus* var. *hispidus* ($\times 0.9$); leaf ($\times 9$); lemma ($\times 18$); glumes ($\times 13$).




Fig. 2. *Carex caryophyllea* ($\times 1.2$); details ($\times 18$).

Mill, R. R. 1985: *Arthraxon* P. Beauv. in Davis, P. H. (ed.), *Flora of Turkey*, vol. 9: 615-616. -Edinburgh University Press.

Naqinezhad, A. & Ghahreman, A. 2002: Two new records of Cyperaceae from coastal flora of the Caspian sea, N. Iran. - *Iran. J. Bot.* 9 (2):171-175.

Naqinezhad, A., Ghahreman, A. & Assadi, M. 2005: Some new record species for the flora of Iran as well as ecological and phytogeographical notes. - *Iran. J. Bot.* 11 (1): 89-95.

Naqinezhad, A., Jalili, A., Attar, F., Ghahreman, & Maassoumi, A. A. 2008: Two new records from

wetland habitats of the central Alborz mountains, Iran. -*Turkish Journal of Botany* 38: 1-5.

Nilsson, O. 1985: *Carex* in Davis, P. H. (ed.), *Flora of Turkey*, vol. 9: 21-158. -Edinburgh University Press.

Tsvelev, N. N. 1976: *Grasses of the Soviet Union*. Part I - II. -Leningrad. (English version 1983, New Delhi).

Van Welzen, P. C. 1981: A taxonomic revision of the genus *Arthraxon* Beauv. (*Gramineae*). -*Blumea* 27: 255-300.