



The Trend of Changes in the Risk Factors of Coronary Heart Disease from 1980 to 2010 in Iran

Manijeh Alimohammadi,¹ Yousef Alimohamadi,^{2,3} Malihe Khoramdad,⁴ Mohammad Shahbaz,⁵ Amir Abbas Shokraeian,⁶ and Firooz Esmaeilzadeh^{7,*}

¹Noor Research Center for Ophthalmic Epidemiology, Noor Eye Hospital, Tehran, Iran

²Pars Advanced and Minimally Invasive Medical Manners Research Center, Pars Hospital, Tehran, Iran

³PhD Student of Epidemiology, Department of Epidemiology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

⁴MSc, Faculty of Health, Kermanshah University of Medical Sciences, Kermanshah, Iran

⁵MSc of Epidemiology, Department of Epidemiology, School of Public Health, Shahid Beheshti University of Medical Sciences, Tehran, Iran

⁶D.V.M Graduated, Faculty of Veterinary Medicine, Shahid Chamran University of Ahvaz, Ahvaz, Iran

⁷Department of Public Health, School of Public Health, Maragheh University of Medical Sciences, Maragheh, Iran

*Corresponding author: Firooz Esmaeilzadeh, Department of Public Health, School of Public Health, Maragheh University of Medical Sciences, Maragheh, Iran. E-mail: firooz.esmaeilzadeh@gmail.com

Received 2016 August 08; Revised 2017 January 25; Accepted 2018 February 07.

Abstract

Background: Industrialization is the significant cause of death and inability in the most developed and developing countries, and a greater part of nourishing inadequacies and infectious diseases is attributed to this outcome of modernity. The current study aimed at determining the status of the cardiovascular disease progression in Iran (as a developing country) by assessing the trend of risk factors such as blood pressure (PB), total cholesterol, and body mass index (BMI) from 1980 to 2010.

Methods: The current cross sectional study was conducted based on aggregate data extracted from the Gapminder Website. No sampling was done in the study. The research was conducted from 1980 to 2010. The bivariate Pearson correlation, independent samples t test, and message sequence charts were used to analyze data with SPSS version 16; $\alpha = 0.05$ was considered the level of significance.

Results: The mean total cholesterol in females over 3 decades showed an increasing trend. The mean BMI increased within the study period. The mean BMI of males decreased from 1980 to 1997 and then, increased until 2008; while it remained constant in 2009 and 2010. The mean PB decreased in females aged above 31 years. The total cholesterol levels in males decreased from 1980 to 1990 and afterwards increased to 2010. The mean BMI of males increased in the same period.

Conclusions: By the improvement of lifestyle, healthy nutrition, purposive training, and increasing the knowledge about risk factors of chronic heart disease (CHD), these risk factors as well as the rate of CHD reduced in the studied population.

Keywords: Risk Factors, Coronary, Heart, Iran

1. Background

The health status and disease profile of human societies are identified based on the economic improvement and social development. By industrialization, a major cause of humans' death and inability, most of the developed and developing societies faced nourishing inadequacies, infectious diseases, and even chronic illnesses such as cardiovascular disease (CVD), cancer, and diabetes (1, 2). The CVD rate in the developed and developing countries vary depending on geographical area. A few studies reported the association between the development of CVD and routine risk factors, such as hypertension, total cholesterol, and smoking, which seem to decrease with aging (3, 4). Some other data suggested aggravating prevalence of other risk factors of CVD; for example, diabetes, obesity,

and the metabolic syndrome (5, 6). CVD, cancers, and respiratory diseases are the 3 leading causes of mortality and morbidity. Obesity increases the risk of deaths due to any of the abovementioned diseases, and is additionally associated with different complications such as infertility, osteoarthritis, etc. (7). Body mass index (BMI), considered as a risk factor in the current study, is a weight: height ratio and the most commonly utilized tool and the basic measure of body size, which is used as often as possible to evaluate the prevalence of obesity in a population (8, 9). BMI is an index, which is reliably associated with predisposition to CVD and type 2 diabetes (10). Hypertension is the most common controllable symptom of heart and vascular problems, representing about half of the risks in the developed countries. It affects 50% of adults and is the leading cause of

drug prescription (11). Numerous studies evaluated the relationship between hypertension and CVD and indicated a significant, predictable, and independent relationship between the 2 diseases (12). Hypercholesterolemia is a conceivably critical modifiable risk factor for cardiovascular problems. It is noteworthy that about 25% of males and 42% of females over 65 years old have total cholesterol of ≥ 240 mg/dL. It is evident that cholesterol is a key contributor to CVD, except for artery diseases (13). For several decades, the rate of changes in the risk factors of CVD in Iran was not measured. The present study aimed at determining the status of CVD in Iran (as a developing country) by assessing the trend of risk factors such as blood pressure (BP), total cholesterol, and BMI from 1980 to 2010.

2. Methods

The present cross sectional study was conducted based on the aggregate data available on the Gapminder Website (14). No sampling was done in the current study. The study population consisted of the Iranian population with regard to the availability of the data. The research was performed from 1980 to 2010.

2.1. Data and Variables

The data were extracted from the Gapminder Website where the information about the risk factors of chronic heart disease (CHD) is provided. The variables studied in the research were as follows:

males systolic BP (SBP) (mmHg), females SBP (mmHg), standardized mean age, males BMI, females BMI, male total cholesterol (mM/L), and female total cholesterol (mM/L).

2.2. Data Analysis

The bivariate Pearson correlation was used to estimate the correlation coefficient between SBP and BMI and also between total cholesterol and BMI in both males and females. To compare the mean SBP, mean total cholesterol, and mean BMI of males and females, the independent samples t test was used. To show the trend of male SBP, female SBP, male BMI, female BMI, male total cholesterol, and female total cholesterol within the period of 31 years, the message sequence charts were employed. To analyze data, SPSS version 16 was used; $\alpha = 0.05$ was considered as the level of significance.

3. Results

The average SBP in males was 127.69 mmHg, while it was 126.12 mmHg for females. The mean BP difference between males and females was statistically significant ($P <$

0.001). The mean total cholesterol in males was 4.95 mM/L, while it was 5.10 mM/L in females and the difference between mean values of males and females was statistically significant ($P < 0.001$). Furthermore, the mean BMI in males was 24.35 kg/m² and it was 27.32 kg/m² in females; the difference between the mean values of the 2 genders was statistically significant ($P < 0.001$). The correlation coefficient between the mean BP and the mean BMI in males was -0.595 ($P < 0.001$), and the correlation coefficient between total cholesterol and BMI was 0.476 ($P = 0.007$); the correlations were also significant. The correlation coefficient between the mean BP and the mean BMI for females was -0.905 ($P < 0.001$), while their correlation coefficient between the total cholesterol and BMI was $r = -0.939$ ($P < 0.001$); the correlation in both comparisons was significant.

3.1. The Trend of Males

The mean BP decreased from 1980 to 1997 and then, increased until 2008, and in 2009 and 2010 remained constant. The changes in the mean PB in males and females during the 31 years study are shown in Figure 1.




Figure 1. The Average Blood Pressure (mmHg) in Males and Females from 1980 to 2010

The total cholesterol levels in males reduced from 1980 to 1990 and afterward increased until 2010. The average total cholesterol in females during 3 decades showed an increasing trend.

Figure 2 shows the changes in the mean total cholesterol in males and females from 1980 to 2010.




Figure 2. The Changes in the Mean Total Cholesterol (mM/L) in Males and Females from 1980 to 2010




Figure 3. The Average BMI Variations in Males and Females from 1980 to 2010

The average BMI variations for males and females are shown in [Figure 3](#).

The Average BMI for males and females increased during the study years.

4. Discussion

The mean PB had a decreasing trend from 131.57 to 128.18 mmHg in males and from 128.03 to 125.35 mmHg in females

from 1980 to 2010. A systematic review on adults above 25 years old in 199 countries and territories showed that males and females in Australia, North America, and Western Europe experienced a remarkable decrease in SBP. On the other hand, SBP increased for in males and females in Oceania, Eastern Africa, South and South-East Asia, and just for females in Western Africa. The current study also indicated a decline in the mean SBP all over the world from 130.5 to 128.1 mmHg from 1980 to 2008; a global decrease of SBP also observed by 0.8 mmHg per decade from 1980 to 2008 (15). A study in London regarding the mean SBP was conducted on a population within the age range of 16 to 54 years and reported BP rate of 127.5 mmHg for males and 117.3 mmHg for females. Hence, SBP increased by 0.2 mmHg in females and decreased by 0.5 mmHg in males in 2009, but a significant decrease was observed in BP rate for both genders aged ≥ 55 years (16). There was also a downward trend in BP rate in the Iranian adolescents from 1998 to 2008 (17).

Although the reason for these changes is unknown, many studies showed that the diet can shift the PB curve to a downward direction. The results of a clinical trial showed that a diet rich in fruits, vegetables, and low-fat dairy products with lower saturated and total fat can decrease SBP by 5.5 mmHg and diastolic BP (DBP) by 3 mmHg that was higher than those of the control group (18). Another clinical trial in the US on 412 companies revealed that cutting back on salt (sodium) from high to middle, reduced SBP by 2.1 mmHg in the control and 1.2 mmHg in the DASH group, and also decreased sodium from middle to lower level from 4.6 to 1.7 mmHg in the control and DASH groups, respectively (19). A meta-analysis showed that the modest and long-term reduction in salt intake for 4 weeks reduced the PB dramatically in both hypertensive and normotensive groups. A 9% decline in stroke death and 14% in coronary death in the hypertensive patients as well as 6% in stroke death and 4% in coronary death in the normotensive patients can serve as a proof of this claim (20). It seems that the increase of awareness about healthy diet in recent years greatly influenced females and they paid more attention to their health status compared with males. In the current study, the mean BMI was changed from 23.16 to 25.32 kg/m² in males and from 26.24 to 28.14 kg/m² in females during the study period. A national health survey of people aged 25 - 64 years showed that the mean BMI increased from 25.03 kg/m² in 1999 to 26.14 kg/m² in 2005 (21). It is not only about Iran, but it is a common even all around the world. A systematic review of 9.1 million participants showed that the global mean BMI increased 0.4 kg/m² per decade for males and 0.5 kg/m² for females from 1980 to 2008 (22). In the USA, BMI increased 0.08 kg/m² in males and 0.19 kg/m² in females per year (23). Kather-

ine reported that the mean BMI remained constant as 28.7 kg/m² from 2009 to 2010 and a remarkable increase was shown in males BMI value during the recent 12 years, but indicated no significant changes in that of females (24). Flegal in a 20-year study showed that the mean BMI increased in both white and black females at different income and educational levels (25). Unfortunately, this upward trend progresses much faster in the developing countries than the developed ones; for example, the annual rate of obesity in adult males and females in European and North American countries is 0.25, while it is 2 - 5 times higher in Latin America, Northern Africa, and Asia (26). The changes in food habits, job structure, leisure activities, and exercises are the reasons for such upwards. In addition, McTigue found that the individuals who born in the earlier decades of the century usually have a larger age-specific BMI compared with the ones born in the later decades. The people who born in 1957 had a lower mean BMI than the ones born in 1964 (27).

4.1. Cholesterol

Based on the results of the current study, the total cholesterol increased in both genders, except a decreasing trend in males from 1980 to 1990. Farzadfar focused on the total cholesterol of global population and reported a significant decrease in the total cholesterol level in high-income nations including Australasia, North America, and Western Europe by 0.19 mM/L per decade in males and 0.21 mM/L per decade in females, whereas it increased by 0.08 mM/L per decade in males and 0.09 mM/L per decade in females in East and South-East Asia and the Pacific region (28). Similar studies in the developing countries also reported the same decreasing trend (29, 30). However, an increase in total cholesterol level was observed in Asian countries such as Malaysia, Singapore, Korea, China, and Philippines. Despite the fact that Japan has the least amount of total cholesterol in Asian countries as well as the world, an increase in the total cholesterol of Japanese was observed once in 1958 (31). A reason underlying this increase might be the ascending trajectory of BMI. The second national health and nutrition examination survey (NHANES II) showed that the changes in BMI are accompanied with the changes in total cholesterol (32). A study by Richard also indicated that the total cholesterol and BMI are increased simultaneously (33). Although some other factors contributing total cholesterol increase, apart from BMI increase, the total cholesterol levels can be controlled by cholesterol-lowering drugs and lifestyle modifications, i.e. physical activities and low-fat diets (30). Therefore, controlling any of these factors may induce changes in the general trend.

4.2. Conclusion

Results of the current study indicated that among the risk factors for CVDs, more attention should be paid to PB, and the average PB had a declining trend in recent years due to effective measures and policies. Meanwhile, less attention is paid to other factors such as cholesterol level and BMI; hence, purposive training and interventions to modify lifestyle and nutrition habits are of great importance.

Footnote

Conflict of Interest: Authors declared no conflict of interest.

References

1. Omran AR. The epidemiologic transition. A theory of the epidemiology of population change. *Milbank Mem Fund Q*. 1971;**49**(4):509-38. doi: [10.2307/3349375](https://doi.org/10.2307/3349375). [PubMed: [5155251](https://pubmed.ncbi.nlm.nih.gov/5155251/)].
2. Murray CJ, Lopez AD. Alternative projections of mortality and disability by cause 1990-2020: Global Burden of Disease Study. *Lancet*. 1997;**349**(9064):1498-504. doi: [10.1016/S0140-6736\(96\)07492-2](https://doi.org/10.1016/S0140-6736(96)07492-2). [PubMed: [9167458](https://pubmed.ncbi.nlm.nih.gov/9167458/)].
3. Kannel WB, Vokonas PS. Primary risk factors for coronary heart disease in the elderly: The Framingham Study. In: Wenger NK, Furberg D, Pitt E, editors. *Coronary Heart Disease in the Elderly*. New York: Elsevier; 1986. p. 60-95.
4. Rose G, Shipley M. Plasma cholesterol concentration and death from coronary heart disease: 10 year results of the Whitehall study. *Br Med J (Clin Res Ed)*. 1986;**293**(6542):306-7. doi: [10.1136/bmj.293.6542.306](https://doi.org/10.1136/bmj.293.6542.306). [PubMed: [3089495](https://pubmed.ncbi.nlm.nih.gov/3089495/)].
5. American Heart Association. *Heart Disease and Stroke Statistics: 2005 Update*. Dallas, Tex: American Heart Association; 2004.
6. Mokdad AH, Ford ES, Bowman BA, Dietz WH, Vinicor F, Bales VS, et al. Prevalence of obesity, diabetes, and obesity-related health risk factors, 2001. *JAMA*. 2003;**289**(1):76-9. doi: [10.1001/jama.289.1.76](https://doi.org/10.1001/jama.289.1.76). [PubMed: [12503980](https://pubmed.ncbi.nlm.nih.gov/12503980/)].
7. Garrow J. Importance of obesity. *BMJ*. 1991;**303**(6804):704-6. doi: [10.1136/bmj.303.6804.704](https://doi.org/10.1136/bmj.303.6804.704). [PubMed: [1912920](https://pubmed.ncbi.nlm.nih.gov/1912920/)].
8. The WHO MONICA Project. Geographical variation in the major risk factors of coronary heart disease in men and women aged 35-64 years. *World Health Stat Q*. 1988;**41**(3-4):115-40. [PubMed: [3232405](https://pubmed.ncbi.nlm.nih.gov/3232405/)].
9. Colditz GA, Willett WC, Rotnitzky A, Manson JE. Weight gain as a risk factor for clinical diabetes mellitus in women. *Ann Intern Med*. 1995;**122**(7):481-6. doi: [10.7326/0003-4819-122-7-199504010-00001](https://doi.org/10.7326/0003-4819-122-7-199504010-00001). [PubMed: [7872581](https://pubmed.ncbi.nlm.nih.gov/7872581/)].
10. Field AE, Coakley EH, Must A, Spadano JL, Laird N, Dietz WH, et al. Impact of overweight on the risk of developing common chronic diseases during a 10-year period. *Arch Intern Med*. 2001;**161**(13):1581-6. doi: [10.1001/archinte.161.13.1581](https://doi.org/10.1001/archinte.161.13.1581). [PubMed: [11434789](https://pubmed.ncbi.nlm.nih.gov/11434789/)].
11. Rothwell PM. Limitations of the usual blood-pressure hypothesis and importance of variability, instability, and episodic hypertension. *Lancet*. 2010;**375**(9718):938-48. doi: [10.1016/S0140-6736\(10\)60309-1](https://doi.org/10.1016/S0140-6736(10)60309-1). [PubMed: [20226991](https://pubmed.ncbi.nlm.nih.gov/20226991/)].
12. Vasan RS, Larson MG, Leip EP, Evans JC, O'Donnell CJ, Kannel WB, et al. Impact of high-normal blood pressure on the risk of cardiovascular disease. *N Engl J Med*. 2001;**345**(18):1291-7. doi: [10.1056/NEJMoa003417](https://doi.org/10.1056/NEJMoa003417). [PubMed: [11794147](https://pubmed.ncbi.nlm.nih.gov/11794147/)].
13. Anderson SA, Waters JH. Executive summary from the third report on nutrition monitoring in the United States. *J Nutr*. 1996;**126**(7):iii-x. 1907S-1936S. [PubMed: [8683334](https://pubmed.ncbi.nlm.nih.gov/8683334/)].

14. *Data in Gapminder World*. 2017. Available from: <https://www.gapminder.org/data/>.
15. Danaei G, Finucane MM, Lin JK, Singh GM, Paciorek CJ, Cowan MJ, et al. National, regional, and global trends in systolic blood pressure since 1980: systematic analysis of health examination surveys and epidemiological studies with 786 country-years and 5.4 million participants. *Lancet*. 2011;**377**(9765):568-77. doi: [10.1016/S0140-6736\(10\)62036-3](https://doi.org/10.1016/S0140-6736(10)62036-3). [PubMed: [21295844](https://pubmed.ncbi.nlm.nih.gov/21295844/)].
16. Kumar RA, Dalton AR. The recent English trend in systolic blood pressure and its relationship with age. *Public Health*. 2013;**127**(10):955-7. doi: [10.1016/j.puhe.2012.11.019](https://doi.org/10.1016/j.puhe.2012.11.019). [PubMed: [23290533](https://pubmed.ncbi.nlm.nih.gov/23290533/)].
17. Hosseini-Esfahani F, Mousavi Nasl Khameneh A, Mirmiran P, Ghanbarian A, Azizi F. Trends in risk factors for cardiovascular disease among Iranian adolescents: the Tehran Lipid and Glucose Study, 1999-2008. *J Epidemiol*. 2011;**21**(5):319-28. doi: [10.2188/jea.JE20100162](https://doi.org/10.2188/jea.JE20100162). [PubMed: [21804294](https://pubmed.ncbi.nlm.nih.gov/21804294/)].
18. Appel LJ, Moore TJ, Obarzanek E, Vollmer WM, Svetkey LP, Sacks FM, et al. A clinical trial of the effects of dietary patterns on blood pressure. DASH Collaborative Research Group. *N Engl J Med*. 1997;**336**(16):1117-24. doi: [10.1056/NEJM199704173361601](https://doi.org/10.1056/NEJM199704173361601). [PubMed: [9099655](https://pubmed.ncbi.nlm.nih.gov/9099655/)].
19. Sacks FM, Svetkey LP, Vollmer WM, Appel LJ, Bray GA, Harsha D, et al. Effects on blood pressure of reduced dietary sodium and the Dietary Approaches to Stop Hypertension (DASH) diet. DASH-Sodium Collaborative Research Group. *N Engl J Med*. 2001;**344**(1):3-10. doi: [10.1056/NEJM200101043440101](https://doi.org/10.1056/NEJM200101043440101). [PubMed: [11136953](https://pubmed.ncbi.nlm.nih.gov/11136953/)].
20. He FJ, MacGregor GA. Effect of modest salt reduction on blood pressure: a meta-analysis of randomized trials. Implications for public health. *J Hum Hypertens*. 2002;**16**(11):761-70. doi: [10.1038/sj.jhh.1001459](https://doi.org/10.1038/sj.jhh.1001459). [PubMed: [12444537](https://pubmed.ncbi.nlm.nih.gov/12444537/)].
21. Esteghamati A, Khalilzadeh O, Mohammad K, Meysamie A, Rashidi A, Kamgar M, et al. Secular trends of obesity in Iran between 1999 and 2007: National Surveys of Risk Factors of Non-communicable Diseases. *Metab Syndr Relat Disord*. 2010;**8**(3):209-13. doi: [10.1089/met.2009.0064](https://doi.org/10.1089/met.2009.0064). [PubMed: [20085488](https://pubmed.ncbi.nlm.nih.gov/20085488/)].
22. Finucane MM, Stevens GA, Cowan MJ, Danaei G, Lin JK, Paciorek CJ, et al. National, regional, and global trends in body-mass index since 1980: systematic analysis of health examination surveys and epidemiological studies with 960 country-years and 9.1 million participants. *Lancet*. 2011;**377**(9765):557-67. doi: [10.1016/S0140-6736\(10\)62037-5](https://doi.org/10.1016/S0140-6736(10)62037-5). [PubMed: [21295846](https://pubmed.ncbi.nlm.nih.gov/21295846/)].
23. Shah M, Hannan PJ, Jeffery RW. Secular trend in body mass index in the adult population of three communities from the upper mid-western part of the USA: the Minnesota Heart Health Program. *Int J Obes*. 1991;**15**(8):499-503. [PubMed: [1938092](https://pubmed.ncbi.nlm.nih.gov/1938092/)].
24. Flegal KM, Carroll MD, Kit BK, Ogden CL. Prevalence of obesity and trends in the distribution of body mass index among US adults, 1999-2010. *JAMA*. 2012;**307**(5):491-7. doi: [10.1001/jama.2012.39](https://doi.org/10.1001/jama.2012.39). [PubMed: [22253363](https://pubmed.ncbi.nlm.nih.gov/22253363/)].
25. Flegal KM, Harlan WR, Landis JR. Secular trends in body mass index and skinfold thickness with socioeconomic factors in young adult women. *Am J Clin Nutr*. 1988;**48**(3):535-43. doi: [10.1093/ajcn/48.3.544](https://doi.org/10.1093/ajcn/48.3.544). [PubMed: [3414568](https://pubmed.ncbi.nlm.nih.gov/3414568/)].
26. Popkin BM, Gordon-Larsen P. The nutrition transition: worldwide obesity dynamics and their determinants. *Int J Obes Relat Metab Disord*. 2004;**28** Suppl 3:S2-9. doi: [10.1038/sj.ijo.0802804](https://doi.org/10.1038/sj.ijo.0802804). [PubMed: [15543214](https://pubmed.ncbi.nlm.nih.gov/15543214/)].
27. McTigue KM, Garrett J, Popkin B. The natural history of obesity: weight change in a large US longitudinal survey. *Ann Intern Med*. 2002;**136**:857-64.
28. Farzadfar F, Finucane MM, Danaei G, Pelizzari PM, Cowan MJ, Paciorek CJ, et al. National, regional, and global trends in serum total cholesterol since 1980: systematic analysis of health examination surveys and epidemiological studies with 321 country-years and 3.0 million participants. *Lancet*. 2011;**377**(9765):578-86. doi: [10.1016/S0140-6736\(10\)62038-7](https://doi.org/10.1016/S0140-6736(10)62038-7). [PubMed: [21295847](https://pubmed.ncbi.nlm.nih.gov/21295847/)].
29. Johnson CL, Rifkind BM, Sempos CT, Carroll MD, Bachorik PS, Briefel RR, et al. Declining serum total cholesterol levels among US adults. The National Health and Nutrition Examination Surveys. *JAMA*. 1993;**269**(23):3002-8. doi: [10.1001/jama.269.23.3002](https://doi.org/10.1001/jama.269.23.3002). [PubMed: [8501842](https://pubmed.ncbi.nlm.nih.gov/8501842/)].
30. Carroll MD, Lacher DA, Sorlie PD, Cleeman JI, Gordon DJ, Wolz M, et al. Trends in serum lipids and lipoproteins of adults, 1960-2002. *JAMA*. 2005;**294**(14):1773-81. doi: [10.1001/jama.294.14.1773](https://doi.org/10.1001/jama.294.14.1773). [PubMed: [16219880](https://pubmed.ncbi.nlm.nih.gov/16219880/)].
31. Khoo KL, Tan H, Liew YM, Deslypere JP, Janus E. Lipids and coronary heart disease in Asia. *Atherosclerosis*. 2003;**169**(1):1-10. doi: [10.1016/S0021-9150\(03\)00009-1](https://doi.org/10.1016/S0021-9150(03)00009-1). [PubMed: [12860245](https://pubmed.ncbi.nlm.nih.gov/12860245/)].
32. Denke MA, Sempos CT, Grundy SM. Excess body weight. An underrecognized contributor to high blood cholesterol levels in white American men. *Arch Intern Med*. 1993;**153**(9):1093-103. doi: [10.1001/archinte.1993.00410090045006](https://doi.org/10.1001/archinte.1993.00410090045006). [PubMed: [8481076](https://pubmed.ncbi.nlm.nih.gov/8481076/)].
33. Hershcopf RJ, Elahi D, Andres R, Baldwin HL, Raizes GS, Schocken DD, et al. Longitudinal changes in serum cholesterol in man: an epidemiologic search for an etiology. *J Chronic Dis*. 1982;**35**(2):101-14. doi: [10.1016/0021-9681\(82\)90111-4](https://doi.org/10.1016/0021-9681(82)90111-4). [PubMed: [7056835](https://pubmed.ncbi.nlm.nih.gov/7056835/)].