

CAN RELAXATION BE USED TO ACHIEVE WEIGHT CONTROL IN YOUTHS?

R Kelishadi MD⁽¹⁾, M Beizaei MS⁽²⁾, M Hashemipour MD⁽³⁾, M Salek MD⁽⁴⁾
A Gholi Amini MD⁽⁵⁾, M Soghrati MD⁽⁶⁾, S Ghatreh Samani MD⁽⁷⁾

Abstract

INTRODUCTION: Childhood obesity is a cultural and medical problem and is usually a refractory to treatment. The objective of this study was to assess the effects of parental therapy behavior and relaxation methods on children in controlling childhood obesity.

METHODS: In this randomized controlled clinical trial, 90 obese children aged 10-17 years were randomly divided into three groups of equal number. All three groups received similar diet and exercise recommendations. The first group of children attended 15 relaxation sessions. Parents of the second group participated in 15 behavior therapy sessions. The third group was considered as control. Data were analyzed by SPSS 13. Mean changes of weight, BMI and WC were calculated.

RESULTS: Mean BMI decreased in all three groups after the intervention. This decrease was more remarkable, but not significant in the first group. Mean weight decreased in the first group but increased in the second group. Mean WC decreased in all three groups. This decrease was more obvious, but not significant in the first group.

CONCLUSIONS: Stress leads to sympathetic system arousal which is associated with many pathologic conditions. Elicitation of relaxation response can help individuals embrace healthy lifestyle choices. Relaxation and parent behavior therapy can be considered as useful methods of controlling childhood obesity.

Keywords: Obesity, children and adolescents, relaxation, behavior therapy, parents.

ARYA Journal, 2006, 2(2): 92-97

Introduction

Childhood obesity is a cultural and medical problem. Although its prevalence is higher in developed countries, it is rapidly increasing in developing, and even in the third world countries.¹⁻⁵ Recent studies in the Iranian society have shown a rapid increase in the childhood obesity rate.^{6,7}

The early and late complications of childhood obesity confirm the importance of weight control from childhood. Lifestyle change is a very effective method to this end, but requires good training and support. In addition, it is difficult to maintain ideal weight after weight loss. In many cases, it is very difficult to draw the children's and adolescents' cooperation in weight-loss programs, so many families look for safe medications to control the weight of their children.

Pharmacologic therapy might have side effects and our previous study on the effects of herbal mixtures on controlling childhood obesity suggested that using placebo and herbal mixtures decreased the compliance of children and their families in following diet and exercise recommendations.⁸ Relaxation can be effective as an adjunct therapy with other treatments. Parent behavior therapy might also have a positive effect on weight control of children and adolescents.^{9,10}

An association between stress and diseases has been examined in recent studies.¹¹⁻¹³

Such an association apparently exists, however, it is complex and specific clinical implications should be further investigated. In general, stress impacts on the immune, circulatory, and nervous systems, hence it

(1) Roya Kelishadi MD. Associate Professor of Pediatrics, Head, Preventive Pediatric Cardiology Department, Isfahan Cardiovascular Research Center (ICRC)

(2) Maryam Beizaei MS. Research Assistant, Isfahan Cardiovascular Research Center (ICRC)

(3) Mahin Hashemipour MD. Professor, Pediatric Endocrinologist, Isfahan University of Medical Sciences, Iran

(4) Mehdi Salek MD. Assistant Professor, Pediatric Endocrinologist, Isfahan University of Medical Sciences, Iran

(5) Abbas Gholi Amini MD. Assistant Professor, Pediatric Hematologist, Isfahan University of Medical Sciences, Iran

(6) Mojgan Soghrati MD. Research Assistant, Isfahan Cardiovascular Research Center (ICRC)

(7) Shohreh Ghatreh Samani MD. Research Assistant, Isfahan Cardiovascular Research Center (ICRC)

Corresponding author: Roya Kelishadi

Date of submission: June 2, 2006

Date of acceptance: September 24, 2006

may play a role in immunological, cardiovascular and neurodegenerative diseases, as well as mental disorders.^{12,13}

Stress can exert either ameliorating or deleterious effects, depending on a multitude of factors (e.g. individual, endogenous, or exogenous elements).¹¹⁻¹³ However, in the clinical and actual medical setting, the negative influences of stress on health seem to predominate.^{12,13}

Therapeutic use of the relaxation response has been documented in several studies.^{14,15} Mind/body practices that elicit the relaxation response (RR) have been practiced for thousands of years to promote health and wellbeing.¹⁶ Numerous mind/body approaches can elicit the RR.^{16,17} The RR can be elicited as individuals repeat a word, sound, phrase, or prayer, or as they focus on their breathing and disregard everyday intrusive thoughts.¹⁸ RR is described as a coordinated physiological response that is characterized by decreased arousal, diminished heart rate, respiratory rate and blood pressure, associated with a state of "wellbeing".¹⁸⁻²⁰

The physiological responses of relaxation occur in the opposite direction to those of the stress response, described as the "fight or flight" response and the "general adaptation response" to stress.²¹

Clinically, the RR has been shown to counteract the negative effects of long-term stress. The RR is often used as an adjunct to medical treatment, in conditions that are caused or exacerbated by stress.²³

These conditions represent a broad range of physiological systems such as, hypertension,²⁴ myocardial ischemia,²⁴ anxiety,²⁶ psychosomatic complaints,^{26,27} insomnia,²⁹⁻³¹ headache,^{32,33} back/neck pain,³⁴ chronic pain,³⁵⁻³⁶ musculoskeletal disorders,³⁷ etc. The clinical effects of the RR have been shown to improve the outcomes of medical and surgical treatment.³⁸⁻⁴⁰

Despite these clear physiological and clinical observations, the underlying mechanisms of the RR remain undefined. Relaxation therapy may also be of benefit in improving weight loss. The majority of studies have assessed behavioral and cognitive behavioral weight reduction strategies.

Cognitive therapy, psychotherapy, relaxation therapy and hypnotherapy have been assessed in a small number of studies. The objective of this study was to evaluate the effects of parental therapy behavior and relaxation methods on children in controlling childhood obesity.

Materials and methods

This randomized controlled clinical trial was carried out in 2005 on 90 children and adolescents aged 10-17 years who had been diagnosed as obese [body mass index (BMI) > age- and gender-specific 95th percentile] and referred by schools and/or physicians to the Metabolic Syndrome Clinic of preventive Pediatric Cardiology Department, Isfahan Cardiovascular Research Center (a WHO collaborating Center). The subjects were randomly divided into three equal groups of 30 by using the table of randomized numbers and the simple random method.

The Ethics Committee of Isfahan Cardiovascular Research Center approved the study. Verbal assent was obtained from children and informed written consent from their patients.

All subjects were examined by the same physician. Eligibility criteria were age 10-17 years and a BMI greater than age- and gender-specific 95th percentile, no physical disorders leading to exercise inability, and living with biological parents. Height (Ht) and weight (Wt) were measured twice to ± 0.2 cm and to ± 0.2 kg, respectively, with subjects being barefoot and lightly dressed; the averages of these measurements were recorded. BMI (weight in kilograms divided by the square of height in meters) was calculated.

Waist circumference (WC) was measured with non-elastic tape at a point midway between the lower border of the rib cage and the iliac crest at the end of normal expiration. All groups received similar diet and exercise recommendations. Children and adolescents in the first group participated in 15 sessions of relaxation. In the second group, parents of obese children participated in 15 sessions of behavior therapy. The third group was considered as control. Techniques and methods employed in the first and second groups were as follows:

1. Relaxation techniques, consisting of body imaging (negative minds were converted to positive ones)
2. Self imaging for organization of mind/body.
3. Methods for decreasing anxiety.
4. Yoga techniques for activation of muscles and repairing body structure.
5. Breathing techniques.


The following two components necessary to elicit the relaxation response were considered as well-focused awareness on a thought, work, sound, or muscular movement, and passive disregard of distracting thoughts.



Overweight or obese individuals may benefit from psychological interventions, particularly behavioral and cognitive-behavioral strategies to enhance weight reduction. These strategies are particularly useful when combined with dietary and exercise strategies. Behavioral therapy was found to result in significantly greater weight reduction than placebo when assessed as a stand-alone weight loss strategy. When behavioral therapy was combined with a diet/exercise approach and compared with diet/exercise alone, the combined intervention resulted in greater weight reduction.³⁹ The mean changes of weight, BMI and WC were compared between the three groups. Data were analyzed by SPSS13. Changes in mean difference were analyzed by Pearson Correlation method; changes in data in all groups were analyzed by analysis of

variance (ANOVA). The study received the approval of the Ethics Committee of Isfahan Cardiovascular Research Center.

Results

In this study, the mean of BMI decreased in all groups after intervention, but those who practiced relaxation methods as an adjunct therapy to diet and exercise displayed more notable, yet insignificant decrease in their BMI (Figure 1). After the intervention, mean weight decreased in the first group (relaxation) but increased in the second group (Figure 2). Mean WC decreased after intervention in all three groups; although the decrease in WC was greater in the relaxation group, the difference was not significant (Figure 3).





Discussion

Our findings demonstrate that adjunct methods such as relaxation and parent behavior therapy can help to decrease BMI in obese children and adolescents. This decrease was more remarkable, but not significant in the group attending relaxation sessions. There was also a decrease in WC in all three groups and was most notable in the first group, however, it was not significant. This group reported peace of mind and greater tendency towards wellbeing, which apparently helped decrease body weight.

Although stress is a well known phenomenon, it is an elusive concept to define. Stress varies from individual to individual and within the same individual, stress triggers vary from day to day. It is also important to acknowledge that not all stress is bad. A certain amount of stress creates interest and challenge in life. Too much stress, however, leads to negative consequences and adversely affects health.⁴⁰ Sympathetic nervous system (SNS) arousal in response to stress has been linked to increase in CVD and their risk factors.^{41,42}

Our intervention included an integrated psychosocial, non-pharmacologic approach to stress management as part of a multiple risk factor reduction approach.⁴³ Benson et al. first described the physiology of relaxation response. Elicitation of relaxation response is associated with decreased SNS activity acutely, and in longer term has been shown to decrease end-organ responsiveness.¹⁸ In addition to these physiological changes, physiological and behavioral changes may also occur as a result of relaxation response (RR).

Patients who regularly elicit the relaxation response report that they can focus attention more easily, and appraise attitudes in a more conscious, objective way. They experience an opportunity to test new possibilities and embrace healthy lifestyle choices. This process of behavioral changes begins when the patient clarifies core values and makes a behavioral plan that is aligned with his or her core values. Many techniques can be used to elicit the RR. It is best to teach a variety of techniques so that the patients will be able to identify the technique(s) that work best for them. We advised patients to elicit the RR, once a day for 20 minutes.⁴³

In a previous study on the effect of relaxation on anxiety and learning improvement, the groups practicing relaxation methods with or without imagery, experienced a more peaceful state while studying or during exams.⁹

Other researches also showed that parent behavior therapy impacts positively to help obese children and adolescents lose their weight.¹⁰

Our study shows that parent behavior therapy contributes to decreased BMI and WC. All mothers who received behavior therapy stated that their behavior modification and relaxed attitude led to better control and decrease in their children's weight.

Our limitations were the short duration of relaxation and behavior therapy sessions (3 months) and the small sample size (30 subjects in each group). In general, our results indicate that relaxation practices, as an adjunct to diet and exercise, are useful in controlling generalized and abdominal obesity. Thus more prolonged studies (6-12 months) with larger sample size are warranted to obtain more conclusive results. This study was approved by High Council of Isfahan Cardiovascular Research Center.

Acknowledgement

This study was funded by Isfahan Cardiovascular Research Center.

References

1. Schonfeld-Warden N, Warden CH. Pediatric obesity: an overview of etiology and treatment. *Pediatr Clin of North Am*1997;44:339-57.
2. Popkin BM, Paeratakul S, Zhai F, Ge K. Dietary and environmental correlates of obesity in a population study in China. *Obes Res* 1995;2:135.
3. Booth ML, Macaskill P, Baur LA. Sociodemographic distribution of measures of body fatness among children and adolescents in New South Wales, Australia. *Int J Obes Relat Metab Disord* 1999;23:456-62.
4. MUSAIGER AO, MATTER AM, ALEKRI SA, MAHDI AR. Prevalence of obesity in Bahrain. *Nutr Health* 1993;9:25-8.
5. Dhurandhar NV, Kulkarni PR. Obesity among secondary school prevalence of obesity in Bombay. *Int J Obes Relat Metab Disord* 1992;16 :367-70.
6. Kelishadi R, Hashemipour M, Sarrafzadegan N, Amiri M. Trend of atherosclerosis risk factors in Isfahan. *Asian cardiovascular thorac Ann* 2001;9:36-40.
7. Kelishadi R, Hashemipour M, Sarrafzadegan N, et al. Obesity and associated modifiable risk factors in Iranian adolescents. *Pediatr Int* 2003;45:435-42.
8. Hashemipour M, Kelishadi R, Asgari S, Talaei M, Salek M, Mohammadifard N, et al. Efficacy of herbal mixtures in childhood obesity control. *J Ped Neonat* 2006;3(2):4-8.
9. Goli F, Amini A, Golabadi M. Study of effect of short time relaxation program with or without body mind self-imagery, on amount of anxiety and student's memory function in elementary levels. *Iran J Higher Health* 2003;2(2-3):156-68.
10. Wrotniak BH, Epstein LH, Paluch RA, Roemmich JN. Parent weight change in family-based behavioral obesity treatment. *Arch Pediatr Adolesc Med* 2004; 158(4): 342-7.
11. Esch T: Health in Stress: Change in the stress concept and its significance for prevention, health and lifestyle. *Gesundheitswesen* 2002;64:73-81.

12. Esch T, Stefano GB, Fricchione GL, Benson H. An overview of stress and its impact in immunological disease. *Mod Asp Immunobiol disease. Med Asp Immunobiol* 2002;2:187-92.
13. Esch T, Stefano GB, Fricchione GL, Benson H. The role of stress in neurodegenerative diseases and mental disorders. *Neuroendocrinol Lett* 2002;23:199-208.
14. Komaroff AL. *mind/body medicine: A special health report.* Boston Harvard Health Publication 2001.
15. Willett WC. Balancing life-style and genomics research for disease prevention. *Science* 2002;296:695-8.
16. Wolsko PM, Eisenberg DM, Davis RB, Phillips RS. Use of mind-body medical therapies. *J Gen Intern Med* 2004;19(1):43-50.
17. Barnes PM, Powell-Griner E, McFann K, Nahin RL. Complementary and alternative medicine use among adults: United States, 2002. *Adv Data* 2004;(343):1-19.
18. Benson H, Beary JF, Carol MP. The relaxation response. *Psychiatry* 1974;37(1):37-46.
19. Benson H. *The relaxation response: William Morrow, New York, 1975.*
20. Wallace PK, Benson H, Wilson AF. A wakeful hypometabolic physiologic state. *Am J physiol* 1971;221(3):795-9.
21. Selye H. *The stress of life.* MC Graw-Hill, New York, 1956.
22. Sternberg E, Gold PW. The mind-body interaction in disease. *Scientific Amer* 1997;7(1):8-15.
23. Benson H et al: Decreased blood-pressure in pharmacologically treated hypertensive patients who regularly elicited the relaxation response. *Lancet* 1974;1(7852):289-91.
24. Blumenthal J et al. Stress management and exercise training in cardiac patients with myocardial ischemia : effects on prognosis and evaluation of mechanisms. *Arch Intern Med.* 1997;157:2213-23.
25. Miller J, Fletcher KE, Kabat-Zinn J. Three year follow-up and clinical. Implications of a mindfulness meditation based stress reduction intervention in the treatment of anxiety disorders. *Gen Hosp Psychiatry*, 1995;17:192-200.
26. Hellman CJ, Budd M, Borysenko J, McClelland DC, Benson H. A study of the effectiveness of two group behavioral medicine interventions for patients with psychosomatic complaints. *Behav Med* 1990;16(4): 165-73.
27. Nakao M, Myers P, Fricchione G, Zuttermeister PC, Barsky AJ, Benson H. Somatization and symptom reduction through a behavioral medicine intervention in a mind/body medicine clinic. *Behav Med* 2001;26(4): 169-76.
28. Jacobs GD, Roisenberg PA, Friedman R, Matheson J, Peavy GM, Domar AD, et al: Multifactor behavioral treatment of chronic sleep-onset insomnia using stimulus control and the relaxation response. A preliminary study. *Behav Modif* 1993;17(4):498-509.
29. Morin C, Culbert J, Schwartz S: Nonpharmacological interventions for insomnia: a meta-analysis of treatment efficacy. *Am J Psychiatry* 1994;151:1172-80.
30. Benson H, Klemchuk HP, Graham JR: The usefulness of the relaxation response in the therapy of headache. *Headache* 1974;14(1):49-52.
31. Benson H, Malvea BP, Graham JR: Physiologic correlates of meditation and their clinical effects in headache: an ongoing investigation. *Headache* 1973;13(1): 23-24.
32. Astin JA: Mind-body therapies for the management of pain. *Clin J Pain* 2004,20(1):27-32.
33. Caudill M et al: Decreased clinic utilization by chronic pain patients after behavioral medicine intervention. *Pain* 1991;45(3): 334-5.
34. Luskin FM, Newell KA, Griffith M, Holmes M, Telles S, DiNucci E, Marvasti FF, Hill M, Pelletier KR, Haskell WL. A review of mind/body therapies in the treatment of musculoskeletal disorders with implications for the elderly. *Altern Ther Altern Ther Health Med.* 2000;6(2):46-56.
35. Astin JA et al: Psychological interventions for rheumatoid arthritis: a meta-analysis of randomized controlled trials. *Arthritis and Rheumatism* 2002;47(3):291-302.
36. Dreher H. Mind-body Interventions for surgery: evidence and exigency. *Adv Mind-Body Med* 1998;14:207-22.
37. Wientjes KA. Mind-body techniques in wound healing. *Ostomy Wound Manage* 2002;48(11):62-7.
38. Lang E, Benotsch EG, Fick LJ, Lutegendorf S, Berbaum ML, Berbaum KS, et al: Adjunctive non-pharmacological analgesia for invasive medical procedures: a randomized trial. *Lancet* 2000;355:1486-90.
39. Shaw K, O'Rourke P, Mar C, Kenardy J. Psycho logical interventions for overweight or obesity *Cochrane Database Syst Rev* 2005;18(2).
40. Well- Federman CL, Stuart EM, Deckro JP, Mandel CL, Baim M, Medich C. the mind- body connection: the psychophysiology of many traditional nursing intervention. *Clin Nurs Specialist* 1995;9:59-66.
41. Mittleman MA, Maclure M, Sherwood JB, Murly RP, Tofler GH, Jacobs SC, et al. Triggering of acute myocardial infraction onset by episodes of anger. *Circulation* 1995;22:1720-5.
42. Muller JE, Abela GS, Nesto RW, Tofler GH. Triggers, acute risk factors and vulnerable plaques; the lexicon of new frontier. *J AM Coll Cardiol* 1999;13:809-13.
43. Stuart Shor EM. *Stress Management In: Nanatte K, Wenger L, et al. Cardiac rehabilitation: A Guide to practice in 21st centuries.* Korcel Pekker 1999;287-94.