

RESEARCH ARTICLE

The effectiveness of life skills training on the quality of life of mothers of children with hearing impairment

Khalil Kakavandi¹, Rogayeh Koochi², Setare Shojaee¹, Guita Movallali^{2*}, Shadi Norozali³

¹- Department of Special Education, School of Education and Psychology, Shiraz University, Shiraz, Iran

²- Pediatric Neurorehabilitation Research Center, University of Social Welfare and Rehabilitation Sciences, Tehran, Iran

³- Department of Rehabilitation Counseling, University of Social Welfare and Rehabilitation Sciences, Tehran, Iran

Received: 20 Sep 2016, Revised: 23 Oct 2016, Accepted: 23 Nov 2016, Published: 15 Jan 2017

Abstract

Background and Aim: Hearing impairment is one of the common disabilities quality of life of mothers of children with hearing impairment are often lower than others. This study aimed to evaluate the effects of life skills training (LST) on the quality of life of mothers who have children with hearing impairment.

Methods: An experimental research method (pretest, post-test with control group design) was followed in this study. The statistical population comprised all mothers who have children with hearing impairment and lived in Shiraz, Iran. Of them, 36 mothers were selected through purposeful sampling method and randomly divided into two 18-member groups (experimental and control groups). The pretest was conducted on both groups. LST program was performed in the experimental group in 12 sessions, and the control group did not receive any intervention. Post-test was conducted on both groups. The quality of life scale (QOLS) was used to assess each mother's quality of life.

Results: Life skills training causes significant improvement in the mean scores of the mothers'

physical health, mental health, social relationships and ultimately quality of life in the experimental group ($p < 0.001$).

Conclusion: According to this study, LST is effective in improving the quality of life of mothers who have children with hearing impairment. This program so it can be considered as a useful tool in this field.

Keywords: Life skills training; quality of life; children with hearing impairment

Introduction

Hearing loss is a prevalent and disabling chronic condition that can impair communication, quality of life, and health [1]. Becoming aware of their child's hearing impairment, mothers show different reactions. Some parents may feel disappointed, whereas others become suspicious [2]. The sources of this stress may be concerns over the future of the child, his or her behavioral problems and disabilities, and how the child will cope with future problems. Other concerns are related to changes in family relationships, heavy economical costs, concerns over non-fulfillment of dreams, loss of recreation, compassionate behaviors of other people or negative attitudes of society, lack of information, restricted access to services and facilities or conflicts about rearing [3].

* **Corresponding author:** Pediatric Neurorehabilitation Research Center, University of Social Welfare and Rehabilitation Sciences, Daneshjoo Blvd., Evin, Tehran, 1985713834, Iran. Tel: 009821-22180042, E-mail: drgmovallali@gmail.com

All these concerns exert extra psychological pressure on parents and affect their personal lives. Based on systemic approaches, family is not a static organization, but a dynamic one in which the members influence each other. Thus, the presence of a disabled child and its related stress can affect the whole family, and their interactions and relationships with the hearing-impaired child. The unwise resolution of these unpleasant effects can impose irreparable psychic damages on both parents and other family members, especially the one with hearing loss [4].

In addition, having a child with hearing impairment can impact the quality of life of the parents. Quality of life is a multidimensional construct that includes evaluation of at least four main aspects of emotional well-being, physical health, social function, and spirituality [3]. Dempsey et al. considered the quality of life as a social, emotional, psychological, and physical function [5]. They reported that stress evaluation is the strongest predictor variable of quality of life of parents who have children with attention deficit hyperactivity disorder. Botvin and Kantor argued that the effects of any stressor event are based on how people deal with that, thus appropriate coping skills to maintain family and quality of life of every member of the family is essential. Parents of disabled children can control and manage their condition through appropriate training and learning programs so as to maintain emotional stability in their lives. One of these programs is life skills training (LST), which can lead to appropriate changes in attitudes and values, strengthen behaviors toward health problems and barriers, promotes mental health [6], and enables people to cope with difficult life situations [7]. Studies on the effectiveness of teaching life skills on mental health have indicated that training these skills impacts the quality of life of individuals. For example, Malouff et al. [8] showed that teaching life skills had significant effects on physical and mental health problems of individuals. Kazemi et al. [9] ran a training program to support parents who had children with disabilities. Their results showed that LST was a beneficial

intervention for health and lifestyle problems and could ameliorate the depressed mood. Yankey and Biswas also reported that LST was effective in reducing stress in young Tibetan refugees [10].

The results of multivariate analysis of variance revealed that LST in students with mathematical problems had positive effects on social competence and its components. Pakdaman Savoji and Ganji [11] found that there was a significant effectiveness of LST program on the mental health of students, also the study girls received more benefit from the program than boys.

Based on the theoretical foundation and the mentioned research background, it seems that one way to reduce stress of mothers of children with hearing impairment is to help them develop skills needed to create a satisfying life. One of the achievements in this field is LST, which is inscribed by WHO as an approach to prevent and improve the mental health of the population. This increases the individuals' ability in efficient adaptation to environment and facing challenges [12]. Therefore, this study aimed to evaluate the effect of LST on the quality of life of mothers who have children with hearing impairment.

Methods

This study had a quasi-experimental design with pretest and post-test along with a control group. The study population comprised mothers of children (age 2 to 8 years) with hearing impairment who visited rehabilitation centers in Shiraz, Iran. Sampling was performed using the purposeful convenience sampling method. In the beginning, questionnaires related to the quality of life components were randomly distributed among 100 mothers in the rehabilitation centers. After collecting the questionnaires and scoring them, 36 mothers, whose quality of life scores were below average and were willing to participate in the second phase of the study with LST workshops, were selected. They were randomly placed into experimental group (n=18) and control group (n=18). The study instrument was the WHO quality of life short scale (WHOQOL-BREF).

Table 1. Programs content of life skills training (LST) [15]

Session one: Getting to know each other and establish a good relationship, describing the workshops laws, life skills training, introducing the life skills and quality of life, the definition of quality of life and improve the quality of life.
Session two: Stressors coping skills: definition of stress, coping skills and effective factors related to deal with stressors, coping types [problem-focused, emotion-focused and inconsistent-focused coping] and strategies against stress.
Session three: Coping with emotions skills: introducing the skills to deal with emotions. In this session, the main discussion was about the emotions, all kinds of emotions and their impact on social relationships and unpleasant emotions control strategies and to train these skills, mothers were asked to share their experiences of family member's emotional responses during a pleasant and unpleasant event and describe their sense in an emotional situations.
Session four: Problem solving skills: during this session, mothers became familiar with problem-solving skills. Also, about the ability to solve problems or cope with the problem and its impact on mental health promotion were discussed. The process of problem solving [problem type, finding different solutions and making decisions about the best solution was spoken and in this regard, some assignments were devolved to mothers].
Session five: Decision-making skills: during the session, about the importance of decision-making skills to achieve the objectives, methods of decision-making and its different stages and critical situations of life were discussed. Then mothers were given assignments.
Session six: Self-awareness: during this session, the definition of self-awareness, self-awareness component, contributing factors in the growth of self-awareness and self-growth obstacles discussed. Then assignments were given to mothers.
Session seven: Empathy skills: during the session, about the definition of empathy, empathy skills, prerequisites for empathy, ways to achieve empathy and other empathy-related skills discussed and assignments were given to mothers.
Session eight: Effective communication skills: during the session, about the importance of effective communication in human flourishing and communication barriers discussed and mothers were asked to describe an experience about the barriers to communication and share them with each other.
Session nine: Interpersonal communication skills, interpersonal skills and its importance in mental health, social communication skills, interpersonal relations and the establishment of security, peace, and respect were discussed in this session, and then mothers were given assignments.
Session ten: Conflict resolution skills [critical thinking]: this meeting was about the conflicts in the personal lives and their preventing strategies and then, mothers were given assignments.
Session eleven: Creative thinking skills: creative thinking was defined in this meeting and mothers were trained to express material thinking and creative thinking and how to raise and nurture their creative thinking.
Session twelve: Summary of educational issues.

The scale has 26 questions that assess four domains of quality of life: physical health, mental health, living environment, and relationships with others. Nejat et al. have reported descriptive reliability coefficient and internal consistency of the instrument as 87.0 and 84.0, respectively, in a sample of 302 students from Shiraz University [13]. Pretest reliability coefficient was obtained as 0.67, and the result of the concurrent validity of the scale of public health was satisfactory. Moreover, their pilot study showed that the Persian-translated questionnaire was valid and reliable [13]. Factor analysis of the Iranian version led to three subscales, unlike the original version, which had four

subscales, as living environment and relationships with others together make up a subscale [14]. The LST program consisted of skills related to group decision-making, problem-solving, effective communication, interpersonal relationship, self-awareness, empathy with others, coping with emotions and stressors, critical thinking, and creative thinking. Both groups completed WHOQOL-BREF, and then, the 12-step model of LST (Table 1) was taught to the experimental group in 12 weekly sessions of 1.5 hours each. In the end, both groups were asked to complete the WHOQOL-BREF scale. The demographic data included mother's education (elementary, cycle, diploma, associate,

Table 2. Distribution of absolute and relative frequencies of demographic characteristics in both experimental and control groups

Variable		Group				p
		Experimental		Control		
		N	Percent	N	Percent	
Mother's education	Elementary	2	11.1	2	11.1	0.79
	Cycle	5	27.8	5	27.8	
	Diploma	5	33.3	6	33.3	
	Associate	5	11.1	2	11.1	
	Bachelor of Science	1	11.1	2	11.1	
	Master of Science and higher	0	5.6	1	5.6	
Employment status	Employed	1	5.6	4	22.2	0.15
	Housewife	17	94.4	14	77.8	
Gender	Male	11	61.1	13	72.2	0.49
	Female	7	38.9	5	27.8	
Birth order	First	10	55.6	14	72.2	0.77
	Second	5	27.8	1	11.1	
	Third	3	16.7	3	16.7	
Type of pregnancy	Programmed	12	66.7	13	72.2	0.32
	Undesirable	6	33.3	5	27.8	
Income	Low	8	44.4	5	27.8	0.24
	Intermediate	8	44.4	9	50.0	
	High	2	11.1	4	22.2	
The mean age of mothers		18	29.05	18	29.66	
The mean age of children		18	4.55	18	5.25	

bachelor, and master and higher), employment status (employed/housewife), gender (male/female), birth order (first, second and third), type of pregnancy (programmed/undesirable), income level (low, intermediate, and high), age of mothers, and age of children. The demographic data of both groups were compared using the Chi-square and independent t-test. To determine whether the obtained changes are stati-

stically significant, the analysis of covariance was used. Before regression analysis of covariance, homogeneity of variances (Leven test) were conducted. The normal distribution of the dependent variables was confirmed with Kolmogorov-Smirnov test.

Results

There were no significant differences in mothers

Table 3. The mean and standard deviation of quality of life in control and experimental groups

Variable	Group	Mean (SD)	
		Pretest	Post-test
Quality of life	Experimental	65.88 (12.30)	74.05 (15.50)
	Control	74.05 (15.14)	73.33 (15.81)

education, employment status, birth, order, type of pregnancy, income level, age of mothers, and age of children between groups ($p > 0.05$, Table 2).

As it can be seen in Table 3, there were differences between two groups in pretest and post-test.

Results of covariance analysis after statistically controlling for the effect of pretest showed statistically significant difference in test scores between the experimental and control groups in physical health ($F=1.37$, $p < 0.001$), mental health ($F=2.33$, $p < 0.001$), social relationships ($F=1.66$, $p < 0.001$) and finally quality of life ($F=35.67$, $p < 0.001$). Thus, the intervention was effective in improving the quality of life of mothers who have children with hearing impairment.

Discussion

Our results showed that the mean score of quality of life in dimensions of physical health, psychological health, social relationships, and environmental health increased in the experimental group. The result is consistent with the results of previous studies [9-12,16-19]. It seems that, by making changes in self-awareness, understanding feelings and thoughts, communication styles, behavioral tendencies and feelings of self-efficacy and self-sufficiency in the mothers, the LST program can lead to behavioral changes and ultimately, enhance their quality of life. In addition, the results showed that LST had a significant impact on the quality of life in the experimental group.

This intervention has increased physical activity and physical health and reduced physical

pain in the experimental group. This finding is consistent with the findings of other studies [20-23]. It was also found that LST reduced psychological problems and increased psychological health of participants. This result is in line with that of other studies [24-26]. To explain these results, it can be said that, at first, LST program topics such as expecting opportunities, attending imbalances, recognizing feelings, classification issues, and peace of mind, has led to the reduction of negative thoughts, worries, and negative emotions and improvement in mental and physical health [23]. In fact, mothers in the experimental group received benefits from LST such as in problem-solving skills, decision-making and anger management, effective communication, and the like and learned to deal more efficiently with personal, social, and environmental problems after better defining and recognizing the moot points. Several studies have shown that LST is effective in improving mental and physical health, strengthening self-confidence and interpersonal relationships, and preventing mental, behavioral, and social problems [27]. In LST, new techniques for each skill are introduced and exercised by participants for an immediate, positive, and significant impact. This effect is useful even if it is short-term. Practical training ensures the participant of his/her mood changes by repetition and practice. Increasing in the reliability of these techniques encourages repetitive practice, and eventually the more usefulness of this method [20]. Since all items of life skills such as self-awareness, empathy skills, effective communication skills, coping skills, anger and aggression management skills,

problem-solving, and so on are interpersonal skills that one experiences through interactions, it seems that group therapy is the most effective method to improve them [28]. In other words, the members of a group through their interactions, self-revelation, support, empathy, and common pain can achieve wider insight and understanding and better adaptation to their problems. Group provides an opportunity for its members to meet in an atmosphere free of judgment, explore their inner space, and focus on experience. In addition, a person who has difficulty in solving the problem probably has no social and environmental protection; such a person becomes disabled in problem solving because of deprivation from environmental protection [29]. Therefore, in such cases, LST can enable the application of effective tactics to deal with everyday problems. In other words, life skills are important coping strategies that can increase people's personal and social development and reduce their physical and mental problems [30]. Taking part in life skills courses was also effective in increasing maternal environmental and social relations. Sorensen et al. [31], Chen [32], and Robinson [33] confirmed the findings of this research. They reported that mothers who received LST, learned communication skills very well along with a wide range of skills and social strategies, including dialogue skills, courage and ability to listen, as well as willingness to listen to others, and respect for the others' feelings and opinions which are very important in social interactions and interpersonal relationships. Furthermore, these trainings often make changes in the lifestyles of individuals. However, one must be careful when generalizing the results of this study because our study had certain limitations, including low statistical population and limited number of sessions. It is suggested that other family members, especially fathers be included in future studies.

Conclusion

People who receive LST may have strong emotional support network. They will also deal more efficiently with problems due to their

increased tolerance and potential in problem solving.

Acknowledgments

The authors wish to appreciate the kind mothers who generously and compassionately cooperated in the conduct of this study.

REFERENCES

1. Movallali G, Amiri M, Yousefi Afrashteh M, Morovati Z. Parental stress and mental health in mothers of children with hearing impairment: the effectiveness of a behavioural training program. *IOSR Journal of Humanities and Social Science*. 2015;20(7):89-95.
2. Aslani L, Azkhosh M, Movallali G, Younesi SJ, Salehy Z. The effectiveness of resiliency training program on the components of quality of life in mothers with hearing-impaired children. *IOSR Journal of Research and Method in Education*. 2014;4(3):62-6.
3. Abbasi S, Sajedi F, Hemmati S, Rezasoltani P. The effectiveness of life skills training on quality of life in mothers of children with down syndrome. *Iranian Rehabilitation Journal*. 2014;12(22):29-34.
4. Movallali G, Dousti M, Abedi-Shapourabadi S. The effectiveness of positive parenting program (triple P) on mental health of parents of hearing impaired children. *J Appl Environ Biol Sci*. 2015;5(5S):286-90.
5. Dempsey I, Keen D, Pennell D, O'Reilly J, Neilands J. Parent stress, parenting competence and family-centered support to young children with an intellectual or developmental disability. *Res Dev Disabil*. 2009;30(3):558-66.
6. Botvin GJ, Kantor LW. Preventing alcohol and tobacco use through life skills training theory, methods, and empirical findings. *Alcohol Res Health*. 2000;24(4):250-7.
7. McClay CA, Collins K, Matthews L, Haig C, McConnachie A, Morrison J, et al. A community-based pilot randomised controlled study of life skills classes for individuals with low mood and depression. *BMC Psychiatry*. 2015;15:17.
8. Malouff JM, Thorsteinsson EB, Schutte NS. The efficacy of problem solving therapy in reducing mental and physical health problems: a meta-analysis. *Clin Psychol Rev*. 2007;27(1):46-57.
9. Kazemi R, Momenia S, Abolghasemi A. The effectiveness of life skill training on self-esteem and communication skills of students with dyscalculia. *Procedia Soc Behav Sci*. 2014;114:863-6.
10. Yankey T, Biswas UN. Life skills training as an effective intervention strategy to reduce stress among tibetan refugee adolescents. *J Refug Stud*. 2012;25(4):514-36.
11. Pakdaman Savoji A, Ganji K. Increasing mental health of university students through life skills training (LST). *Procedia Soc Behav Sci*. 2013;84:1255-9.
12. Hosseinkhanzadeh AA, Yeganeh T. The effects of life skills training on marital satisfaction. *Procedia Soc Behav Sci*. 2013;84:769-72.
13. Nejat S, Montazeri A, Holakouie Naieni K, Mohammad K, Majdzadeh SR. The world health organization quality of life (WHOQOL-BREF) questionnaire: translation and validation study of the Iranian version. *Journal of School*

- of Public Health and Institute of Public Health Research. 2006;4(4):1-12. Persian.
14. Wenzel V, Weichold K, Silbereisen RK. The life skills program IPSY: positive influences on school bonding and prevention of substance misuse. *J Adolesc.* 2009;32(6):1391-401.
 15. World Health Organization. Plan for life skills education (Noori R, Mohamadkhani P, Trans.). Tehran: Branch of the mental psychological health and prevention from the abuse of the substances, World Health Organization; 2000. Persian.
 16. Esmailinasab M, Malek Mohamadi D, Ghiasvand Z, Bahrami S. Effectiveness of life skills training on increasing self-esteem of high school students. *Procedia Soc Behav Sci.* 2011;30:1043-7.
 17. Mohammadi A, Adalatzadeh Aghdam G, Kiyani R, Sattarzadeh L. Investigating effect of life skills training on family functioning of epileptic people's in Tabriz. *Procedia Soc Behav Sci.* 2011;30:2316-8.
 18. Prasertcharoensuk T, Somprach K, Ngang TK. Influence of teacher competency factors and students' life skills on learning achievement. *Procedia Soc Behav Sci.* 2015; 186:566-72.
 19. Pohan MN, Hinduan ZR, Riyanti E, Mukaromah E, Mutiara T, Tasya IA, et al. Hiv-Aids prevention through a life-skills school based program in Bandung, west java, Indonesia: evidence of empowerment and partnership in education. *Procedia Soc Behav Sci.* 2011;15:526-30.
 20. Botvin GJ, Griffin KW. School-based programmes to prevent alcohol, tobacco and other drug use. *Int Rev Psychiatry.* 2007;19(6):607-15.
 21. Taylor JL, Smith PJ, Babyak MA, Barbour KA, Hoffman BM, Sebring DL, et al. Coping and quality of life in patients awaiting lung transplantation. *J Psychosom Res.* 2008;65(1):71-9.
 22. Gamble B. Teaching life skills for student success. *Techniques: Connecting Education and careers.* 2006; 81(6):40-1.
 23. Jenaabadi H, Azizi Nejad B, Mostafapour R, Haghi R. The effect of life skills training on the mental health and level of resilience among teachers of normal students and teachers of exceptional students in Zahedan. *Open Journal of Medical Psychology.* 2015;4(2):17-22.
 24. Barlow JH, Powell LA, Gilchrist M, Fotiadou M. The effectiveness of the Training and Support Program for parents of children with disabilities: a randomized controlled trial. *J Psychosom Res.* 2008;64(1):55-62.
 25. Seid M, Varni JW, Gidwani P, Gelhard LR, Slymen DJ. Problem-solving skills training for vulnerable families of children with persistent asthma: report of a randomized trial on health-related quality of life outcomes. *J Pediatr Psychol.* 2010;35(10):1133-43.
 26. Sobhi-Gharamaleki N, Rajabi S. Efficacy of life skills training on increase of mental health and self esteem of the students. *Procedia Soc Behav Sci.* 2010;5:1818-22.
 27. Turner NE, Macdonald J, Somerset M. Life skills, mathematical reasoning and critical thinking: a curriculum for the prevention of problem gambling. *J Gamb Stud.* 2008;24(3):367-80.
 28. Tayyebi Ramin Z, Dadkhah A, Bahmani B, Movallali G. Effectiveness of group logo therapy on increasing the quality of the mothers' life of hearing impaired children. *Applied Psychology.* 2014;2(6):18-26.
 29. Timmer SG, Ho LK, Urquiza AJ, Zebell NM, Fernandez Y Garcia E, Boys D. The effectiveness of parent-child interaction therapy with depressive mothers: the changing relationship as the agent of individual change. *Child Psychiatry Hum Dev.* 2011;42(4):406-23.
 30. Vatankhahah H, Daryabari D, Ghadami V, Khanjan Shoeibi E. Teaching how life skills (anger control) affect the happiness and self-esteem of Tonekabon female students. *Procedia Soc Behav Sci.* 2014;116:123-6.
 31. Sorensen G, Gupta PC, Nagler E, Viswanath K. Promoting life skills and preventing tobacco use among low-income Mumbai youth: effects of Salaam Bombay Foundation intervention. *PLoS One.* 2012;7(4):e34982.
 32. Chen G. Social support, spiritual program, and addiction recovery. *Int J Offender Ther Comp Criminol.* 2006;50(3):306-23.
 33. Naseri A, Babakhani N. The effect of life skills training on physical and verbal aggression male delinquent adolescents marginalized in Karaj. *Procedia Soc Behav Sci.* 2014;116:4875-9.