

RESEARCH ARTICLE

A pilot study on the short-term effects of myofascial release and core stability exercises on balance in chronic low back pain

Mina Mavajian¹, Zahra Fakhari^{1*}, Soofia Naghdi^{1,2,3}, Hossein Bagheri¹, Shohreh Jalaie⁴

¹- Department of Physiotherapy, School of Rehabilitation, Tehran University of Medical Sciences, Tehran, Iran

²- Sports Medicine Research Center, Neuroscience Institute, Tehran University of Medical Sciences, Tehran, Iran

³- Neuromusculoskeletal Research Center, Iran University of Medical Sciences, Tehran, Iran

⁴- School of Rehabilitation, Tehran University of Medical Sciences, Tehran, Iran

Received: 26 Jan 2020, Revised: 13 Feb 2020, Accepted: 19 Feb 2020, Published: 15 Apr 2020

Abstract

Background and Aim: Balance disorders are among common complications of chronic low back pain (LBP). Since the tissue and echogenicity of thoracolumbar fascia changes in chronic LBP, the independent motion of fascia layers would negatively alter, and it may affect the control of spine movements. It has been shown that myofascial release (MFR) improves the function of transversus abdominis as a core muscle in LBP. So, MFR may affect the balance and postural control in these patients. The aim of this study was to evaluate the immediate effects of MFR and core stability (CS) exercises on dynamic balance and pain in women with chronic non-specific LBP.

Methods: In this pilot study, with a pretest-posttest design, 10 females aged 30 ± 9.04 participated. The participants received one session of MFR on thoracolumbar fascia and CS exercises. The outcome measures were dynamic balance by Y balance test (YBT) and pain with visual analogue scale (VAS), which were assessed before and immediately after the intervention.

Results: Dynamic balance improved significantly ($p < 0.02$) after intervention. The VAS as an indicator of pain intensity showed significant improvement ($p = 0.005$).

Conclusion: The application of MFR plus CS exercises for a single session is effective in improving dynamic balance and pain in women with chronic non-specific LBP.

Keywords: Dynamic balance; core stability; low back pain; myofascial release

Citation: Mavajian M, Fakhari Z, Naghdi S, Bagheri H, Jalaie S. A pilot study on the short-term effects of myofascial release and core stability exercises on balance in chronic low back pain. *Aud Vestib Res.* 2020;29(2):109-16.

Introduction

Postural stability is an essential component for different daily activities. Evidence suggest that subjects with low back pain (LBP) have altered automatic postural coordination and thus demonstrated some sort of balance disorders [1]. Lifetime prevalence of non-specific LBP is about 80% so it is a high-priced issue to healthcare services worldwide [2]. Most of people experiencing LBP, will develop chronic symptoms lasting longer than one year [3]. In female

* **Corresponding author:** Department of Physiotherapy, School of Rehabilitation, Tehran University of Medical Sciences, Piche-Shemiran, Enghelab Ave., Tehran, 1148965141, Iran. Tel: 009821-77535132, E-mail: fakhari@tums.ac.ir

subjects, the overall mean prevalence of LBP is significantly higher [4].

In chronic LBP, alternations of passive structures and muscle recruitment patterns are common [5-8]. These factors could easily affect postural stability and balance in subjects with LBP [9]. Postural instability and balance dysfunction would also remarkably affect daily activities of these patients [10,11]. Numerous neuromuscular processes such as sensory input, central processing, and neuromuscular responses are prerequisite for good postural control and balance. The sensory components include the vestibular, visual and proprioceptive systems. An appropriate motor response requires an intact neuromuscular system and sufficient muscle strength [12,13]. Proprioceptive impairment and decreased core muscles' strength has also been known as one of the plausible causes for balance impairments in LBP [13].

Core muscles also make a major contribution to controlling postural alignment, stability and maintaining dynamic balance in different daily activities [14]. Among the therapeutic and strengthening exercises, core stabilization with special focus on transversus abdominis (TrA) got particular attention. An anticipatory activation of TrA helps better stabilizing the spine and therefore providing better dynamic control in various functions [15]. In patients with LBP decreased amplitude of TrA muscle activity or delayed onset of its activation were pointed out that could represent changes in motor control and planning [16].

Manual therapy includes a wide variety of techniques. Myofascial release (MFR) is a form of manual therapy which consists of a low load, long duration stretch to the myofascial complex. It helps enhancing function, reducing pain and retrieving desired structure of the fascia [17]. The thoracolumbar fascia (TLF), that placed posteriorly, from iliac crest and sacrum up to rib cage, plays an essential role in low back pain. It is an important component for transmitting forces and making stability through multiple connections with muscles and vertebral column [18].

In chronic LBP, TLF underwent some changes in

its tissue and echogenicity [19], therefore the independent motion of fascia layers would negatively alter and movement control might be changed [8]. There is some evidence suggesting that fascia may act like a smooth muscle and in this way affects musculoskeletal dynamics [20,21]. Chen et al. indicated that MFR on the TLF is efficient in terms of increasing sliding of TrA during contraction [22]. Other researches also investigated the effects of MFR or core stability (CS) exercises on functional mobility, pain, balance separately [17,18,23,24], but to the best of our knowledge this study is the first to investigate the possible effects of MFR beside CS exercises on balance in women with chronic LBP.

Considering the strong relationship between TLF, spinal column and core muscles, and also regarding different changes of TLF and TrA in chronic LBP, we postulated that the combination of MFR on TLF and core stability exercises might have further beneficial effects on pain and dynamic balance in female patients with chronic non-specific LBP.

Methods

In this pilot study with a pretest-posttest design, 10 women with chronic non-specific LBP participated. Inclusion criteria were women aged 18 to 50 years old with a history of LBP for at least three months, absence of conditions like congenital low back disorders, back, abdomen or lower extremities surgery in past three months, lumbar vertebrae fracture, malignancies, chronic vertigo, vestibular disorders, pregnancy and being active athletes. Either the participants who simultaneously participate in other treatment protocols or were unwilling to continue the protocol were excluded from this study. All subjects read and signed an informed consent form before the beginning of study. The Review Board and the Ethics Committee of Tehran University of Medical Sciences approved the study protocol (Approval ID: IR.TUMS.FNM.REC.1398.066).

All participants were interviewed to collect demographic data including age, weight, height and period of low back pain. Outcome measures including dynamic balance and pain were

assessed by Y balance test (YBT) and visual analogue scale (VAS), respectively before and immediately after the intervention.

YBT is a reliable and valid tool for measuring dynamic balance and neuromuscular aspects of core stability [25]. The assessor instructed YBT protocols in detail and all patients were allowed to practice a trial on each leg, for all directions prior to performing main trial. The test was performed with bare feet, hands placed on hips and foot in contact with the ground. To perform the test, while the patient stood on the center point, and maintaining single leg stance, tried to reach the free limb to the most possible distal part of the scaled tape in anterior, postromedial and postrolateral directions respectively. Between three trials of each leg for each direction, the best records, normalized to leg length were selected to be analyzed. The lower limb length was measured with the patient in supine lying position and straight legs, from Anterior Superior Iliac Spine (ASIS) to the most prominent part of medial malleolus [26]. Each single trial was rejected and repeated if the patient failed to 1) maintain unilateral stance 2) return the reach foot to the starting position 3) maintain full contact of standing foot to the ground (heel off the ground) [27]. VAS was used for measuring pain intensity. The patients were asked to highlight their pain intensity on the scaled continuum (from none to an extreme amount of pain). For data recording a precise ruler was used. The ruler (from the zero point) placed exactly between the start point of the continuum and the patient's mark to the nearest 1 mm so, the possible range was from 1 to 10 cm [28]. Reliability and validity of VAS had been adequately tested for patients with LBP by various studies [29,30].

MFR techniques in this study included compression release and cross-hand release in lower back and TLF which were performed in both prone and side lying positions (illustrated as Fig. 1 a-d) the therapist's hands in compression release was side by side and for cross-hand release, was crossed in opposite directions. Each of the mentioned techniques held for 120s and repeated three times in tissue barriers. To perform the techniques physical therapist applied adequate

pressure in direction of facial restriction just until the first tissue barrier was felt. Then, the sufficient pressure was engaged for about 120s without gliding over the skin or pushing the tissue tightly [31], until the physical therapist felt a softening feeling under the hands and thoracolumbar fascia starts to yield. As it achieved, the technique would be progressed to the deeper tissue barrier [32]. After MFR, patients performed core stability with abdominal drawing-in maneuver (ADIM) in six positions (illustrated as Fig. 2 a-f) under the therapist supervision. At first, therapist instructed the correct form of ADIM with pelvic floor muscles' activation and all the detailed important points of exercises. Patients were also instructed that breath holding, ribcage depression or abdominal hollowing should not be done during the exercise program. On average exercise program took 15 minutes for each patient.

SPSS statistical software (version 23) was used for data analyzing. One-sample Kolmogorov-Smirnov test was used to determine the distribution of data and showed that all the data were normally distributed. Paired t-test was conducted to compare dependent variables including YBT and VAS before and after the intervention. Cohen's d was also calculated to determine treatment effect size [33].

Results

This study conducted on 10 women with chronic non-specific low back pain (mean and standard deviation of age 30 ± 9.04 year) participated in the present study. All of them had a history of chronic low back pain (mean and standard deviation of low back pain duration 5.02 ± 4.09 year) (Table 1).

For an accurate comparison of balance between subjects all data extracted from YBT were normalized to the patients' leg length and the normalized data were used in analyses. Participants improved significantly on pain intensity measured by visual analogue scale (VAS) ($p = 0.005$) and also dynamic balance measured by YBT of both feet in all directions (anterior, postromedial and postrolateral) ($p \leq 0.02$; Table 2). For all variables including dynamic balance




Fig. 1. Myofascial release techniques, a) compression release in prone position, b) cross-hand release in prone position, c) compression release in side-lying position, and d) cross-hand release in side-lying position.

and pain, effect size measures were medium to large ($p < 0.05$; Table 2). Test-retest (intra-rater) reliability of YBT scores for all directions of both feet were excellent ($r > 0.97$).

Discussion

The primary findings of the present study were in favor of our hypothesis, so the intervention was effective in terms of improving dynamic balance and reducing pain in women with chronic non-specific LBP. These findings are confirmed by

other research articles that showed the useful effects of MFR treatment in different population [17,34], but concurrent investigation of MFR beside CS exercises on dynamic balance of these patients seems limited. Tozzi et al. showed positive short-term improvements in fascial layers' mobility and pain perception of patients with neck or low back pain following MFR [35]. Even on healthy subjects after MFR, TLF stiffness reduction was reported [18]. Findings of another study by Chen et al. showed that sliding




Fig. 2. Core stability exercises, a) abdominal drawing-in maneuver, crook lying 10s hold 10 repetitions, b) abdominal drawing-in maneuver, heel slide, each leg 10 repetitions, c) bridging in supine, 10s hold, 10 repetitions, d) abdominal drawing-in maneuver with moving the swiss ball, 10 repetitions, e) abdominal drawing-in maneuver in quadruped, 10s hold, 10 repetitions, and f) bird dog exercise 10 repetitions on each side.

of musculofascial junctions of transversus abdominis muscle was significantly and clinically improved following MFR in patients with chronic LBP [22]. Some of possible long term complications of chronic LBP are: impaired neuromuscular control [13], altered recruitment patterns of different muscles [36], fewer mechanoreceptors in lumbar fascia [37], impaired lumbopelvic proprioception [38] and increased connective tissue adhesions. All of these factors could lead to mobility restriction, postural control deficits and pain. Some examinations have

demonstrated the presence of unmyelinated terminal nerve endings in the lumbodorsal fascia [5]. In some situation like stiffness, they could be entrapped by the myofibrosis parts in tissue [39] so, one possible mechanism of pain reduction in our study could be pressure release of these sensitive free nerve endings [40], after fascial release and improving the layers' mobility. Concerning the researches on CS exercises, Kim and Park showed that these types of exercises could lead to immediate significant improvements in balance [23] also Trampas et al. [24], evaluated

Table 1. Demographic characteristics of patients with chronic low back pain (n = 10)

| Characteristics | Mean ± SD |
|--------------------------|--------------|
| Age (y) | 30 ± 9.04 |
| Weight (kg) | 63.5 ± 12 |
| Height (m) | 1.64 ± 0.05 |
| BMI (kg/m ²) | 23.34 ± 3.66 |
| Duration of LBP | 5.2 ± 4.09 |

BMI; Body Mass Index, LBP; low back pain

the immediate effects of clinical massage and CS exercises on dynamic balance performance in chronic LBP. Results demonstrated significant improvements in balance performance and pressure pain threshold. These type of exercises could play an important role in decreasing pain and improving function of patients with chronic LBP [41,42]. However, in long term evaluation CS exercises are not superior to other forms of active exercises [42,43]. Core stability exercises could have the potential to stimulate adaptive processes in the neuromuscular system like: more efficient neural recruitment patterns,

faster response of nervous system, increased synchronization of motor units and decreasing of neural inhibitory reflexes [44]. These processes might improve balance performance and functional mobility in older adults [45]. Although for a better understanding of this topic, more research is required.

CS exercises of the present study were executed by special focus on TrA muscle activation. An anticipatory action of this muscle helps stabilizing the spine and provide a better basis for daily movements [15]. Schleip et al. [21], showed that fascia may also express some kind of contractile behavior, therefore it seems that TrA activation alongside with TLF's accurate function would help evoking better stabilization function of core muscles and dynamic balance performances.

Regarding the close connection of TLF with abdominal and back muscles and also considering its alternations in chronic LBP and the beneficial effects of MFR in releasing fascial layers, it seems that implementation of MFR plus CS exercises might be effective in better force transforming mechanics by facilitating the mobility and gliding of layers and hence, increasing stability. Further studies are needed to investigate the long-term effects of this treatment or to compare different population.

Table 2. Mean and standard deviation of outcome measures pre and post intervention in patients with chronic low back pain (n = 10)

| Variable | Pre | Post | p | Effect size* |
|-------------|-----------|-----------|---------|--------------|
| YBT. L. ant | 0.89±0.08 | 0.96±0.12 | 0.022 | 0.69 |
| YBT. L. pm | 0.97±0.15 | 1.11±0.12 | < 0.001 | 0.99 |
| YBT. L. pl | 1.04±0.1 | 1.15±0.1 | < 0.001 | 1.13 |
| YBT. R. ant | 0.86±0.1 | 0.94±0.14 | 0.029 | 0.57 |
| YBT. R. pm | 0.99±0.12 | 1.11±0.12 | < 0.001 | 0.93 |
| YBT. R. pl | 0.99±0.12 | 1.1±0.14 | < 0.001 | 0.79 |
| VAS | 4.12±1.73 | 2.4±2.36 | 0.005 | 0.83 |

YBT; Y balance test, L; left, R; right, ant; anterior, pm; postromedial, pl; postrolateral, VAS; visual analogue scale
*Cohen's d

Conclusion

In conclusion, the application of myofascial release (MFR) plus core stability (CS) exercises could immediately improve dynamic balance in anterior, postromedial and postrolateral directions also significantly improvement in pain intensity was achieved in women with chronic non-specific low back pain (LBP).

Acknowledgments

The information contained in this article was extracted from MSc. thesis of M. Mavajian, which submitted to Tehran University of Medical Sciences.

Conflict of interest

No potential conflict of interest relevant to this article was reported.

References

1. Henry SM, Hitt JR, Jones SL, Bunn JY. Decreased limits of stability in response to postural perturbations in subjects with low back pain. *Clin Biomech (Bristol, Avon)*. 2006;21(9):881-92. doi: [10.1016/j.clinbiomech.2006.04.016](https://doi.org/10.1016/j.clinbiomech.2006.04.016)
2. Nachemson AL, Jonsson E. Neck and back pain: the scientific evidence of causes, diagnosis, and treatment. Philadelphia: Lippincott Williams & Wilkins; 2000.
3. Hestbaek L, Leboeuf-Yde C, Manniche C. Low back pain: what is the long-term course? A review of studies of general patient populations. *Eur Spine J*. 2003;12(2):149-65. doi: [10.1007/s00586-002-0508-5](https://doi.org/10.1007/s00586-002-0508-5)
4. Hoy D, Bain C, Williams G, March L, Brooks P, Blyth F, et al. A systematic review of the global prevalence of low back pain. *Arthritis Rheum*. 2012;64(6):2028-37. doi: [10.1002/art.34347](https://doi.org/10.1002/art.34347)
5. Wilke J, Schleip R, Klingler W, Stecco C. The lumbo-dorsal fascia as a potential source of low back pain: a narrative review. *Biomed Res Int*. 2017;2017:5349620. doi: [10.1155/2017/5349620](https://doi.org/10.1155/2017/5349620)
6. Selkow NM, Eck MR, Rivas S. Transversus abdominis activation and timing improves following core stability training: a randomized trial. *Int J Sports Phys Ther*. 2017;12(7):1048-56. doi: [10.26603/ijsp20171048](https://doi.org/10.26603/ijsp20171048)
7. Moghadam N, Ghaffari MS, Noormohammadpour P, Rostami M, Zarei M, Moosavi M, et al. Comparison of the recruitment of transverse abdominis through drawing-in and bracing in different core stability training positions. *J Exerc Rehabil*. 2019;15(6):819-25. doi: [10.12965/jer.1939064.352](https://doi.org/10.12965/jer.1939064.352)
8. Langevin HM, Fox JR, Koptiuch C, Badger GJ, Greenan-Naumann AC, Bouffard NA, et al. Reduced thoracolumbar fascia shear strain in human chronic low back pain. *BMC Musculoskelet Disord*. 2011;12:203. doi: [10.1186/1471-2474-12-203](https://doi.org/10.1186/1471-2474-12-203)
9. della Volpe R, Popa T, Ginanneschi F, Spidalieri R, Mazzocchio R, Rossi A. Changes in coordination of postural control during dynamic stance in chronic low back pain patients. *Gait Posture*. 2006;24(3):349-55. doi: [10.1016/j.gaitpost.2005.10.009](https://doi.org/10.1016/j.gaitpost.2005.10.009)
10. Koch C, Hänsel F. Non-specific low back pain and postural control during quiet standing-A systematic review. *Front Psychol*. 2019;10:586. doi: [10.3389/fpsyg.2019.00586](https://doi.org/10.3389/fpsyg.2019.00586)
11. Ghamkhar L, Kahlaee AH. Trunk muscles activation patterns during walking in subjects with and without chronic low back pain: a systematic review. *PM R*. 2015;7(5):519-26. doi: [10.1016/j.pmrj.2015.01.013](https://doi.org/10.1016/j.pmrj.2015.01.013)
12. Alexandrov AV, Frolov AA, Horak FB, Carlson-Kuhta P, Park S. Feedback equilibrium control during human standing. *Biol Cybern*. 2005;93(5):309-22. doi: [10.1007/s00422-005-0004-1](https://doi.org/10.1007/s00422-005-0004-1)
13. Karimi N, Ebrahimi I, Kahrizi S, Torkaman G. Evaluation of postural balance using the biodex balance system in subjects with and without low back pain. *Pak J Med Sci*. 2008;24(3):372-7.
14. Clark MA, Fater D, Reuteman P. Core (trunk) stabilization and its importance for closed kinetic chain rehabilitation. *Orthop Phys Ther Clin N Am*. 2000;9:119-35.
15. Hodges PW, Richardson CA. Contraction of the abdominal muscles associated with movement of the lower limb. *Phys Ther*. 1997;77(2):132-42. doi: [10.1093/ptj/77.2.132](https://doi.org/10.1093/ptj/77.2.132)
16. Ferreira PH, Ferreira ML, Hodges PW. Changes in recruitment of the abdominal muscles in people with low back pain: ultrasound measurement of muscle activity. *Spine (Phila Pa 1976)*. 2004;29(22):2560-6. doi: [10.1097/01.brs.0000144410.89182.f9](https://doi.org/10.1097/01.brs.0000144410.89182.f9)
17. Ajimsha MS, Al-Mudahka NR, Al-Madzhar JA. Effectiveness of myofascial release systematic review of randomized controlled trials. *J Bodyw Mov Ther*. 2015;19(1):102-12. doi: [10.1016/j.jbmt.2014.06.001](https://doi.org/10.1016/j.jbmt.2014.06.001)
18. Wong KK, Chai HM, Chen YJ, Wang CL, Shau YW, Wang SF. Mechanical deformation of posterior thoracolumbar fascia after myofascial release in healthy men: A study of dynamic ultrasound. *Musculoskelet Sci Pract*. 2017;27:124-30. doi: [10.1016/j.math.2016.10.011](https://doi.org/10.1016/j.math.2016.10.011)
19. Langevin HM, Stevens-Tuttle D, Fox JR, Badger GJ, Bouffard NA, Krag MH. Ultrasound evidence of altered lumbar connective tissue structure in human subjects with chronic low back pain. *BMC Musculoskelet Disord*. 2009;10:151. doi: [10.1186/1471-2474-10-151](https://doi.org/10.1186/1471-2474-10-151)
20. Spector M. Musculoskeletal connective tissue cells with muscle: expression of muscle actin in and contraction offibroblasts, chondrocytes, and osteoblasts. *Wound Repair Regen*. 2001;9(1):11-8. doi: [10.1046/j.1524-475x.2001.00011.x](https://doi.org/10.1046/j.1524-475x.2001.00011.x)
21. Schleip R, Klingler W, Lehmann-Horn F. Active fascial contractility: Fascia may be able to contract in a smooth muscle-like manner and thereby influence musculoskeletal dynamics. *Med Hypotheses*. 2005;65(2):273-7. doi: [10.1016/j.mehy.2005.03.005](https://doi.org/10.1016/j.mehy.2005.03.005)
22. Chen YH, Chai HM, Shau YW, Wang CL, Wang SF. Increased sliding of transverse abdominis during contraction after myofascial release in patients with chronic low back pain. *Man Ther*. 2016;23:69-75. doi: [10.1016/j.math.2015.10.004](https://doi.org/10.1016/j.math.2015.10.004)
23. Kim JJ, Park SY. Immediate effects of the trunk stabilizing exercise on static balance parameters in double-leg and one leg stance. *J Phys Ther Sci*. 2016;28(6):1673-5. doi: [10.1589/jpts.28.1673](https://doi.org/10.1589/jpts.28.1673)
24. Trampas A, Mpeneka A, Malliou V, Godolias G, Vlachakis P. Immediate effects of core-stability exercises and clinical massage on dynamic-balance performance of patients with chronic specific low back pain. *J Sport Rehabil*. 2015;24(4):373-83. doi: [10.1123/jsr.2014-0215](https://doi.org/10.1123/jsr.2014-0215)
25. Okubo Y, Kaneoka K, Imai A, Shiina I, Tatsumura M, Izumi S, et al. Electromyographic analysis of transversus abdominis and lumbar multifidus using wire electrodes during lumbar stabilization exercises. *J Orthop Sports Phys Ther*. 2010;40(11):743-50. doi: [10.2519/jospt.2010.3192](https://doi.org/10.2519/jospt.2010.3192)
26. Plisky PJ, Gorman PP, Butler RJ, Kiesel KB, Underwood FB, Elkins B. The reliability of an instrumented device for measuring components of the star excursion balance test. *N Am J Sports Phys Ther*. 2009;4(2):92-9.
27. Butowicz CM, Ebaugh DD, Noehren B, Silfies SP. Validation of two clinical measures of core stability. *Int J Sports Phys Ther*. 2016;11(1):15-23.
28. Hawker GA, Mian S, Kendzerska T, French M. Measures of adult pain: Visual Analog Scale for Pain (VAS Pain), Numeric Rating Scale for Pain (NRS Pain), McGill Pain Questionnaire (MPQ), Short-Form McGill Pain Questionnaire (SF-MPQ), Chronic Pain Grade Scale (CPGS), Short Form-36 Bodily Pain Scale (SF-36 BPS), and Measure of Intermittent and Constant Osteoarthritis Pain (ICOAP). *Arthritis Care Res (Hoboken)*. 2011;63(Suppl

- 11):S240-52. doi: [10.1002/acr.20543](https://doi.org/10.1002/acr.20543)
29. Smeets RJ, Vlaeyen JW, Kester AD, Knottnerus JA. Reduction of pain catastrophizing mediates the outcome of both physical and cognitive-behavioral treatment in chronic low back pain. *J Pain*. 2006;7(4):261-71. doi: [10.1016/j.jpain.2005.10.011](https://doi.org/10.1016/j.jpain.2005.10.011)
30. Carlsson AM. Assessment of chronic pain. I. Aspects of the reliability and validity the visual analogue scale. *Pain*. 1983;16(1):87-101. doi: [10.1016/0304-3959\(83\)90088-x](https://doi.org/10.1016/0304-3959(83)90088-x)
31. Manheim CJ, The myofascial release manual. 3rd ed. New Jersey: Slack Incorporated; 2001.
32. Duncan R. Myofascial release (hands-on guides for therapists). 1st ed. Champaign, IL: Human Kinetics, Inc.; 2014.
33. Middel B, van Sonderen E. Statistical significant change versus relevant or important change in (quasi) experimental design: some conceptual and methodological problems in estimating magnitude of intervention-related change in health services research. *Int J Integr Care*. 2002;2:e15. doi: [10.5334/ijic.65](https://doi.org/10.5334/ijic.65)
34. Beardsley C, Škarabot J. Effects of self-myofascial release: A systematic review. *J Bodyw Mov Ther*. 2015; 19(4):747-58. doi: [10.1016/j.jbmt.2015.08.007](https://doi.org/10.1016/j.jbmt.2015.08.007)
35. Tozzi P, Bongiorno D, Vitturini C. Fascial release effects on patients with non-specific cervical or lumbar pain. *J Bodyw Mov Ther*. 2011;15(4):405-16. doi: [10.1016/j.jbmt.2010.11.003](https://doi.org/10.1016/j.jbmt.2010.11.003)
36. Jacobs JV, Henry SM, Nagle KJ. People with chronic low back pain exhibit decreased variability in the timing of their anticipatory postural adjustments. *Behav Neurosci*. 2009;123(2):455-8. doi: [10.1037/a0014479](https://doi.org/10.1037/a0014479)
37. Bednar DA, Orr FW, Simon GT. Observations on the pathomorphology of the thoracolumbar fascia in chronic mechanical back pain. A microscopic study. *Spine (Phila Pa 1976)*. 1995;15;20(10):1161-4. doi: [10.1097/00007632-199505150-00010](https://doi.org/10.1097/00007632-199505150-00010)
38. Radebold A, Cholewicki J, Polzhofer GK, Greene HS. Impaired postural control of the lumbar spine is associated with delayed muscle response times in patients with chronic idiopathic low back pain. *Spine (Phila Pa 1976)*. 2001;26(7):724-30. doi: [10.1097/00007632-200104010-00004](https://doi.org/10.1097/00007632-200104010-00004)
39. Staud R. Peripheral pain mechanisms in chronic widespread pain. *Best Pract Res Clin Rheumatol*. 2011; 25(2):155-64. doi: [10.1016/j.berh.2010.01.010](https://doi.org/10.1016/j.berh.2010.01.010)
40. Fischer AA. Pressure algometry over normal muscles. Standard values, validity and reproducibility of pressure threshold. *Pain*. 1987;30(1):115-26. doi: [10.1016/0304-3959\(87\)90089-3](https://doi.org/10.1016/0304-3959(87)90089-3)
41. Smith BE, Littlewood C, May S. An update of stabilisation exercises for low back pain: a systematic review with meta-analysis. *BMC Musculoskelet Disord*. 2014; 9;15:416. doi: [10.1186/1471-2474-15-416](https://doi.org/10.1186/1471-2474-15-416)
42. Standaert CJ, Weinstein SM, Rumpeltes J. Evidence-informed management of chronic low back pain with lumbar stabilization exercises. *Spine J*. 2008;8(1):114-20. doi: [10.1016/j.spinee.2007.10.015](https://doi.org/10.1016/j.spinee.2007.10.015)
43. May S, Johnson R. Stabilisation exercises for low back pain: a systematic review. *Physiotherapy*. 2008;94(3): 179-89. doi: [10.1016/j.physio.2007.08.010](https://doi.org/10.1016/j.physio.2007.08.010)
44. Staron RS, Karapondo DL, Kraemer WJ, Fry AC, Gordon SE, Falkel JE, et al. Skeletal muscle adaptations during early phase of heavy-resistance training in men and women. *J Appl Physiol (1985)*. 1994;76(3):1247-55. doi: [10.1152/jappl.1994.76.3.1247](https://doi.org/10.1152/jappl.1994.76.3.1247)
45. Granacher U, Lacroix A, Muehlbauer T, Roettger K, Gollhofer A. Effects of core instability strength training on trunk muscle strength, spinal mobility, dynamic balance and functional mobility in older adults. *Gerontology*. 2013;59(2):105-13. doi: [10.1159/000343152](https://doi.org/10.1159/000343152)