



Biosurfactant Producing Bacteria on Oily Areas of Ruminant Skin

Azizollah Ebrahimi^{a,*}, Najmeh Tashi^a, Saeid Karimi^b

^aDepartment of Pathobiology, School of Veterinary Science, Shahrekord University,
P. O. Box: 115, Postal Code, 88186/34141, Shahrekord, Iran.

^bDepartment of Animal Science, Agricultural College of TarbiatModares, Tehran, Iran

Abstract

Biosurfactants are surface-active compounds produced by microorganisms. In this study, we collected 60 inguinal area and ear canal samples from cows, sheep, and goats (each, 10 animals) and screened for biosurfactant-producing bacteria. We also determined the genera of culturing strains. Fifty six hemolytic bacterial strains (27, 22 and 7, from cows, sheep and goats, respectively) were isolated. Oil spreading test and bioemulsifying activities were measured for all isolates. The cows' samples had higher population of positive strains than other animals, so that 5 isolates from inguinal area and 4 from ear canal samples (16.1%) were positive for all tests. For sheep, the numbers were 6 and one (12.5%) while for goats one and two (5.3%), respectively. Totally, 19 isolates (33.9%) were positive for all examinations out of them 12 were gram positives. The microorganisms isolated in this study could well be sources of novel biosurfactants. Further investigation into the composition of the biosurfactants and phylogenetic determination of biosurfactant producing bacteria is suggested.

Keywords: Biosurfactant; Emulsification; Oil spreading; Ruminant skin.

Received: February 2, 2011; *Accepted:* April 5, 2011

1. Introduction

Biosurfactants are amphiphilic compounds produced on living surfaces, mostly on microbial cell surfaces, or excreted extracellularly and contain hydrophobic and hydrophilic moieties that confer the ability to accumulate between fluid phases, thus reducing surface and interfacial tension at the surface and interface, respectively [1]. They are a structurally diverse group of surface-

active molecules synthesized by microorganisms [2].

Rosenberg and Ron [3] suggested that biosurfactants can be divided into low-molecular-mass molecules, which efficiently lower surface and interfacial tension, and high molecular-mass polymers, which are more effective as emulsion-stabilizing agents.

Apart from their obvious role as agents that decrease surface and interfacial tension, thus promoting the formation and stabilization of emulsions, surfactants can have several other functions. They improve consistency and texture of fat-based products [4]. Several

*Correspondence author: Azizollah Ebrahimi, Department of Pathobiology, College of Veterinary Science, P.O. BOX 115, Postal Code, 88186/34141, Shahrekord University, Shahrekord, Iran
E-mail: A_kahrizsangi@yahoo.com
Phone: (+98)3814424427; Fax: (+98)3814424427

Table 1. Biosurfactant-producing bacteria isolated from Cows

Isolate Ear/IA*	Ear			IA		
	E _{24h} %	E _{72h} %	O.S. _{SD} (cm)**	E _{24h} %	E _{72h} %	O.S. _{SD} (cm)
<i>Bacillus</i> spp/ <i>Escherichia</i> spp	56	56	4.55±0.05	40	47.8	5.75±0.75
<i>Bacillus</i> spp/ <i>Bacillus</i> spp	40	52	3.2±0.3	44	52.3	5.00±0.5
Staphylococci/ <i>Providentia</i> spp	40	48	3.55±0.25	50	63.6	5.55±0.25
<i>Lactobacillus</i> spp/ <i>Aeromonas</i> spp	44	48	5.25±0.65	52.3	47.8	5.75±0.15
<i>Bacillus</i> spp/ <i>Staphylococcus</i> spp	45	44	6.45±1.05	42.8	45	5.65±0.35
<i>Pasteurella</i> spp/ <i>Bacillus</i> spp	47	60	5.4±0.2	52.3	47.8	4.5±0.3
<i>Bacillus</i> spp/ <i>Bacillus</i> spp	48	48	4.75±0.15	50	55	5.75±0.45
<i>Lactobacillus</i> spp/ <i>Lactobacillus</i> spp	56	56	4.55±0.05	54.5	60.8	4.75±0.15
<i>Acinetobacter</i> spp/ <i>Acinetobacter</i> spp	56.5	52.1	6.02 ±0.25	59	58.3	5.4±0.1
<i>Lactobacillus</i> spp/ <i>Bacillus</i> spp	52	52	4.15±0.25	45.4	43.4	4.55±0.25
<i>Bacillus</i> spp/ <i>Falavobacterium</i> spp	48	48	5.4±0.2	52.1	48	5.25±0.25
<i>Bacillus</i> spp/ <i>Lactobacillus</i> spp	48	52	5.05±0.05	45.4	56	4.6±0.0
Staphylococcus spp./ <i>Lactobacillus</i> spp	40	48	5.7±0.1	54.5	56	4.9±0.3
<i>Bacillus</i> spp/-	44	44	7.05±0.25	-	-	-
Control	50	50	3.55±0.05	50	50	3.55±0.05

*IA stands for inguinal area, ** O.S. SD for oil spreading and the standard deviation

biosurfactants have shown antimicrobial action against bacteria, fungi, algae and viruses [5].

There are many advantages of biosurfactants compared to their chemically synthesized counterpart. Research in this subject, will make them highly sought after biomolecules for present and future applications as fine specialty chemicals, biological control agents and new generation molecules for pharmaceutical, cosmetic and health care industries.

Although a large number of biosurfactant producers have been reported in the literature, reports regarding screening and isolation of these microorganisms from animals are scarce. The primary aim of the present study was to investigate biosurfactant producing bacteria (PBB) habitats in ear canal and inguinal areas (IA) (as oily skin areas) of ruminants.

2. Materials and methods

2.1. Sample collection

The study was carried out through June 2009 to December 2010 on 30, Holstein cows, native sheep and goats, (each 10) randomly selected from animals in farms of Shahrekord University.

All animals were adults and were found to be apparently healthy. Samples were collected,

by inserting sterile cotton-tipped applicator sticks into the ear canal and rubbing on inguinal areas. The surfaces were thoroughly rubbed by rolling the swabs to attain effective contact. The swabs were put in separate sterile test tubes containing sterile pepton water (Merck cat. QB-65-5015), labeled and kept in a cool box and transported to the veterinary microbiology laboratory of veterinary college of Shahrekord University on the day of sampling for further processing.

For bacteriological examination, the swabs were removed from the bottles and streaked over the plates of blood agar-base (Scharlau 01-352) supplemented with 7% sheep blood. The streaking was further spread with inoculating loop to aid colony isolation. The plates were labeled and incubated aerobically at 37 °C for 24-48 h [6].

One colony was selected from those colonies that have similar morphologies and sub-cultured on blood agar plates for further analysis.

2.2. Screening methods

The first screening test for identification and isolation of BPB is hemolysis test [7]. For assaying hemolytic activity, each strain was streaked onto blood agar plates and incubated for 48 h at 37 °C. The plates were visually inspected for zones of clearing around the

Table 2. Biosurfactant-producing bacteria isolated from Sheep

Isolate Ear/ IA*	Ear			IA		
	E _{24h} %	E _{72h} %	O.S. SD ** (cm)	E _{24h} %	E _{72h} %	O.S. SD (cm)
Lactobacillus spp /Lactobacillus spp	44	44	4.95±0.35	47	54.5	5.25 ±0.05
Lactobacillus spp/Streptococcus spp.	40	44	5.55±0.15	47	43.5	5.800 ±0.15
Bacillus spp / Bacillus spp	44	45.5	3.2±0.4	48	52	3.4 ±0.2
Bacillus spp /Lactobacillus spp	52	48	4.4±0.1	47.6	45.5	6.5 ±0.25
Bacillus spp / Bacillus spp	40	38.5	4.85±0.25	42.9	45.5	4.5 ±0.25
Bacillus spp /Lactobacillus spp	44	52	4.4±0.2	50	43.3	5.5 ±0.05
Aeromonas spp/ Bacillus spp	48	50	3.4±0.15	47.8	45.8	3.4 ±0.25
Staphylococcus spp. / Bacillus spp	52	48	5.3±0.15	48.8	50	5.7 ±0.05
Escherichia spp./ Bacillus spp	42.3	37.1	5.5±0.2	34.8	40	6.1 ±0.2
Staphylococcus spp. /Lactobacillus spp	40	46.2	3.5±0.05	40	40.7	3.6 ±0.4
Pseudomonas spp/Pseudomonas spp	44	51.9	5.2±0.1	50	60	6.2 ±0.15
Control	40	40	4.35±0.15	40	40	4.4 ±0.15

*IA stands for inguinal area, **O.S. SD for oil spreading and the standard deviation.

colonies, indicative of biosurfactant production. After gram staining, catalase and oxidase tests, identification of the isolated hemolytic positive strains were done using a standard biochemical scheme according to Balows et al [8].

Each hemolytic isolate was inoculated in tubes containing Lauria bertani broth (LB, Biomark-B699) media and incubated at 37 °C for 72 h with shaking (~50 rpm). For each set of cultures one tube of sterile LB was also incubated to use as control in further analysis.

For the oil spreading technique (OS), 50 ml of distilled water was added to a large petri dish (25 cm diameter) followed by addition of 20 µl of n-Decane (Merck, UN 2247) to the surface of the water. Ten microliters of cell-free broth of LB culture (Centrifuged at 10000 rpm for 10 min.) were then added to the surface of oil [9]. The diameter of the clear zone on the oil surface was measured. The diameters of triplicate samples from the same culture of each strain were determined.

The emulsifying capacity was evaluated by an emulsification index (E₂₄). The E₂₄ of culture samples was determined by adding 1.5 ml of kerosene and 1.5 ml of the cell-free broth in test tube, vortexed at high speed for 2 min and allowed to stand for 24 h and 72 h. The E₂₄ (and E₇₂) index is given as the percentage of the height of emulsified layer divided by the total height of the liquid column (cm). The percentage of

emulsification index calculated by using the following equation [10],

$$E_{24} = \frac{\text{Height of emulsion formed} \times 100}{\text{Total height of solution}}$$

For each test strain, centrifuged samples of incubated tubes of sterile LB were used as control.

3. Results

After culture and incubation of 60 samples (20 from each animal species, 10 ear and 10 IA) 56 hemolytic strains (27, 22 and 7, from cows, sheep and goats, respectively) were isolated. OS and bioemulsifying activities were measured for all isolates (Tables 1-3).

The cow's samples had higher population of E₂₄, E₇₂ and OS positives than other animals, so that 5 isolates from IA and 4 from ear canal samples (16.1%) were positive for all tests. For sheep the numbers were 6 and one (12.5%) while for goats one and two (5.3%) respectively. Totally 19 isolates (33.9%) were positive for all examinations, out of them 12 were gram positives.

More sensitive OS test was positive for 13 IA and 12 ear canal isolates of cows (44.6%), for sheep the numbers were 7, 1 (14.3%) and for goat 2, 3 (8.9%), respectively. Totally 38 isolates (67.8%) were positive for this test (Tables 1-3).

4. Discussion

Hemolytic activity appears to be a good

Table 3. Biosurfactant-producing bacteria isolated from goats

Isolate	Ear			IA			
	Ear/ IA*	E _{24h} %	E _{72h} %	O.S.** SD (cm)	E _{24h} %	E _{72h} %	O.S. SD (cm)
<i>Providentia</i> spp / <i>Bacillus</i> spp		54	54	7.1 ±2.05	50	54.2	5.5 ±0.05
<i>Bacillus</i> spp / <i>Bacillus</i> spp		45.8	45.5	5.4 ±0.1	45.8	50	6.3 ±0.75
<i>Lactobacillus</i> spp/ <i>Lactobacillus</i> spp		54.1	50	6.4 ±0.1	45.8	45.5	4.1 ±0.6
<i>Staphylococcus</i> spp. / -		45.8	47.8	4.9 ±0.4	-	-	-
Control		45	45	3.55 ±0.05	45	45	3.55 ±0.05

*IA stands for inguinal area, **O.S. SD for oil spreading and the standard deviation

screening criterion in the search for BPB (7). Such screening can be used to limit the number of samples. Further screening of BPB is generally carried out using monitoring parameters that estimate surface activity, such as ability to emulsify oils and dispersing or solubilization activity [11].

Comparatively high abundances of surfactant-producing bacteria were isolated from the cows and sheep (9 and 7 out of 56 isolates were positive for all tests respectively). In contrast, goats had lower surfactant producing bacteria (3 isolates).

These results suggest that probably the oily places of the skin of only some ruminants might be potential sources of surfactant-producing bacteria. However, some skin areas did not study here may contain even more surfactants produced by BPB as compared to studied areas.

Biosurfactant production by many of the isolated strains suggests that the resident bacteria could be a source of surfactants in the studied areas. A relatively biosurfactant producing *Bacillus* spp and *Lactobacillus* spp domination are represented in the isolated strains.

The function and composition of surfactants in the organisms of the examined areas has not been established. It might be suggested that the surfactants assist in the surface fat layer removal process by solubilizing hydrophobic fat layer or preventing destructive function of skin lytic substances. It may also dissolve organic matter of skin surface secreted by the different body systems or has some roles in the bacterial community formation of the skin surfaces.

Biosurfactants are often superior to commercial surfactants at solubilizing different chemicals and are more easily biodegraded [5]. Viewing biosurfactant producing bacteria in tables 1-3, the genera isolated from the studied areas, are well documented to be present in different oily environments as BPB [5, 11].

The microorganisms isolated in this study could well be sources of novel biosurfactants. Given demonstrated biosurfactant production by ruminant isolates, further investigation into the composition of the biosurfactants and phylogenetic determination of BPB is suggested.

Acknowledgments

Funding for this work was provided by School of Veterinary Science of Shahrekord University as costs for DVM research project.

References

- [1] Karanth NGK, Deo PG, Veenanadig NK. Microbial production of biosurfactants and their importance. *Curr Sci* 1999; 77: 116-23.
- [2] Lu JR, Zhao XB, Yaseen M. Biomimetic amphiphiles: biosurfactants. *Curr Opin Colloid Interface Sci* 2007; 12: 60-7.
- [3] Rosenberg E, Ron EZ. High- and low-molecular-mass microbial surfactants. *Appl Microbiol Biotechnol* 1999; 52: 154-62.
- [4] Kachholz T, Schlingmann M. Possible food and agricultural applications of microbial surfactants: an assessment. In: *Biosurfactants and biotechnology*. New York: (eds); Marcel Dekker, 1987; pp. 183-210.
- [5] Krishnaswamy M, Subbucchettiar G, Thiengungal KR, Panchaksharam S. Biosurfactants: properties, commercial production and application. *Curr Sci* 2008; 94: 736-47.
- [6] Carter GR. Isolation and identification of bacteria

- from clinical specimens. In: *Diagnostic procedures in veterinary bacteriology and mycology*, 4th ed., Charles C. Thomas, USA, 1984; pp. 19-30.
- [7] Carrillo PG, Mardaraz C, Pitta-Alvarez SI, Giuliett AM. Isolation and selection of biosurfactant producing bacteria. *World J Microbial Biotechnol* 1996; 12: 62-4.
- [8] Balows A, Hausler WJ, Kenthl JR, Isenberg HHD, Shadomy HJ. *Manual of clinical microbiology* 5th ed. AMS, 1991; pp. 222- 440.
- [9] Morikawa M, Hirata Y, Imanaka TA. Study on the structure-function relationship of the lipopeptide biosurfactants. *Biochim Biophys Acta* 2000; 1488: 211-8.
- [10] Sarubbo LA. Production and stability studies of the bioemulsifier obtained from a strain of *Candida glabrata* UCP 1002. *J Biotechnol* 2006; 9: 400-6.
- [11] Tabatabaee A., Assadi MM, Noohi AA, Sajadian VA. Isolation of biosurfactant producing bacteria from oil reservoirs. *Iranian J Env Health Sci Eng* 2005; 2: 6-12.

